



## DAFTAR PUSTAKA

1. Lakshminarayanan B, Davenport M. Biliary atresia: a comprehensive review. *J Autoimmun.* 2016;73:1–9.
2. Agün M, Tümgör G, Alkan M, et al. Clues to the diagnosis of biliary atresia in neonatal cholestasis. *Turk J Gastroenterol.* 2016;27:37–41.
3. Gunadi, Gunawan TA, Widiyanto G, Yuanita A, Mulyani NS, Makhmudi A. Liver transplant score for prediction of biliary atresia patients' survival following Kasai procedure. *BMC Res Notes.* 2018;11:381.
4. Chiu CY, Chen PH, Chan CF, et al. Biliary atresia in preterm infants in Taiwan: a nationwide survey. *J Pediatr.* 2013;163:100–3.
5. Livesey E, Cortina Borja M, Sharif K, et al. Epidemiology of biliary atresia in England and Wales (1999–2006). *Arch Dis Child Fetal Neonatal Ed.* 2009;94: F451–5.
6. Muthukanagarajan SJ, Karnan I, Srinivasan P, et al. Diagnostic and prognostic significance of various histopathological features in extrahepatic biliary atresia. *J Clin Diagn Res.* 2016;10:EC23–7.
7. Czubkowski P, Cielecka-Kuszyk J, Rurarz M, et al. The limited prognostic value of liver histology in children with biliary atresia. *Ann Hepatol.* 2015;14: 902–9.
8. Gupta L, Gupta SD, Bhatnagar V. Extrahepatic biliary atresia: correlation of histopathology and liver function tests with surgical outcomes. *J Indian Assoc Pediatr Surg.* 2012;17:147–52.
9. Rashed YK, Saber MA, Tawfik M, Mourad WS. Histopathological features and accuracy for diagnosis biliary atresia by prelaparotomy liver biopsy in Egypt. *Egypt Pediatr Assoc Gaz.* 2013;61:42–5.
10. Santos JL, Kieling CO, Meurer L, et al. The extent of biliary proliferation in liver biopsies from patients with biliary atresia at portoenterostomy is associated with the postoperative prognosis. *J Pediatr Surg.* 2009;44: 695–701.
11. Kahn E. Biliary atresia revisited. *Pediatr Dev Pathol.* 2004;7:109–24.
12. Morotti RA, Jain D. Pediatric cholestatic disorders. Approach to pathologic diagnosis. *Surg Pathol.* 2013;6:205–25.
13. Davenport M, Tizzard SA, Underhill J, et al. The biliary atresia splenic malformation syndrome: a 28-year single-center retrospective study. *J Pediatr.* 2006;149:393–400.
14. Rastogi A, Krishnani N, Yachha SK, et al. Histopathological features and accuracy for diagnosing biliary atresia by prelaparotomy liver biopsy in developing countries. *J Gastroenterol Hepatol.* 2009;24:97–102.
15. Azarow KS, Phillips MJ, Sandler AD, et al. Biliary atresia: should all patients undergo a portoenterostomy? *J Pediatr Surg.* 1997;32:168–72.



16. Yeh MM. Pathologic diagnosis of biliary atresia on liver biopsy: is tissue the issue? *J Gastroenterol Hepatol.* 2009;24:933–40.
17. Russo P, Magee JC, Anders RA, Bove KE, Chung C, Cummings OW, et al. Key histopathologic features of liver biopsies that distinguish biliary atresia from other causes of infantile cholestasis and their correlation with outcome: a multicenter study. *Am J Surg Pathol.* 2016;40:1601–15.
18. Shneider BL, Magee JC, Karpen SJ, Rand EB, Narkewicz MR, Bass LM, et al. Total serum bilirubin within 3 months of hepatoperoenterostomy predicts short-term outcomes in biliary atresia. *J Pediatr.* 2016;170:211–7.
19. Serinet MO, Wildhaber BE, Broué P, Lachaux A, Sarles J, Jacquemin E, et al. Impact of age at Kasai operation on its results in late childhood and adolescence: a rational basis for biliary atresia screening. *Pediatrics.* 2009;123:1280–6.
20. Nio M, Ohi R, Miyano T, Saeki M, Shiraki K, Tanaka K, et al. Five- and 10-year survival rates after surgery for biliary atresia: a report from the Japanese biliary atresia registry. *J Pediatr Surg.* 2003;38:997–1000.
21. Wong KK, Chung PH, Chan IH, Lan LC, Tam PK. Performing Kasai portoenterostomy beyond 60 days of life is not necessarily associated with a worse outcome. *J Pediatr Gastroenterol Nutr.* 2010;51:631–4.
22. Schoen BT, Lee H, Sullivan K, Ricketts RR. The Kasai portoenterostomy: when is it too late? *J Pediatr Surg.* 2001;36:97–9.
23. Ramos-Gonzalez G, Elisofon S, Dee EC, Staffa SJ, Medford S, Lillehei C, et al. Predictors of need for liver transplantation in children undergoing hepatoperoenterostomy for biliary atresia. *J Pediatr Surg.* 2019;54: