

DAFTAR PUSTAKA

- Armstrong, R. A. and Mouton, R. (2018) 'Definitions of anaesthetic technique and the implications for clinical research', *Anaesthesia*. Blackwell Publishing Ltd, pp. 935–940. doi: 10.1111/anae.14200.
- Basbaum, A. I., Bautista, D. M., Scherrer, G. and Julius, D. (2009) 'Cellular and Molecular Mechanisms of Pain', *Cell*. Cell, pp. 267–284. doi: 10.1016/j.cell.2009.09.028.
- Bokoch, M. P. and Eilers, H. (2017) 'Intravenous Anesthetics', *Basic of Anesthesia*. Elsevier, pp. 104-106.
- Borendal Wodlin, N., Nilsson, L., Carlsson, P. and Kjølhede, P. (2011) 'Cost-effectiveness of general anesthesia vs spinal anesthesia in fast-track abdominal benign hysterectomy', *American Journal of Obstetrics and Gynecology*, 205(4), pp. 326.e1-326.e7. doi: 10.1016/j.ajog.2011.05.043.
- Braz, L. G., Módolo, N. S. P., do Nascimento, P., Bruschi, B. A. M., Castiglia, Y. M. M., Ganem, E. M., de Carvalho, L. R. and Braz, J. R. C. (2006) 'Perioperative cardiac arrest: A study of 53 718 anaesthetics over 9 yr from a Brazilian teaching hospital', *British Journal of Anaesthesia*, 96(5), pp. 569–575. doi: 10.1093/bja/ael065.
- Brown, E. N., Purdon, P. L. and Van Dort, C. J. (2011) 'General anesthesia and altered states of arousal: A systems neuroscience analysis', *Annual Review of Neuroscience*, 34, pp. 601–628. doi: 10.1146/annurev-neuro-060909-153200.
- Brown, I., Jellish, W. S., Kleinman, B., Fluder, E., Sawicki, K., Katsaros, J. and Rahman, R. (2008) 'Use of postanesthesia discharge criteria to reduce discharge delays for inpatients in the postanesthesia care unit', *Journal of Clinical Anesthesia*, 20(3), pp. 175–179. doi: 10.1016/j.jclinane.2007.09.014.
- Buchanan, F. F., Myles, P. S. and Cicuttini, F. (2011) 'Effect of patient sex on general anaesthesia and recovery', *British Journal of Anaesthesia*, 106(6), pp. 832–839. doi: 10.1093/bja/aer094.
- Corey, J. M., Bulka, C. M. and Ehrenfeld, J. M. (2014) 'Is regional anesthesia associated with reduced PACU length of stay? A retrospective analysis from a tertiary medical center', in *Clinical Orthopaedics and Related Research*. Springer New York LLC, pp. 1427–1433. doi: 10.1007/s11999-013-3336-5.
- Davenport, DL., Xenos, ES., Hosokawa, P., Radford, J., Henderson, EG. and Endean, ED. (2009) 'The influence of body mass index obesity status on vascular surgery 30-day morbidity and mortality', *Journal of vascular surgery*, 49(1). doi: 10.1016/J.JVS.2008.08.052.
- Devereaux, P. J., Xavier, D., Pogue, J., Guyatt, G., Sigamani, A., Garutti, I., Leslie, K., Rao-Melacini, P., Chrolavicius, S., Yang, H., MacDonald, C., Avezum,

- A., Lanthier, L., Hu, W. and Yusuf, S. (2011) 'Characteristics and short-term prognosis of perioperative myocardial infarction in patients undergoing noncardiac surgery cohort study', *Annals of Internal Medicine*, 154(8), pp. 523–528. doi: 10.7326/0003-4819-154-8-201104190-00003.
- Eroglu, A., Apan, A., Erturk, E. and Ben-Shlomo, I. (2015) 'Comparison of the anesthetic techniques', *Scientific World Journal*. Hindawi Publishing Corporation. doi: 10.1155/2015/650684.
- Gabriel, R. A., Waterman, R. S., Kim, J. and Ohno-Machado, L. (2017) 'A predictive model for extended postanesthesia care unit length of stay in outpatient surgeries', *Anesthesia and Analgesia*, 124(5), pp. 1529–1536. doi: 10.1213/ANE.0000000000001827.
- Forman, S. A. and Benkowitz, C. (2018) 'Inhalation Anesthetics', *Anesthesiology*, McGraw-Hills Education, pp. 551-553.
- Gabriel, R. A., Waterman, R. S., Kim, J. and Ohno-Machado, L. (2017) 'A predictive model for extended postanesthesia care unit length of stay in outpatient surgeries', *Anesthesia and Analgesia*, 124(5), pp. 1529–1536. doi: 10.1213/ANE.0000000000001827.
- Gabriel, R. A., Wu, A., Huang, C. C., Dutton, R. P. and Urman, R. D. (2016) 'National incidences and predictors of inefficiencies in perioperative care', *Journal of Clinical Anesthesia*, 31, pp. 238–246. doi: 10.1016/j.jclinane.2016.01.007.
- Ganter, M. T., Blumenthal, S., Dübendorfer, S., Brunnschweiler, S., Hofer, T., Klaghofer, R., Zollinger, A. and Hofer, C. K. (2014) 'The length of stay in the post-anaesthesia care unit correlates with pain intensity, nausea and vomiting on arrival', *Perioperative Medicine*, 3(1). doi: 10.1186/s13741-014-0010-8.
- Gerlach, R. M. and Sweitzer, B. J. (2018) 'Overview of Preoperative Assessment and Management', *Anesthesiology*, McGraw-Hills Education, pp. 35-38.
- Gonano, C., Kettner, S. C., Ernstbrunner, M., Schebesta, K., Chiari, A. and Marhofer, P. (2009) 'Comparison of economical aspects of interscalene brachial plexus blockade and general anaesthesia for arthroscopic shoulder surgery', *British Journal of Anaesthesia*, 103(3), pp. 428–433. doi: 10.1093/bja/aep173.
- Greco, L., Bittner, E. A., George, E. (2018) 'Recovery of Healthy Patient', *Anesthesiology*, McGraw-Hills Education, pp. 1187-1197.
- Hadzic, A., Kerimoglu, B., Loreio, D., Karaca, P. E., Claudio, R. E., Yufa, M., Wedderburn, R., Santos, A. C. and Thys, D. M. (2006) 'Paravertebral blocks provide superior same-day recovery over general anesthesia for patients undergoing inguinal hernia repair', *Anesthesia and Analgesia*, 102(4), pp. 1076–1081. doi: 10.1213/01.ane.0000196532.56221.f2.
- Hadzic, A., Williams, B. A., Karaca, P. E., Hobeika, P., Unis, G., Dermksian, J., Yufa, M., Thys, D. M. and Santos, A. C. (2005) 'For outpatient rotator cuff surgery, nerve block anesthesia provides superior same-day recovery over

- general anesthesia', *Anesthesiology*, 102(5), pp. 1001–1007. doi: 10.1097/0000542-200505000-00020.
- Harris, M. and Chung, F. (2013) 'Complications of general anesthesia', *Clinics in Plastic Surgery*. Clin Plast Surg, pp. 503–513. doi: 10.1016/j.cps.2013.07.001.
- Hausman, M. S., Jewell, E. S. and Engoren, M. (2015) 'Regional versus general anesthesia in surgical patients with chronic obstructive pulmonary disease: Does avoiding general anesthesia reduce the risk of postoperative complications?', *Anesthesia and Analgesia*, 120(6), pp. 1405–1412. doi: 10.1213/ANE.0000000000000574.
- Horlocker, T. T. (2011) 'Complications of regional anesthesia and acute pain management', *Anesthesiology Clinics*. Anesthesiol Clin, pp. 257–278. doi: 10.1016/j.anclin.2011.04.006.
- Irita, K. (2011) 'Risk and crisis management in intraoperative hemorrhage: Human factors in hemorrhagic critical events', *Korean Journal of Anesthesiology*. Korean Society of Anesthesiologists, pp. 151–160. doi: 10.4097/kjae.2011.60.3.151.
- Kadry, B., Press, C. D., Alish, H., Opper, I. M., Orsini, J., Popov, I. A., Brodsky, J. B. and Macario, A. (2014) 'Obesity increases operating roomtimes in patients undergoing primary hip arthroplasty: A retrospective cohort analysis', *PeerJ*, 2014(1). doi: 10.7717/peerj.530.
- Kehlet, H. (2005) 'Postoperative opioid sparing to hasten recovery: What are the issues?', *Anesthesiology*. Anesthesiology, pp. 1083–1085. doi: 10.1097/0000542-200506000-00004.
- Kehlet, H. and White, P. F. (2001) 'Optimizing anesthesia for inguinal herniorrhaphy: General, regional, or local anesthesia?', *Anesthesia and Analgesia*. Lippincott Williams and Wilkins, pp. 1367–1369. doi: 10.1097/0000539-200112000-00001.
- Kilic, E. T. and Naderi, S. (2019) 'Effects of anesthesia protocol on perioperative outcomes and costs of Lumbar Microdiscectomies', *Turkish Neurosurgery*, 29(6), pp. 843–850. doi: 10.5137/1019-5149.JTN.25737-18.4.
- Lang, L. H., Parekh, K., Tsui, B. Y. K. and Maze, M. (2017) 'Perioperative management of the obese surgical patient', *British Medical Bulletin*. Oxford University Press, pp. 135–155. doi: 10.1093/bmb/ldx041.
- Lau, L., Hung, C., Chan, C., Chow, B., Chui, P., Ho, B., Kung, M., Lui, J., Hui, T., Ho, E., Chan, S. and So, H. (2001) *Anaesthetic clinical indicators in public hospitals providing anaesthetic care in Hong Kong: prospective study*.
- Leslie, K., Myles, P. S., Forbes, A., Chan, M. T. V., Short, T. G. and Swallow, S. K. (2005) 'Recovery from bispectral index-guided anaesthesia in a large randomized controlled trial of patients at high risk of awareness', *Anaesthesia and Intensive Care*, 33(4), pp. 443–451. doi: 10.1177/0310057x0503300404.
- Leslie, K., Myles, P. S., Forbes, A., Chan, M. T. V., Short, T. G. and Swallow, S. K. (2005) 'Recovery from bispectral index-guided anaesthesia in a large

- randomized controlled trial of patients at high risk of awareness', *Anaesthesia and Intensive Care*, 33(4), pp. 443–451. doi: 10.1177/0310057x0503300404.
- Li, J., Lam, D., King, H., Credaroli, E., Harmon, E. and Vadivelu, N. (2019) 'Novel Regional Anesthesia for Outpatient Surgery', *Current Pain and Headache Reports*. Current Medicine Group LLC 1. doi: 10.1007/s11916-019-0809-6.
- Lunn, T. H., Kristensen, B. B., Gaarn-Larsen, L., Husted, H. and Kehlet, H. (2012) 'Post-anaesthesia care unit stay after total hip and knee arthroplasty under spinal anaesthesia', *Acta Anaesthesiologica Scandinavica*, 56(9), pp. 1139–1145. doi: 10.1111/j.1399-6576.2012.02709.x.
- Martin, D., Mantziari, S., Demartines, N., Hübner, M., Bismuth, H., Zaninotto, G., *et al.* (2020) 'Defining Major Surgery: A Delphi Consensus Among European Surgical Association (ESA) Members', *World Journal of Surgery*, 44(7), pp. 2211–2219. doi: 10.1007/s00268-020-05476-4.
- McKay, R. E. (2017) 'Inhaled Anesthetics', *Basic of Anesthesia*, Elsevier, pp. 83–86.
- Misal, U., Joshi, S. and Shaikh, M. (2016) 'Delayed recovery from anesthesia: A postgraduate educational review', *Anesthesia: Essays and Researches*, 10(2), p. 164. doi: 10.4103/0259-1162.165506.
- Moore, J. G., Ross, S. M. and Williams, B. A. (2013) 'Regional anesthesia and ambulatory surgery', *Current Opinion in Anaesthesiology*. *Curr Opin Anaesthesiol*, pp. 652–660. doi: 10.1097/ACO.0000000000000011.
- Neuman, M. D., Silber, J. H., Elkassabany, N. M., Ludwig, J. M. and Fleisher, L. A. (2012) 'Comparative effectiveness of regional versus general anesthesia for hip fracture surgery in adults', *Anesthesiology*, 117(1), pp. 72–92. doi: 10.1097/ALN.0b013e3182545e7c.
- Nizamuddin, J., O'Connor, M. (2019) 'Anesthesia for Surgery Patients', *Schwartz's Principles of Surgery*, McGraw-Hills Education, pp. 2036–2039.
- Pain terms: a list with definitions and notes on usage. Recommended by the IASP Subcommittee on Taxonomy. (1979). *Pain*, 6(3), 249.
- Puruhito and Wirjoatmodjo, K. (1997) 'Anestesi', *Buku-Ajar Ilmu Bedah*, Penerbit Buku Kedokteran EGC, pp. 300–334.
- Raja, S. N., Carr, D. B., Cohen, M., Finnerup, N. B., Flor, H., Gibson, S., Keefe, F. J., Mogil, J. S., Ringkamp, M., Sluka, K. A., Song, X. J., Stevens, B., Sullivan, M. D., Tutelman, P. R., Ushida, T. and Vader, K. (2020) 'The revised International Association for the Study of Pain definition of pain: concepts, challenges, and compromises', *Pain*, 161(9), pp. 1976–1982. doi: 10.1097/j.pain.0000000000001939.
- Rathmell, J. P., Wu, C. L., Sinatra, R. S., Ballantyne, J. C., Ginsberg, B., Gordon, D. B., Liu, S. S., Perkins, F. M., Reuben, S. S., Rosenquist, R. W. and Viscusi, E. R. (2006) 'Acute Post-Surgical Pain Management: A Critical Appraisal of Current Practice', *Regional Anesthesia and Pain Medicine*, 31(4 SUPPL.), pp. 1–42. doi: 10.1016/j.rapm.2006.05.002.

- Sastroasmoro, S., and Ismael, S. (2011) 'Studi Kohort', *Dasar-dasar Metodologi Penelitian Klinis*, CV Sagung Seto, pp. 167–179.
- Seago, J. A., Weitz, S. and Walczak, S. (1998) 'Factors influencing stay in the postanesthesia care unit: A prospective analysis', *Journal of Clinical Anesthesia*, 10(7), pp. 579–587. doi: 10.1016/S0952-8180(98)00084-1.
- Seago, J. A., Weitz, S. and Walczak, S. (1998) 'Factors influencing stay in the postanesthesia care unit: A prospective analysis', *Journal of Clinical Anesthesia*, 10(7), pp. 579–587. doi: 10.1016/S0952-8180(98)00084-1.
- Smetana, G. W., Lawrence, V. A. and Cornell, J. E. (2006) 'Preoperative pulmonary risk stratification for noncardiothoracic surgery: Systematic review for the American College of Physicians', *Annals of Internal Medicine*. American College of Physicians, pp. 581–595. doi: 10.7326/0003-4819-144-8-200604180-00009.
- Smith, G. and Goldman, J. (2018) *General Anesthesia for Surgeons*, StatPearls Publishing. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/29630251> (Accessed: 17 April 2021).
- Son, Y. (2010) 'Molecular mechanisms of general anesthesia', *Korean Journal of Anesthesiology*. Korean Society of Anesthesiologists, pp. 3–8. doi: 10.4097/kjae.2010.59.1.3.
- Vadivelu, N., Whitney, C. J., Sinatra, R. S. (2009) 'Pain Pathway and Acute Pain Processing', *Acute Pain Management*, Cambridge University Press, pp. 3-20.
- Valchanov, K. P. and Arrowsmith, J. E. (2012) 'The role of venodilators in the perioperative management of heart failure', *European Journal of Anaesthesiology*. Eur J Anaesthesiol, pp. 121–128. doi: 10.1097/EJA.0b013e32835016be.
- Warren, D. E., Nelson, K., Neal. J. M. (2018) 'Regional Anesthesia', *Anesthesiology*, McGraw-Hills Education, pp. 727-739.
- White, P. F. (2001) 'Optimizing Anesthesia for Inguinal Herniorrhaphy: General, Regional, or Local Anesthesia?', *Anesthesia & Analgesia*, 93(6), pp. 1367–1369. doi: 10.1097/00000539-200112000-00001.
- Zhong, S., Zhong, Xiaoxue, Zhong, Xiaomei and Liu, Y. (2019) 'Comparison between the effect of epidural anesthesia combined with epidural analgesia and general anesthesia combined with intravenous analgesia on prognosis of ovarian cancer patients', *Oncology Letters*, 17(6), pp. 5662–5668. doi: 10.3892/ol.2019.10216.