

Daftar Pustaka

- Abbasi, S., Gharaghani, S., Benvidi, A., & Rezaeinasab, M. (2018). New insights into the efficiency of thymol synergistic effect with p-cymene in inhibiting advanced glycation end products: A multi-way analysis based on spectroscopic and electrochemical methods in combination with molecular docking study. *J Pharm Biomed Anal*, *150*, 436–451. <https://doi.org/10.1016/j.jpba.2017.12.042>
- Adisakwattana, S. (2017). Cinnamic acid and its derivatives: Mechanisms for prevention and management of diabetes and its complications. *Nutrients*, *9*(2). <https://doi.org/10.3390/nu9020163>
- Adisakwattana, S., Sompong, W., Meeprom, A., Ngamukote, S., & Yibchok-Anun, S. (2012). Cinnamic acid and its derivatives inhibit fructose-mediated protein glycation. *Int. J. Mol. Sci.*, *13*(2), 1778–1789. <https://doi.org/10.3390/ijms13021778>
- Adnette, F. N., Ulbad, T. P., Magloire, N., Ruffine, F., Koutinhoun, & Akadiri, Y. (2019). Diabetes Mellitus: Classification, Epidemiology, Physiopathology, Immunology, Risk Factors, Prevention And Nutrition. *Int. J. of Adv Research Journal*, *7*(7), 855–863. <https://doi.org/10.21474/IJAR01/9433>
- Ahmad S, Siddiqui Z, Rehman S, Khan MY, Khan H, Khanum S, *et al.* (2017). A Glycation Angle to Look into the Diabetic Vasculopathy: Cause and Cure. *Curr Vasc Pharmacol*. *15*(4):352-364. doi: 10.2174/1570161115666170327162639. PMID: 28356033.
- Akash, M. S. H., Rehman, K., & Liaqat, A. (2018). Tumor Necrosis Factor-Alpha: Role in Development of Insulin Resistance and Pathogenesis of Type 2 Diabetes Mellitus. *J. Cell. Biochem.*, *119*(1), 105–110. <https://doi.org/10.1002/jcb.26174>
- Alam, M. M., Ahmad, I., & Naseem, I. (2015). Inhibitory effect of quercetin in the formation of advanced glycation end products of human serum albumin: An in vitro and molecular interaction study. *Int. J. Biol. Macromol.*, *79*, 336–343. <https://doi.org/10.1016/j.ijbiomac.2015.05.004>
- Al-Dalaeen AM, AL-Domi HA. (2018). Advanced Glycation End Products: Mechanisms in the Pathogenesis of Type 2 Diabetes and its Complications. *J Res Diabetes Metab*.*4*(1): 016-020
- Allam-Ndoul, B., Guénard, F., Barbier, O., & Vohl, M. C. (2016). Effect of n-3 fatty acids on the expression of inflammatory genes in THP-1 macrophages. *Lipids Health Dis.*, *15*(1), 1–7. <https://doi.org/10.1186/s12944-016-0241-4>
- Allen W, C., Yang, C., Zheleznova, N. N., Staruschenko, A., Kurth, T., Rein, L., *et al.* (2016). Evidence Of The Importance Of Nox4 In Production Of Hypertension In Dahl Salt-Sensitive Rats. *Hypertension*, *67*(2), 440–450. <https://doi.org/10.1183/09031936.00063810>.
- Al-Qattan, K. K., Mansour, M. H., Thomson, M., & Ali, M. (2016). Garlic decreases liver and kidney receptor for advanced glycation end products expression in experimental diabetes. *Pathophysiology*, *23*(2), 135–145. <https://doi.org/10.1016/j.pathophys.2016.02.003>

- American Diabetes Association. (2020). Standards of medical care in diabetes - 2020. *Diabetes Care*, 43(SUPPL. 1). <https://doi.org/10.2337/dc12-s011>
- American Association Diabetes. (2014). Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*, 37(Supplement 1, January 2014), 581. <https://doi.org/10.2337/dc14-S081>
- Anderson, M. M., & Heinecke, J. W. (2003). Production of Nε-(carboxymethyl) Lysine Is Impaired in Mice Deficient in NADPH Oxidase. *Diabetes*, (19), 2137–2143.
- Andreasen, A. S., Kelly, M., Berg, R. M. G., Møller, K., & Pedersen, B. K. (2011). Type 2 diabetes is associated with altered NF-κB DNA binding activity, JNK phosphorylation, and AMPK phosphorylation in skeletal muscle after LPS. *PLoS ONE*, 6(9). <https://doi.org/10.1371/journal.pone.0023999>
- Anguizola, J., Matsuda, R., Barnaby, O. S., Hoy, K. S., Wa, C., DeBolt, E., et al. (2013). Review: Glycation of human serum albumin. *Clin. Chim. Acta*, 425, 64–76. <https://doi.org/10.1016/j.cca.2013.07.013>
- Ansari, N. A., Moinuddin, & Ali, R. (2011). Glycated lysine residues: A marker for non-enzymatic protein glycation in age-related diseases. *Dis. Markers*, 30(6), 317–324. <https://doi.org/10.3233/DMA-2011-0791>
- Arifin, W. N., & Zahiruddin, W. M. (2017). Sample size calculation in animal studies using resource equation approach. *Malaysian J. of Med. Sciences*, 24(5), 101–105. <https://doi.org/10.21315/mjms2017.24.5.11>
- Asgharpour Dil F, Ranjkesh Z, Goodarzi MT. (2019). A systematic review of antiglycation medicinal plants. *Diabetes Metab Syndr*. Mar-Apr;13(2):1225-1229. doi: 10.1016/j.dsx.2019.01.053. Epub 2019 Jan 29. PMID: 31336468.
- Asmat U, Abad K, Ismail K. (2016). Diabetes mellitus and oxidative stress-A concise review. *Saudi Pharm J*. Sep;24(5):547-553. doi: 10.1016/j.jsps.2015.03.013. Epub 2015 Mar 21. PMID: 27752226; PMCID: PMC5059829.
- Awasthi, S., Murugan, N. A., & Saraswathi, N. T. (2015). Advanced glycation end products modulate structure and drug binding properties of albumin. *Mol. Pharmaceutics*, 12(9), 3312–3322. <https://doi.org/10.1021/acs.molpharmaceut.5b00318>
- Bains, Y., & Gugliucci, A. (2017). Ilex paraguariensis and its main component chlorogenic acid inhibit fructose formation of advanced glycation endproducts with amino acids at conditions compatible with those in the digestive system. *Fitoterapia*, 117, 6–10. <https://doi.org/10.1016/j.fitote.2016.12.006>
- Balaji, R., Duraisamy, R., & Santhosh Kumar, M. P. (2019). Complications of diabetes mellitus: A review. *Drug Invention Today*, 12(1), 98–103.
- Baraka-Vidot, J., Navarra, G., Leone, M., Bourdon, E., Militello, V., & Rondeau, P. (2014). Deciphering metal-induced oxidative damages on glycated albumin structure and function. *Biochim. Biophys. Acta, Gen. Subj.*, 1840(6), 1712–1724. <https://doi.org/10.1016/j.bbagen.2013.12.017>
- Bavkar, L. N., Patil, R. S., Rooge, S. B., Nalawade, M. L., & Arvindekar, A. U. (2019). Acceleration of protein glycation by oxidative stress and comparative role of antioxidant and protein glycation inhibitor. *Mol. Cell. Biochem.*, 459(1–2), 61–

71. <https://doi.org/10.1007/s11010-019-03550-7>
- Baynest, H. W. (2015). Classification, pathophysiology, diagnosis and management of diabetes mellitus. *J Diabetes Metab*, *06*(05), 1–10. <https://doi.org/10.4172/2155-6156.1000541>
- Bedard, K., & Krause, K.-H. (2007). The NOX Family of ROS-Generating NADPH Oxidases: Physiology and Pathophysiology. *Physiol. Rev.*, *87*(1), 245–313. <https://doi.org/10.1152/physrev.00044.2005>
- Bhattacharjee, N., Barma, S., Konwar, N., Dewanjee, S., & Manna, P. (2016). Mechanistic insight of diabetic nephropathy and its pharmacotherapeutic targets: An update. *Eur. J. Pharmacol.*, *791*, 8–24. <https://doi.org/10.1016/j.ejphar.2016.08.022>
- Bhonsle, H. S., Korwar, A. M., Kote, S. S., Golegaonkar, S. B., Chougale, A. D., Shaik, M. L., *et al.* (2012). Low plasma albumin levels are associated with increased plasma protein glycation and HbA1c in diabetes. *J Proteome Res*, *11*(2), 1391–1396. <https://doi.org/10.1021/pr201030m>
- Bigagli, E., & Lodovici, M. (2019). Circulating oxidative stress biomarkers in clinical studies on type 2 diabetes and its complications. *Oxid. Med. Cell. Longevity*, *2019*. <https://doi.org/10.1155/2019/5953685>
- Blaser, H., Dostert, C., Mak, T. W., & Brenner, D. (2016). TNF and ROS Crosstalk in Inflammation. *Trends Cell Biol.*, *26*(4), 249–261. <https://doi.org/10.1016/j.tcb.2015.12.002>
- Bohlooli, M., Moosavi-Movahedi, A. A., Taghavi, F., Saboury, A. A., Maghami, P., Seyedarabi, A., *et al.* (2014). Inhibition of fluorescent advanced glycation end products (AGEs) of human serum albumin upon incubation with 3-β-hydroxybutyrate. *Mol. Biol. Rep.*, *41*(6), 3705–3713. <https://doi.org/10.1007/s11033-014-3235-1>
- Bongarzone, S., Savickas, V., Luzi, F., & Gee, A. D. (2017). Targeting the Receptor for Advanced Glycation Endproducts (RAGE): A Medicinal Chemistry Perspective. *J. Med. Chem.*, *60*(17), 7213–7232. <https://doi.org/10.1021/acs.jmedchem.7b00058>
- Brings, S., Fleming, T., Freichel, M., Muckenthaler, M. U., Herzig, S., & Nawroth, P. P. (2017). Dicarboxyls and advanced glycation end-products in the development of diabetic complications and targets for intervention. *Int. J. Mol. Sci.*, *18*(5). <https://doi.org/10.3390/ijms18050984>
- Byun, K., Yoo, Y. C., Son, M., Lee, J., Jeong, G. B., Park, Y. M., *et al.* (2017). Advanced glycation end-products produced systemically and by macrophages: A common contributor to inflammation and degenerative diseases. *Pharmacol. Ther.*, *177*, 44–55. <https://doi.org/10.1016/j.pharmthera.2017.02.030>
- Celikler, S., Tas, S., Vatan, O., Ziyank-Ayvalik, S., Yildiz, G., & Bilaloglu, R. (2009). Anti-hyperglycemic and antigenotoxic potential of *Ulva rigida* ethanolic extract in the experimental diabetes mellitus. *Food Chem. Toxicol.*, *47*(8), 1837–1840. <https://doi.org/10.1016/j.fct.2009.04.039>
- Charan, J., & Biswas, T. (2013). How to calculate sample size for different study

- designs in medical research? *Indian J Psychol Med.*, 35(2), 121–126.
<https://doi.org/10.4103/0253-7176.116232>
- Chen, Y. H., Chen, Z. W., Li, H. M., Yan, X. F., & Feng, B. (2018). AGE/RAGE-induced EMP release via the NOX-derived ROS pathway. *J Diabetes Res*, Mar 18;2018:6823058. <https://doi.org/10.1155/2018/6823058>
- Cho, N. H., Shaw, J. E., Karuranga, S., Huang, Y., da Rocha Fernandes, J. D., Ohlrogge, A. W., *et al.* (2018). IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract*, 138, 271–281. <https://doi.org/10.1016/j.diabres.2018.02.023>
- Chojnacka, K., Saeid, A., Witkowska, Z., & Tuhy, Ł. (2012). Biologically active compounds in seaweed extracts the prospects for the application. *The Open Conference Proceedings Journal*, (3), 20–28. <https://doi.org/10.2174/1876326X01203020020>
- Ciaccio, M. (2019). Introduction of glycated albumin in clinical practice. *J of Lab and Precision Med*, 4, 28–28. <https://doi.org/10.21037/jlpm.2019.08.02>
- Coli, J. C., & Bowden, B. F. (1986). The application of vacuum liquid chromatography to the separation of terpene mixtures. *J. Nat. Prod.*, 49(5), 934–936. <https://doi.org/10.1021/np50047a033>
- Cyriac, B., & Eswaran, K. (2016). Anti- hyperglycemic effect of aqueous extract of *kappaphycus alvarezii* (Doty) doty ex. p. silva in alloxan-induced diabetic rats. *J. Appl. Phycol.*, 28(4), 2507–2513. <https://doi.org/10.1007/s10811-015-0762-7>
- D’Orazio, N., Gammone, M. A., Gemello, E., De Girolamo, M., Cusenza, S., & Riccioni, G. (2012). Marine bioactives: Pharmacological properties and potential applications against inflammatory diseases. *Mar. Drugs*, 10(4), 812–833. <https://doi.org/10.3390/md10040812>
- Daffu, G., del Pozo, C. H., O’Shea, K. M., Ananthakrishnan, R., Ramasamy, R., & Schmidt, A. M. (2013). Radical roles for RAGE in the pathogenesis of oxidative stress in cardiovascular diseases and beyond. *Int. J. Mol. Sci.*, 14(10), 19891–19910. <https://doi.org/10.3390/ijms141019891>
- Dal, S., & Sigrist, S. (2016). The Protective Effect of Antioxidants Consumption on Diabetes and Vascular Complications. *Diseases*, 4(3), 24. <https://doi.org/10.3390/diseases4030024>
- Dahiya, R. S., Kaur, N., Kishore, L., & Gupta, G. K. (2013). Management of diabetic complications : A Chemical constituents based. *J. Ethnopharmacol.*, 150(October 2017), 51–70. <https://doi.org/10.1016/j.jep.2013.08.051>
- Delgado-Andrade, C. (2016). Carboxymethyl-lysine: Thirty years of investigation in the field of AGE formation. *Food Funct*, 7(1), 46–57. <https://doi.org/10.1039/c5fo00918a>
- Desai K, Wu L. (2007). Methylglyoxal and advanced glycation endproducts: new therapeutic horizons? *Recent Pat Cardiovasc Drug Discov*. Jun;2(2):89-99. doi: 10.2174/157489007780832498. PMID: 18221107.
- Di Marzo, N., Chisci, E., & Giovannoni, R. (2018). The role of hydrogen peroxide in redox-dependent signaling: homeostatic and pathological responses in

- mammalian cells. *Cells*, 7(10), 156. <https://doi.org/10.3390/cells7100156>
- Ding, Q., & Keller, J. N. (2005). Evaluation of rage isoforms, ligands, and signaling in the brain. *Biochim. Biophys. Acta - Molecular Cell Research*, 1746(1), 18–27. <https://doi.org/10.1016/j.bbamcr.2005.08.006>
- Dludla, P. V., Joubert, E., Muller, C. J. F., Louw, J., & Johnson, R. (2017). Hyperglycemia-induced oxidative stress and heart disease-cardioprotective effects of rooibos flavonoids and phenylpyruvic acid-2-O-β-D-glucoside. *Nutr. Metab*, 14(1), 1–18. <https://doi.org/10.1186/s12986-017-0200-8>
- Dos Santos, J. M., Tewari, S., & Mendes, R. H. (2019). The role of oxidative stress in the development of diabetes mellitus and its complications. *J Diabetes Res.*, May 5;2019:4189813, 10–13. <https://doi.org/10.1155/2019/4189813>
- Dugasani, S., Pichika, M. R., Nadarajah, V. D., Balijepalli, M. K., Tandra, S., & Korlakunta, J. N. (2010). Comparative antioxidant and anti-inflammatory effects of [6]-gingerol, [8]-gingerol, [10]-gingerol and [6]-shogaol. *J. Ethnopharmacol.*, 127(2), 515–520. <https://doi.org/10.1016/j.jep.2009.10.004>
- Fakhrudin, S., Alanazi, W., & Jackson, K. E. (2017). Diabetes-induced reactive oxygen species: mechanism of their generation and role in renal injury. *J Diabetes Res*, 2017. <https://doi.org/10.1155/2017/8379327>
- Festing, M. F. W., & Altman, D. G. (2002). Guidelines for the design and statistical analysis of experiments using laboratory animals. *ILAR J.*, 43(4), 244–257. <https://doi.org/10.1093/ilar.43.4.244>
- Filip-Ciobotaru, F., Manciu, C., Stoleriu, G., & Foia, L. (2016). NADPH Oxidase: structure and activation mechanisms (review). Note I. *Rev Med Chir Soc Med Nat Iasi*. Jan-Mar;120(1):29-33 Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/27125069>
- Freitas, P. A. C., Ehlert, L. R., & Camargo, J. L. (2017). Glycated albumin: a potential biomarker in diabetes. *Arch Endocrinol Metab*, 61(3), 296–304. <https://doi.org/10.1590/2359-3997000000272>
- Ghasemi, A., Khalifi, S., & Jedi, S. (2014). Streptozotocin-nicotinamide-induced rat model of type 2 diabetes (review). *Acta Physiol Hung.*, 101(4), 408–420. <https://doi.org/10.1556/APhysiol.101.2014.4.2>
- Giacco, F., & Brownlee, M. (2010). Oxidative stress and diabetic complications. *Cir. Res.*, 107(9), 1058–1070. <https://doi.org/10.1161/CIRCRESAHA.110.223545>
- Gilmore, T. D., & Herscovitch, M. (2006). Inhibitors of NF- κB signaling: 785 and counting. *Oncogene*. 2006 Oct 30;25(51):6887-99. doi: 10.1038/sj.onc.1209982. PMID: 17072334.
- Gkogkolou, P., & Böhm, M. (2012). Advanced glycation end products: Keyplayers in skin aging? *Dermatoendocrinol*, 4(3), 259–270. <https://doi.org/10.4161/derm.22028>
- Gokce, G., & Haznedaroglu, M. Z. (2008). Evaluation of antidiabetic, antioxidant and vasoprotective effects of *Posidonia oceanica* extract. *J. Ethnopharmacol.*, 115(1), 122–130. <https://doi.org/10.1016/j.jep.2007.09.016>
- Goldin, A., Beckman, J. A., Schmidt, A. M., & Creager, M. A. (2006a). Advanced

- glycation end products: Sparking the development of diabetic vascular injury. *Circulation*, 114(6), 597–605. <https://doi.org/10.1161/CIRCULATIONAHA.106.621854>
- Gorin, Y., & Block, K. (2013). Nox as a target for diabetic complications. *Clin Sci (Lond)*, 125(8), 361–382. <https://doi.org/10.1042/cs20130065>
- Gorin, Y., & Block, K. (2013). Nox4 and diabetic nephropathy: With a friend like this, who needs enemies? *Free Radical Biol. Med.*, 61(210), 130–142. <https://doi.org/10.1016/j.freeradbiomed.2013.03.014>
- Gugliucci, A., & Teresita, M. (2014). The axis AGE-RAGE-soluble RAGE and oxidative stress in chronic kidney disease, *Adv Exp Med Biol.* 2014;824:191-208. <https://doi.org/10.1007/978-3-319-07320-0>
- Guiry, M.D. & Guiry, G.M. (2020). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M.D. Guiry). *Kappaphycus alvarezii* (Doty) Doty ex P.C.Silva, 1996. Accessed through: World Register of Marine Species at: <http://www.marinespecies.org/aphia.php?p=taxdetails&id=371403> on 2020-11-06
- Gupta, P., Jain, V., Pareek, A., Kumari, P., Singh, R., Agarwal, P., *et al.* (2017). Evaluation of effect of alcoholic extract of heartwood of *Pterocarpus marsupium* on in vitro antioxidant, anti-glycation, sorbitol accumulation and inhibition of aldose reductase activity. *J Tradit Complement Med.*, 7(3), 307–314. <https://doi.org/10.1016/j.jtcme.2016.11.001>
- Gutsol, A., Szyndralewicz, C., Kennedy, C. R. J., Page, P., Burns, K. D., Touyz, R. M., *et al.* (2010). Critical role of Nox4-based NADPH oxidase in glucose-induced oxidative stress in the kidney: implications in type 2 diabetic nephropathy. *Am J Physiol Renal Physiol.*, 299(6), F1348–F1358. <https://doi.org/10.1152/ajprenal.00028.2010>
- Habibi, F., Soufi, F. G., Ghiasi, R., Khamaneh, A. M., & Alipour, M. R. (2016). Alteration in Inflammation-related miR-146a Expression in NF-KB signaling pathway in diabetic rat hippocampus alteration in inflammation-related mir-146a expression in NF-KB signaling pathway in diabetic rat hippocampus. *Adv Pharm Bull*, 6(1), 99–103. <https://doi.org/10.15171/apb.2016.015>
- Hafsa, J., Hammi, K. M., Khedher, M. R. Ben, Smach, M. A., Charfeddine, B., *et al.* (2016). Inhibition of protein glycation, antioxidant and antiproliferative activities of *Carpobrotus edulis* extracts. *Biomed Pharmacother.*, 84, 1496–1503. <https://doi.org/10.1016/j.biopha.2016.11.046>
- Hamasaki, S., Kobori, T., Yamazaki, Y., Kitaura, A., Niwa, A., Nishinaka, T., *et al.* (2018). Effects of scavenger receptors-1 class A stimulation on macrophage morphology and highly modified advanced glycation end product-protein phagocytosis. *Sci. Rep.*, 8(1), 1–6. <https://doi.org/10.1038/s41598-018-24325-y>
- Harnedy, P. A., & FitzGerald, R. J. (2013). In vitro assessment of the cardioprotective, anti-diabetic and antioxidant potential of *Palmaria palmata* protein hydrolysates. *J. Appl. Phycol.*, 25(6), 1793–1803. <https://doi.org/10.1007/s10811-013-0017-4>

- Hayashi, L., & Reis, R. P. (2012). Cultivation of the red algae *Kappaphycus alvarezii* in Brazil and its pharmacological potential. *Brazilian J. Pharmacogn.*, 22(4), 748–752. <https://doi.org/10.1590/S0102-695X2012005000055>
- Heo, S. J., Hwang, J. Y., Choi, J. I., Han, J. S., Kim, H. J., & Jeon, Y. J. (2009). Diploretrohydroxycarmalol isolated from *Ishige okamurae*, a brown algae, a potent α -glucosidase and α -amylase inhibitor, alleviates postprandial hyperglycemia in diabetic mice. *Eur. J. Pharmacol.*, 615(1–3), 252–256. <https://doi.org/10.1016/j.ejphar.2009.05.017>
- Hocine, A., Belmokhtar, K., Bauley, K., Jaisson, S., Gaha, K., Oubaya, N., *et al.* (2015). Serum and tissue accumulation of advanced glycation end-products correlates with vascular changes. *Perit. Dial. Int.*, 35(5), 591–592. <https://doi.org/10.3747/pdi.2014.00118>
- Hong, C. O., Nam, M. H., Oh, J. S., Lee, J. W., Kim, C. T., Park, K. W., *et al.* (2015). Pheophorbide a from *Capsosiphon fulvescens* inhibits advanced glycation end products mediated endothelial dysfunction. *Planta Med.*, 82(1–2), 46–57. <https://doi.org/10.1055/s-0035-1557829>
- Hsia, S. M., Lee, W. H., Yen, G. C., & Wu, C. H. (2016). Capsaicin, an active ingredient from chilli peppers, attenuates glycative stress and restores sRAGE levels in diabetic rats. *J. Funct. Foods*, 21, 406–417. <https://doi.org/10.1016/j.jff.2015.11.043>
- Huang, S. M., Wu, C. S., Chiu, M. H., Yang, H. J., Chen, G. S., & Lan, C. C. E. (2017). High-glucose environment induced intracellular O-GlcNAc glycosylation and reduced galectin-7 expression in keratinocytes: Implications on impaired diabetic wound healing. *J Dermatol Sci*, 87(2), 168–175. <https://doi.org/10.1016/j.jdermsci.2017.04.014>
- Iwai, K. (2008). Antidiabetic and antioxidant effects of polyphenols in brown alga *Ecklonia stolonifera* in genetically diabetic KK-Ay mice. *Plant Foods Hum. Nutr.*, 63(4), 163–169. <https://doi.org/10.1007/s11130-008-0098-4>
- Iwamoto, M., Mizuiri, S., Arita, M., & Hemmi, H. (2005). Nuclear factor- κ B activation in diabetic rat kidney: Evidence for involvement of P-selectin in diabetic nephropathy. *Tohoku J Exp Med.*, 206(2), 163–171. <https://doi.org/10.1620/tjem.206.163>
- Iwasaki, Y., Kambayashi, M., Asai, M., Yoshida, M., Nigawara, T., & Hashimoto, K. (2007). High glucose alone, as well as in combination with proinflammatory cytokines, stimulates nuclear factor kappa-B-mediated transcription in hepatocytes in vitro. *J Diabetes Complications*, 21(1), 56–62. <https://doi.org/10.1016/j.jdiacomp.2006.02.001>
- Jennifer, N., Kiruba, M., Pradeep, M. A., Jemima, S., & Juliana, B. (2015). Study of Phytoconstituents and antibacterial activity of *Kappaphycus alvarezii*. *Int.J.Curr.Microbiol.App.Sci*, 4(5), 1209–1217.
- Justino, A. B., Pereira, M. N., Vilela, D. D., Peixoto, L. G., Martins, M. M., Teixeira, R. R., *et al.* (2016). Peel of araticum fruit (*Annona crassiflora* Mart.) as a source of antioxidant compounds with α -amylase, α -glucosidase and glycation inhibitory

- activities. *Bioorg. Chem.*, *69*, 167–182.
<https://doi.org/10.1016/j.bioorg.2016.11.001>
- Kaewseejan, N., & Siriamornpun, S. (2015). Bioactive components and properties of ethanolic extract and its fractions from *Gynura procumbens* leaves. *Ind. Crops Prod.*, *74*, 271–278. <https://doi.org/10.1016/j.indcrop.2015.05.019>
- Kaku, K. (2010). Pathophysiology of type 2 diabetes and its treatment policy. *Japan Med. Assoc. J.*, *53*(1), 41–46.
- Kang, C., Jin, Y. B., Lee, H., Cha, M., Sohn, E. Tae, Moon, J., *et al.*. (2010). Brown alga *Ecklonia cava* attenuates type 1 diabetes by activating AMPK and Akt signaling pathways. *Food Chem. Toxicol.*, *48*(2), 509–516. <https://doi.org/10.1016/j.fct.2009.11.004>
- Kemendes RI. (2018). Laporan Nasional Riskesdas 2018. *Badan Penelitian Dan Pengembangan Kesehatan*. Retrieved from http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf
- Kerner, W., & Brückel, J. (2014). Definition, classification and diagnosis of diabetes mellitus. *Exp Clin Endocrinol Diabetes*, *122*, 384–386. <https://doi.org/http://dx.doi.org/10.1055/s-0034-1366278>
- Khosravi, F., Kharazmi, F., Kamran, M., Malekzadeh, K., Talebi, A., & Soltani, N. (2018). The role of PPAR- γ and NFkB genes expression in muscle to improve hyperglycemia in STZ-induced diabetic rat following magnesium sulfate administration. *Int J Physiol Pathophysiol Pharmacol.*, *10*(3), 124–131. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/30042813> <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC6055085>
- Kim, K. J., & Lee, B. W. (2012). The roles of glycated albumin as intermediate glycation index and pathogenic protein. *Diabetes Metab J.*, *36*(2), 98–107. <https://doi.org/10.4093/dmj.2012.36.2.98>
- Kim, K. Y., Nam, K. A., Kurihara, H., & Kim, S. M. (2008). Potent α -glucosidase inhibitors purified from the red alga *Grateloupia elliptica*. *Phytochem*, *69*(16), 2820–2825. <https://doi.org/10.1016/j.phytochem.2008.09.007>
- Kim, Y. S., Jung, D. H., Lee, I. S., Pyun, B. J., & Kim, J. S. (2016). *Osteomeles schwerinae* extracts inhibits the binding to receptors of advanced glycation end products and TGF- β 1 expression in mesangial cells under diabetic conditions. *Phytomedicine*, *23*(4), 388–397. <https://doi.org/10.1016/j.phymed.2016.02.005>
- King, G. L., & Loeken, M. R. (2004). Hyperglycemia-induced oxidative stress in diabetic complications. *Histochem Cell Biol*, *122*(4), 333–338. <https://doi.org/10.1007/s00418-004-0678-9>
- Koga, M. (2014). Glycated albumin; clinical usefulness. *Clin. Chim. Acta*, *433*, 96–104. <https://doi.org/10.1016/j.cca.2014.03.001>
- Koo, Y. C., Pyo, M. C., Nam, M. H., Hong, C. O., Yang, S. Y., & Lee, K. W. (2016). Chebulic acid prevents hepatic fibrosis induced by advanced glycation end-products in LX-2 cell by modulating Nrf2 translocation via ERK pathway.

- Toxicol. in Vitro*, 34, 8–15. <https://doi.org/10.1016/j.tiv.2016.03.013>
- Kuda, T., Eda, M., Kataoka, M., Nemoto, M., Kawahara, M., Oshio, S., *et al.*. (2016). Anti-glycation properties of the aqueous extract solutions of dried algae products and effect of lactic acid fermentation on the properties. *Food Chem.*, 192, 1109–1115. <https://doi.org/10.1016/j.foodchem.2015.07.073>
- Kuerban, A., Moselhy, S. S., Yaaser Q. A, Razvi, S.S, Hasan, N.M. Abulnaja, K.O. *et al.* (2017). Natural compounds that inhibit protein glycation: a review for recent findings. *Indo Am. J. Pharm. Sci.*, 4(11), 4027–4042. <https://doi.org/10.5281/ZENODO.1045213>
- Kuo, C. T., Liu, T. H., Hsu, T. H., Lin, F. Y., & Chen, H. Y. (2015). Antioxidant and antiglycation properties of different solvent extracts from Chinese olive (*Canarium album* L.) fruit. *Asian Pac. J. Trop. Med*, 8(12), 1013–1021. <https://doi.org/10.1016/j.apjtm.2015.11.013>
- Lee, H. B., & Blaufox, M. D. (1985). Blood volume in the rat the estimation of blood volume is important in studies. *J Nucl Med*, 25, 72–76.
- Lee, J., Wang, J., & Teo, S. (2017). Antidiabetic effect of *Kappaphycus alvarezii* extracts on streptozotocin-induced Type II Diabetic ICR Mice. *Ann. Pharm. & Pharm. Sci.*, 2(10), 1–5.
- Lee, K. H., & Whang, W. K. (2018). Inhibitory effects of bioassay-guided isolation of anti-glycation components from taraxacum coreanum and simultaneous quantification. *Molecules*, 23(9). <https://doi.org/10.3390/molecules23092148>
- Lee, Y. H., Kown, M. H., Kim, K. J., Lee, E. Y., Kim, D., Lee, B. W., *et al.* (2014). Inverse association between glycated albumin and insulin secretory function may explain higher levels of glycated albumin in subjects with longer duration of diabetes. *PLoS ONE*, 9(9). <https://doi.org/10.1371/journal.pone.0108772>
- Liang, D., Li, F., Fu, Y., Cao, Y., Song, X., Wang, T., *et al.* (2014). Thymol inhibits LPS-stimulated inflammatory response via down-regulation of NF-κB and MAPK signaling pathways in mouse mammary epithelial cells. *Inflammation*, 37(1), 214–222. <https://doi.org/10.1007/s10753-013-9732-x>
- Lin, J. A., Wu, C. H., & Yen, G. C. (2017). Breadfruit flavonoid derivatives attenuate advanced glycation end products (AGEs)-enhanced colon malignancy in HCT116 cancer cells. *J. Funct. Foods*, 31, 248–254. <https://doi.org/10.1016/j.jff.2017.01.050>
- Ling, X., Nagai, R., Sakashita, N., Takeya, M., Takahashi, K., & Horiuchi, S. (2002). Immunohistochemical distribution and quantitative biochemical detection of advanced glycation end products in rats from fetal to adult life. *Int Congr Ser*, 1245(C), 137–142. [https://doi.org/10.1016/S0531-5131\(02\)00961-5](https://doi.org/10.1016/S0531-5131(02)00961-5)
- Liu, J. L, He, Y. L, Wang, S., He, Y., Wang, W. Y Li, Q. J, *et al.*. (2018). Ferulic acid inhibits advanced glycation end products (AGEs) formation and mitigates the AGEs-induced inflammatory response in HUVEC cells. *J. Funct. Foods*, 48(June), 19–26. <https://doi.org/10.1016/j.jff.2018.06.024>
- Liu, T., Zhang, L., Joo, D., & Sun, S. C. (2017). NF-κB signaling in inflammation. *Signal Transduction Targeted Ther.*, 2(April).

- <https://doi.org/10.1038/sigtrans.2017.23>
- Liu, Z., Duan, Y. L., Ge, S. L., Zhang, C. X., Gong, W. H., & Xu, J. J. (2019). Effect of estrogen on right ventricular remodeling of monocrotaline-induced pulmonary arterial hypertension in rats and its mechanism. *Eur. Rev. Med. Pharmacol. Sci.*, 23(4), 1742–1750. https://doi.org/10.26355/eurrev_201902_17136
- Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods*, 25(4), 402–408. <https://doi.org/10.1006/meth.2001.1262>
- Lotfy, M., Adeghate, J., Kalasz, H., Singh, J., & Adeghate, E. (2016). Chronic complications of diabetes mellitus: a mini review. *Curr Diabetes Rev.*, 13(1), 3–10. <https://doi.org/10.2174/1573399812666151016101622>
- Mardiah, Zakaria, F. R., Prangdimurti, E., & Damanik, R. (2015). Anti-inflammatory of purple roselle extract in diabetic rats induced by streptozotocin. *Procedia Food Sci.*, 3(December), 182–189. <https://doi.org/10.1016/j.profoo.2015.01.020>
- Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillaire-Buys, D., *et al.* (1998). Experimental NIDDM: Development of a new model in adult rats administered streptozotocin and nicotinamide. *Diabetes*, 47(2), 224–229. <https://doi.org/10.2337/diab.47.2.224>
- Matough, F. A., Budin, S. B., Hamid, Z. A., Alwahaibi, N., & Mohamed, J. (2012). The role of oxidative stress and antioxidants in diabetic complications, *Sultan Qaboos Univ Med J.* 2012 Feb;12(1):5-18. doi: 10.12816/0003082 PMID: 22375253; PMCID: PMC3286717.
- Matsui, T., Higashimoto, Y., Nishino, Y., Nakamura, N., Fukami, K., & Yamagishi, S. I. (2017). RAGE-aptamer blocks the development and progression of experimental diabetic nephropathy. *Diabetes*, 66(6), 1683–1695. <https://doi.org/10.2337/db16-1281>
- Meenatchi, P., Purushothaman, A., & Maneemegalai, S. (2017). Antioxidant, antiglycation and insulinotrophic properties of *Coccinia grandis* (L.) in vitro: Possible role in prevention of diabetic complications. *J. Tradit. Complement. Med.*, 7(1), 54–64. <https://doi.org/10.1016/j.jtcme.2016.01.002>
- Meng Ji Wu. 1998. Extraction process of pheophorbide A from algae. China No CN1230546A
- Mestry, S. N., Dhodi, J. B., Kumbhar, S. B., & Juvekar, A. R. (2017). Attenuation of diabetic nephropathy in streptozotocin-induced diabetic rats by *Punica granatum* Linn. leaves extract. *J. Tradit. Complement. Med.*, 7(3), 273–280. <https://doi.org/10.1016/j.jtcme.2016.06.008>
- Mildner-Szkudlarz, S., Siger, A., Szwengiel, A., & Bajerska, J. (2015). Natural compounds from grape by-products enhance nutritive value and reduce formation of CML in model muffins. *Food Chem.*, 172, 78–85. <https://doi.org/10.1016/j.foodchem.2014.09.036>
- Moselhy, S. S., Razvi, S. S., Al shibili, F. A., Kuerban, A., Hasan, M. N., Balamash, K. S., *et al.* (2018). m-Coumaric acid attenuates non-catalytic protein glycosylation in the retinas of diabetic rats. *J Pestic Sci.*, 43(3), 180–185.

- <https://doi.org/10.1584/jpestics.D17-091>
- Motawi, T. M. K., Abou-Seif, M. A., Bader, A. M. A., & Mahmoud, M. O. (2013). Effect of glycemic control on soluble RAGE and oxidative stress in type 2 diabetic patients. *BMC Endocr. Disord.*, *13*(1), 1. <https://doi.org/10.1186/1472-6823-13-32>
- Mukai, Y., Shibata, H., Nakamura, T., Yoshioka, Y., Abe, Y., Nomura, T., *et al.* (2009). Structure-function relationship of tumor necrosis factor (TNF) and its receptor interaction based on 3D structural analysis of a fully active TNFR1-selective TNF mutant. *J. Mol. Biol.*, *385*(4), 1221–1229. <https://doi.org/10.1016/j.jmb.2008.11.053>
- Muramatsu, D., Uchiyama, H., Kida, H., & Iwai, A. (2019). Cell cytotoxicity and anti-glycation activity of taxifolin-rich extract from Japanese larch, *Larix kaempferi*. *Heliyon*, *5*(7), e02047. <https://doi.org/10.1016/j.heliyon.2019.e02047>
- Murini, T., Wahyuningsih, M. S. H., Satoto, T. B. T., Fudholi, A., & Hanafi, M. (2018). Isolation and identification of naturally occurring larvicidal compound isolated from zingiber zerumbet (L).J.E. smith. *Asian J. Pharm. Clin. Res.*, *11*(2), 189–193. <https://doi.org/10.22159/ajpcr.2018.v11i2.21703>
- Nagarani, N., & Kumaraguru, A. K. (2013). Evaluation of anti-inflammatory, antidiabetic, cytotoxic activity of *Kappaphycus alvarezii*. *Int J Pharm Bio Sci*, *4*(1)(January 2013), 921–929.
- Negre-Salvayre, A., Salvayre, R., Augé, N., Pamplona, R., & Portero-Otín, M. (2009). Hyperglycemia and glycation in diabetic complications. *Antioxid. Redox Signaling*, *11*(12), 3071–3109. <https://doi.org/10.1089/ars.2009.2484>
- Nevin, C., McNeil, L., Ahmed, N., Murgatroyd, C., Brison, D., & Carroll, M. (2018). Investigating the glycyating effects of glucose, glyoxal and methylglyoxal on human sperm. *Sci. Rep.*, *8*(1), 1–12. <https://doi.org/10.1038/s41598-018-27108-7>
- Niedowics, D. M., & Daleke, D. L (2005). The role of oxidative stress in diabetic complications the role of oxidative stress. *Cell Biochem Biophys*, *43*(October 2015), 289–290. <https://doi.org/10.1385/CBB>
- Nowotny, K., Jung, T., Höhn, A., Weber, D., & Grune, T. (2015). Advanced glycation end products and oxidative stress in type 2 diabetes mellitus. *Biomolecules*, *5*(1), 194–222. <https://doi.org/10.3390/biom5010194>
- Nurilmala, M., Anwar, E., & Luthfiyana, N. (2018). Identification of bioactive compounds of seaweed *Sargassum* sp . and *Euclima cottonii* Doty as a raw sunscreen cream, *Proceedings of the Pakistan Academy of Sciences* *54*(February), 311–318.
- Nwosu, F., Morris, J., Lund, V. A., Stewart, D., Ross, H. A., & McDougall, G. J. (2011). Anti-proliferative and potential anti-diabetic effects of phenolic-rich extracts from edible marine algae. *Food Chem.*, *126*(3), 1006–1012. <https://doi.org/10.1016/j.foodchem.2010.11.111>
- Oliveira-Marques, V., Marinho, H. S., Cyrne, L., & Antunes, F. (2009). Role of hydrogen peroxide in NF-κB activation: From inducer to modulator. *Antioxid Redox Signal*, *11*(9), 2223–2243. <https://doi.org/10.1089/ars.2009.2601>

- Ott, C., Jacobs, K., Haucke, E., Navarrete Santos, A., Grune, T., & Simm, A. (2014). Role of advanced glycation end products in cellular signaling. *Redox Biol.*, 2(1), 411–429. <https://doi.org/10.1016/j.redox.2013.12.016>
- Oyedemi, S. O., Oyedemi, B. O., Coopoosamy, R. M., Prieto, J. M., Stapleton, P., & Gibbons, S. (2017). Antibacterial and norfloxacin potentiation activities of *Ocimum americanum* L. against methicillin resistant *Staphylococcus aureus*. *S. Afr. J. Bot.*, 109, 308–314. <https://doi.org/10.1016/j.sajb.2016.12.025>
- Ozougwu, J. C., Obimba, K. C., Belonwu, C. D., & Unakalamba, C. B. (2013). The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *J. Physiol. Pathophysiol.*, 4(4), 46–57. <https://doi.org/10.5897/jpap2013.0001>
- Papagrigroraki, A., Maurelli, M., Del Giglio, M., Gisondi, P., & Girolomoni, G. (2017). Advanced glycation end products in the pathogenesis of psoriasis. *Int. J. Mol. Sci.*, 18(11), 1–9. <https://doi.org/10.3390/ijms18112471>
- Papatheodorou, K., Banach, M., Bekiari, E., Rizzo, M., & Edmonds, M. (2018). Complications of diabetes 2017. *J Diabetes Res.*, Mar 11, 10–13. <https://doi.org/10.1155/2018/3086167>
- Papatheodorou, K., Papanas, N., Banach, M., Papazoglou, D., & Edmonds, M. (2016). Complications of diabetes 2016. *J Diabetes Res*, 2016(September). <https://doi.org/10.1155/2016/6989453>
- Parameswaran, N., & Patial, S. (2010). Tumor Necrosis Factor- α signalling in macrophages. *Crit Rev Eukaryot Gene Expr.*, 20(2), 87–103. <https://doi.org/10.1016/j.dcn.2011.01.002>
- Park, K. H., Kim, S. J., Kim, J. Y., Min, K. J., Kim, J. R., & Cho, K. H. (2016). Cold-water extract of Korean Red Ginseng exhibits potent inhibitory effects against cholesteryl ester transfer protein activity and fructose-mediated glycation along with lipid-lowering activity in hyperlipidemic zebrafish. *J. Ginseng Res.*, 1–12. <https://doi.org/10.1016/j.jgr.2017.01.005>
- Patel, S., & Santani, D. (2009). Role of NF- κ B in the pathogenesis of diabetes and its associated complications. *Pharmacol. Rep.*, 61(4), 595–603. [https://doi.org/10.1016/S1734-1140\(09\)70111-2](https://doi.org/10.1016/S1734-1140(09)70111-2)
- Peng, X., Ma, J., Chen, F., & Wang, M. (2011). Naturally occurring inhibitors against the formation of advanced glycation end-products. *Food Funct.*, 2(6), 289–301. <https://doi.org/10.1039/c1fo10034c>
- Perez Gutierrez, R. M., & Velazquez, E. G. (2020). Glucopyranoside flavonoids isolated from leaves of *Spinacia oleracea* (spinach) inhibit the formation of advanced glycation end products (AGEs) and aldose reductase activity (RLAR). *Biomed. Pharmacother.*, 128(April), 110299. <https://doi.org/10.1016/j.biopha.2020.110299>
- Perrone, A., Giovino, A., Benny, J., & Martinelli, F. (2020). Advanced Glycation End Products (AGEs): biochemistry, signaling, analytical methods, and epigenetic Effects. *Oxid. Med. Cell. Longevity*, 2020(Cml). <https://doi.org/10.1155/2020/3818196>
- Pia de la Maza, M., Garrido, F., Escalante, N., Leiva, L., Barrera, G., Schnitzler, S., et

- al.* (2012). Fluorescent advanced glycation end-products (ages) detected by spectro-photofluorimetry, as a screening tool to detect diabetic microvascular complications. *J. Diabetes Mellit.*, 02(02), 221–226. <https://doi.org/10.4236/jdm.2012.22035>
- Picot, M. C. N., Bender, O., Atalay, A., Zengin, G., Loffredo, L., Hadji-Minaglou, F., *et al.* (2017). Multiple pharmacological targets, cytotoxicity, and phytochemical profile of *Aphloia theiformis* (Vahl.) Benn. *Biomed Pharmacother.*, 89, 342–350. <https://doi.org/10.1016/j.biopha.2017.02.031>
- Picu, A., Petcu, L., Stefan, S., Mitu, M., Lixandru, D., Ionescu-Tîrgoviste, C., *et al.* (2017). Markers of oxidative stress and antioxidant defense in romanian patients with type 2 diabetes mellitus and obesity. *Molecules*, 22(5), 1–14. <https://doi.org/10.3390/molecules22050714>
- Popov, A. M., & Krivoschapko, O. N. (2013). Protective effects of polar lipids and redox-active compounds from marine organisms at modeling of hyperlipidemia and diabetes, *J. Biomed. Sci. Eng.* 2013(May), 543–550. <https://doi.org/10.4236/jbise.2013.65069>
- Popova, E. A., Mironova, R. S., & Odjakova, M. K. (2010). Non-enzymatic glycosylation and deglycating enzymes. *Biotechnol. Biotechnol. Equip.*, 24(3), 1928–1935. <https://doi.org/10.2478/V10133-010-0066-7>
- Prasad, K., & Mishra, M. (2018). AGE–RAGE stress, stressors, and antistressors in health and disease. *Int J Angiol*, 27(1), 1–12. <https://doi.org/10.1055/s-0037-1613678>
- Prieto-bermejo, R., & Hern, A. (2017). The Importance of NADPH Oxidases and redox signaling in angiogenesis. *Antioxidants*, 6(2), 32. <https://doi.org/10.3390/antiox6020032>
- Qais, F. A., Alam, M. M., Naseem, I., & Ahmad, I. (2016). Understanding the mechanism of non-enzymatic glycation inhibition by cinnamic acid: An in vitro interaction and molecular modelling study. *RSC Advances*, 6(70). <https://doi.org/10.1039/c6ra12321j>
- Quehenberger, P., Bierhaus, A., Fasching, P., Muellner, C., Klevesath, M., Hong, M., *et al.* (2000). Endothelin 1 transcription is controlled by nuclear factor- κ B in AGE-stimulated cultured endothelial cells. *Diabetes*, 49(9), 1561–1570. <https://doi.org/10.2337/diabetes.49.9.1561>
- Radoi, V., Lixandru, D., Mohora, M., & Virgolici, B. (2012). Advanced glycation end products in diabetes mellitus: mechanism of action and focused treatment. *Proc Rom Acad Series B*, 1, 9–19.
- Rahimi-Madiseh, M., Malekpour-Tehrani, A., Bahmani, M., & Rafieian-Kopaei, M. (2016). The research and development on the antioxidants in prevention of diabetic complications. *Asian Pac J Trop Med*, 9(9), 825–831. <https://doi.org/10.1016/j.apjtm.2016.07.001>
- Rajasulochana P., K. P. and D. R. (2012). Potential application of *Kappaphycus alvarezii* in agricultural and pharmaceutical industry. *J Chem Pharm Res*, 4(1), 33–37.

- Ramana, K. V., Friedrich, B., Srivastava, S., Bhatnagar, A., & Srivastava, S. K. (2004). Activation of nuclear factor- κ B by hyperglycemia in vascular smooth muscle cells is regulated by aldose reductase. *Diabetes*, 53(11), 2910–2920. <https://doi.org/10.2337/diabetes.53.11.2910>
- Ramadhan, Y. M. (2018). Pengelolaan budidaya rumput laut berbasis daya dukung lingkungan perairan di pesisir kabupaten Dompu, Provinsi Nusa Tenggara Barat. *Seminar Nasional Geomatika*, 2(February), 01. <https://doi.org/10.24895/SNG.2017.2-0.391>
- Ramkissoon, J. S., Mahomoodally, M. F., Subratty, A. H., & Ahmed, N. (2016). Inhibition of glucose- and fructose-mediated protein glycation by infusions and ethanolic extracts of ten culinary herbs and spices. *Asian Pac J Trop Biomed*, 6(6), 492–500. <https://doi.org/10.1016/j.apjtb.2016.01.016>
- Raupbach, J., Ott, C., Koenig, J., & Grune, T. (2020). Proteasomal degradation of glycated proteins depends on substrate unfolding: Preferred degradation of moderately modified myoglobin. *Free Radical Biology and Medicine*, 152(July 2019), 516–524. <https://doi.org/10.1016/j.freeradbiomed.2019.11.024>
- Rhee, S. Y., & Kim, Y. S. (2018). The role of advanced glycation end products in diabetic vascular complications. *Diabetes Metab J*, 42(3), 188–195. <https://doi.org/10.4093/dmj.2017.0105>
- Rimando, A. M., Olofsdotter, M., Dayan, F. E., & Duke, S. O. (2001). Searching for rice allelochemicals. *Agron. J.*, 93(1), 16–20. <https://doi.org/10.2134/agronj2001.93116x>
- Sadowska-Bartosz, I., & Bartosz, G. (2015). Prevention of protein glycation by natural compounds. *Molecules*, 20(2), 3309–3334. <https://doi.org/10.3390/molecules20023309>
- Saide, A., Lauritano, C., & Ianora, A. (2020). Pheophorbide A: State of the art. *Mar. Drugs*, 18(5), 1–12. <https://doi.org/10.3390/md18050257>
- Salami, M., Rahimmalek, M., & Ehtemam, M. H. (2016). Inhibitory effect of different fennel (*Foeniculum vulgare*) samples and their phenolic compounds on formation of advanced glycation products and comparison of antimicrobial and antioxidant activities. *Food Chem.*, 213, 196–205. <https://doi.org/10.1016/j.foodchem.2016.06.070>
- Sampath, C., Rashid, M. R., Sang, S., & Ahmedna, M. (2017). Green tea epigallocatechin 3-gallate alleviates hyperglycemia and reduces advanced glycation end products via nrf2 pathway in mice with high fat diet-induced obesity. *Biomed. Pharmacother.*, 87, 73–81. <https://doi.org/10.1016/j.biopha.2016.12.082>
- Sanz, A. B., Sanchez-Niño, M. D., Ramos, A. M., Moreno, J. A., Santamaria, B., Ruiz-Ortega, M., et al. (2010). NF- κ B in renal inflammation. *J. Am. Soc. Nephrol.*, 21(8), 1254–1262. <https://doi.org/10.1681/ASN.2010020218>
- Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., & Latha, L. Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants' extracts. *Afr J Tradit Complement Altera Med*, 8(1), 1–10.

- <https://doi.org/10.4314/ajtcam.v8i1.60483>
- Schinzela, R., Münchb, G., Heidlandc, A., & Sebekova, K. (2001). Advanced glycation end products in end-stage renal disease and their removal. *Nephron*, 87(4), 295–303. <https://doi.org/10.1159/000045934>
- Schmidt, A. M., Yan, S. Du, Yan, S. F., & Stern, D. M. (2000). The biology of the receptor for advanced glycation end products and its ligands. *Biochim. Biophys. Acta, Mol. Cell Res.*, 1498(2–3), 99–111. [https://doi.org/10.1016/S0167-4889\(00\)00087-2](https://doi.org/10.1016/S0167-4889(00)00087-2)
- Schröder, K. (2015). NADPH oxidases in bone homeostasis and osteoporosis. *Cell. Mol. Life Sci.*, 72(1), 25–38. <https://doi.org/10.1007/s00018-014-1712-2>
- Selvan, K. B., Piriya, S. P., & Vennison, J. S. (2014). Macro algae (*Euचेuma cottoni* and *Sargassum* Sp.) are reservoirs of biodiesel and bioactive compounds. *J. Chem. Pharm. Sci JCHPS Special Issue*, 2(2), 974–2115.
- Semba, R. D., Nicklett, E. J., & Ferrucci, L. (2010). Does accumulation of advanced glycation end products contribute to the aging phenotype? *J Gerontol A Biol Sci Med Sci*, 65 A(9), 963–975. <https://doi.org/10.1093/gerona/gdq074>
- Serasanambati, M., & Chilakapati, S. R. (2016). Function of Nuclear Factor Kappa B (NF-kB) in human diseases-a review. *South Indian J. Biol. Sci.*, 2(4), 368. <https://doi.org/10.22205/sijbs/2016/v2/i4/103443>
- Serban, A. I., Stanca, L., Geicu, O. I., & Dinischiotu, A. (2015). Ages-induced IL-6 synthesis precedes RAGE up-regulation in HEK 293 cells: An alternative inflammatory mechanism? *Int. J. Mol. Sci.*, 16(9), 20100–20117. <https://doi.org/10.3390/ijms160920100>
- Serdiati, N., & Widiastuti, I. M. (2010). Pertumbuhan dan produksi rumput laut *Euचेuma cottonii* pada kedalaman penanaman yang berbeda. *Media Litbang Sulteng*, III(1), 21–26.
- Shah, S., Baez, E. A., Felipe, D. L., Maynard, J. D., Hempe, J. M., & Chalew, S. A. (2013). Advanced glycation endproducts in children with diabetes. *J. Pediatr.*, 163(5), 1427–1431. <https://doi.org/10.1016/j.jpeds.2013.06.044>
- Shi, D., Guo, S., Jiang, B., Guo, C., Wang, T., Zhang, L., *et al.* (2013). HPN, a synthetic analogue of bromophenol from red alga *Rhodomela confervoides*: Synthesis and anti-diabetic effects in C57BL/KsJ-db/db mice. *Mar. Drugs*, 11(2), 350–362. <https://doi.org/10.3390/md11020350>
- Siddiqui, H., Bashir, M. A., Javaid, K., Nizamani, A., Bano, H., Yousuf, S., *et al.* (2016). Ultrasonic synthesis of tyramine derivatives as novel inhibitors of α -glucosidase in vitro. *J. Enzyme Inhib. Med. Chem.*, 31(6), 1392–1403. <https://doi.org/10.3109/14756366.2016.1142983>
- Sifuentes-Franco, S., Padilla-Tejeda, D. E., Carrillo-Ibarra, S., & Miranda-Díaz, A. G. (2018). Oxidative stress, apoptosis, and mitochondrial function in diabetic nephropathy. *Int. J. Endocrinol.*, 2018. <https://doi.org/10.1155/2018/1875870>
- Singh, V. P., Bali, A., Singh, N., & Jaggi, A. S. (2014a). Advanced glycation end products and diabetic complications. *Korean J Physiol Pharmacol*, 18(1), 1–14. <https://doi.org/10.4196/kjpp.2014.18.1.1>

- Singh, M. P., & Pathak, K. (2015). Animal models for biological screening of anti-diabetic drugs: An overview. *Eur. J. Exp. Biol.*, 5(5), 37–48. Retrieved from <http://www.imedpub.com/articles/animal-models-for-biological-screening-of-antidiabetic-drugs-an-overview.pdf>
- Silván, J. M., Assar, S. H., Srey, C., Dolores Del Castillo, M., & Ames, J. M. (2011). Control of the Maillard reaction by ferulic acid. *Food Chem.*, 128(1), 208–213. <https://doi.org/10.1016/j.foodchem.2011.03.047>
- Siriamornpun, S., Weerapreeyakul, N., & Barusrux, S. (2015). Bioactive compounds and health implications are better for green jujube fruit than for ripe fruit. *J. Funct. Foods*, 12, 246–255. <https://doi.org/10.1016/j.jff.2014.11.016>
- Soenardjo, N. (2011). Aplikasi budidaya rumput laut *Eucheuma cottonii* (Weber van Bosse) dengan Metode Jaring Lepas Dasar (Net Bag) Model Cidaun. *Buletin Oseanografi Marina*, 1(1), 36–44. Retrieved from <https://ejournal.undip.ac.id/index.php/buloma/article/view/2970>
- Stirban, A., Gawlowski, T., & Roden, M. (2014). Vascular effects of advanced glycation endproducts : Clinical effects and molecular mechanisms. *Mol. Metab.*, 3(2), 94–108. <https://doi.org/10.1016/j.molmet.2013.11.006>
- Sudirman, S., Hsu, Y., He, J., & Kong, Z. (2018). Red seaweed *Eucheuma cottonii* ethanol extract regulates inflammatory mediators production and prevents colonic injury on acute colitis disease in mice, (June), 1–11. <https://doi.org/10.20944/preprints201806.0120.v1>
- Suganya, A. M., Sanjivkumar, M., Chandran, M. N., Palavesam, A., & Immanuel, G. (2016). Pharmacological importance of sulphated polysaccharide carrageenan from red seaweed *Kappaphycus alvarezii* in comparison with commercial carrageenan. *Biomed. Pharmacother.*, 84, 1300–1312. <https://doi.org/10.1016/j.biopha.2016.10.067>
- Sumayya, S., & Murugan, K. (2017). Phytochemical screening, RP-HPLC and FTIR analysis of *Kappaphycus alvarezii* (Doty) Doty EX P.C Silva: Macro red algae. *J. Pharmacogn. Phytochem.*, 6(1), 325–330.
- Sun, Q., Li, J., & Gao, F. (2014). New insights into insulin: The anti-inflammatory effect and its clinical relevance. *World Journal of Diabetes*, 5(2), 89. <https://doi.org/10.4239/wjd.v5.i2.89>
- Suryavanshi, S. V., & Kulkarni, Y. A. (2017). NF- κ B: A potential target in the management of vascular complications of diabetes. *Front Pharmacol*, 8(NOV), 1–12. <https://doi.org/10.3389/fphar.2017.00798>
- Swaroop, J. J., Rajarajeswari, D., & Naidu, J. N. (2012). Association of TNF- α with insulin resistance in type 2 diabetes mellitus. *Indian J Med Res*, 135(1), 127–130. <https://doi.org/10.4103/0971-5916.93435>
- Syarif, R. A., Wahyuningsih, M. S. H., Mustofa, M., & Ngatidjan, N. (2018). Antiplasmodial and onset speed of growth inhibitory activities of *Tithonia diversifolia* (Hemsley) A gray leaf fractions against *Plasmodium falciparum*. *Trop J Pharm Res*, 17(11), 2213–2218. <https://doi.org/10.4314/tjpr.v17i11.15>
- Szymczak, I., Śliwińska, A., & Drzewoski, J. (2014). DNA damage and efficacy of

- DNA repair in patients with type 2 diabetes and coexisting colorectal cancer. *Pol Arch Intern Med*, 124(7–8), 352–358. <https://doi.org/10.20452/pamw.2337>
- Takahashi, S., Uchino, H., Shimizu, T., Kanazawa, A., Tamura, Y., Sakai, K., *et al.* (2007). Comparison of Glycated Albumin (GA) and Glycated Hemoglobin (HbA1c) in type 2 diabetic patients: Usefulness of GA for evaluation of short-term changes in glycemic control. *Endocr. J.*, 54(1), 139–144. <https://doi.org/10.1507/endocrj.K06-103>
- Tangko, A. M. (2009). Budidaya Rumput Laut Di Provinsi Sulawesi Selatan.
- Tanji, N., Markowitz, G. S., Fu, C., Kislinger, T., Taguchi, A., Pischetsrieder, M., *et al.* (2000). Expression of advanced glycation end products and their cellular receptor RAGE diabetic nephropathy and nondiabetic renal disease. *J. Am. Soc. Nephrol.*, 11(9), 1656–1666.
- Thornalley, P. J., Häussinger, D., Lüthen, R., Heidland, A., Schinzel, R., Ahmed, N., *et al.* (2005). Increased protein glycation in cirrhosis and therapeutic strategies to prevent it. *Ann. N. Y. Acad. Sci.*, 1043(1), 718–724. <https://doi.org/10.1196/annals.1333.083>
- Tiwari, B. K., Pandey, K. B., Abidi, A. B., & Rizvi, S. I. (2013). Markers of oxidative stress during diabetes mellitus. *J. Biomark.*, 2013, 1–8. <https://doi.org/10.1155/2013/378790>
- Tomobe, Y. I., Morizawa, K., Tsuchida, M., Hibino, H., Nakano, Y., & Tanaka, Y. (2000). Dietary docosahexaenoic acid suppresses inflammation and immunoresponses in contact hypersensitivity reaction in mice. *Lipids*, 35(1), 61–69. <https://doi.org/10.1007/s11745-000-0495-0>
- Trask, O. J., Jr (2012). Nuclear Factor Kappa B (NF-κB) Translocation assay development and validation for high content screening. In S. Markossian (Eds.) *et. al.*, *Assay Guidance Manual*. Eli Lilly & Company and the National Center for Advancing Translational Sciences. PMID: 23035273. P 554
- Tripathi, B. K., & Srivastava, A. K. (2006). Diabetes mellitus: Complications and therapeutics. *Med. Sci. Monit.*, 12(7), RA130–RA147. <https://doi.org/4187> [pii]
- Tupe, R. S., Kulkarni, A., Adeshara, K., Shaikh, S., Shah, N., & Jadhav, A. (2015). *Syzygium jambolanum* and *Cephalandra indica* homeopathic preparations inhibit albumin glycation and protect erythrocytes: An invitro study. *Homeopathy*, 104(3), 197–204. <https://doi.org/10.1016/j.homp.2015.02.009>
- Venskutonis, P. R., Miliauskas, G., & Sivik, B. (2009). Extraction and fractionation of bioactive compounds. *J. Liq. Chromatogr.*, 25, 317. <https://doi.org/10.1186/1471-2105-14-370>
- Vlassara, H., & Uribarri, J. (2014). Advanced Glycation End Products (AGE) and Diabetes: cause, effect, or both? *Curr Diab Rep*, 14(1), 453. <https://doi.org/10.1007/s11892-013-0453-1>.Advanced
- Volpe, C. M. O., Villar-Delfino, P. H., Dos Anjos, P. M. F., & Nogueira-Machado, J. A. (2018). Cellular death, reactive oxygen species (ROS) and diabetic complications review-Article. *Cell Death Dis.*, 9(2). <https://doi.org/10.1038/s41419-017-0135-z>

- Wahyuningsih, M. S. H., Mubarika, S., Ganjar, I. G., Wahyuono, S., & Takeya, T. (2017). 5 α -Oleandrin reduce Bcl-2 protein and increase Bax protein expression on Hela cervical cancer cell. *UnivMed*, 36(2), 102. <https://doi.org/10.18051/univmed.2017.v36.102-109>
- Wahyuningsih, M. S. H., Wijayanti, M. A., Budiyanto, A., & Hanafi, M. (2015). Isolation and identification of potential cytotoxic compound from kembang bulan [*tithonia diversifolia* (Hemsley) a gray] leaves. *Int J Pharm Pharm Sci*, 7(6), 298–301.
- Wang, W., Okada, Y., Shi, H., Wang, Y., & Okuyama, T. (2005). Structures and aldose reductase inhibitory effects of bromophenols from the red alga *Symphyocladia latiuscula*. *J. Nat. Prod.*, 68(4), 620–622. <https://doi.org/10.1021/np040199j>
- Weiss, M. F., Erhard, P., Kader-Attia, F. A., Wu, Y. C., Deoreo, P. B., Araki, A., *et al.* (2000). Mechanisms for the formation of glycoxidation products in end-stage renal disease. *Kidney Int*, 57(6), 2571–2585. <https://doi.org/10.1046/j.1523-1755.2000.00117.x>
- Wu, L., Ma, L., Nicholson, L. F. B., & Black, P. N. (2011). Advanced glycation end products and its receptor (RAGE) are increased in patients with COPD. *Respir. Med.*, 105(3), 329–336. <https://doi.org/10.1016/j.rmed.2010.11.001>
- Xue, J., Rai, V., Frolov, S., Singer, D., Chabierski, S., Xie, J., *et al.* (2012). Advanced glycation end product (AGE) recognition by the receptor for AGEs (RAGE), *Structure* 19(5), 722–732. <https://doi.org/10.1016/j.str.2011.02.013>.
- Yamada, H., Umemoto, T., Kakei, M., Momomura, S. I, Kawakami, M., Ishikawa, S. e., *et al.* (2017). Eicosapentaenoic acid shows anti-inflammatory effect via GPR120 in 3T3-L1 adipocytes and attenuates adipose tissue inflammation in diet-induced obese mice. *Nutr. Metab.*, 14(1), 1–11. <https://doi.org/10.1186/s12986-017-0188-0>
- Yao, D., & Brownlee, M. (2010). Hyperglycemia-induced reactive oxygen species increase expression of the receptor for advanced glycation end products (RAGE) and RAGE ligands. *Diabetes*, 59(1), 249–255. <https://doi.org/10.2337/db09-0801>
- Yazdanpanah, S., Rabiee, M., Tahriri, M., Abdolrahim, M., Rajab, A., Jazayeri, H. E., *et al.* (2017). Evaluation of glycated albumin (GA) and GA/HbA1c ratio for diagnosis of diabetes and glycemic control: A comprehensive review. *Crit. Rev. Clin. Lab. Sci.*, 54(4), 219–232. <https://doi.org/10.1080/10408363.2017.1299684>
- Yang, P., Feng, J., Peng, Q., Liu, X., Fan, Z., & Luca, M. (2019). Advanced Glycation End Products: potential mechanism and therapeutic target in cardiovascular complications under diabetes. *Oxid. Med. Cell. Longev.*, 2019. <https://doi.org/10.1155/2019/9570616>
- Yehualashet, A. S. (2020). Toll-like receptors as a potential drug target for diabetes mellitus and diabetes-associated complications. *Diabetes Metab Syndr Obes.: Targets and Therapy*, 13, 4763–4777. <https://doi.org/10.2147/DMSO.S274844>
- Yoon, H. J., Lee, Y. H., Kim, K. J., Kim, S. R., Kang, E. S., Cha, B. S., *et al.* (2015). Glycated albumin levels in patients with type 2 diabetes increase relative to HbA1c with time. *Biomed Res. Int*, 2015. <https://doi.org/10.1155/2015/576306>

- Yoshiuchi, K., Matsuhisa, M., Katakami, N., Nakatani, Y., Sakamoto, K., Matsuoka, T., Hori, M. (2008). Glycated albumin is a better indicator for glucose excursion than glycated hemoglobin in type 1 and type 2 diabetes. *Endocr. J.*, 55(3), 503–507. <https://doi.org/10.1507/endocrj.K07E-089>
- Younus, H., & Anwar, S. (2016). Prevention of non-enzymatic glycosylation (glycation): Implication in the treatment of diabetic complication. *Int. J. Health Sci.*, 10(2). Retrieved from <http://ijhs.org.sa/index.php/journal/article/view/1310>
- Yu, P., Xu, X. B., & Yu, S. J. (2017). Inhibitory effect of sugarcane molasses extract on the formation of Nε-(carboxymethyl)lysine and Nε-(carboxyethyl)lysine. *Food Chem.*, 221, 1145–1150. <https://doi.org/10.1016/j.foodchem.2016.11.045>
- Yuan, T., Yang, T., Chen, H., Fu, D., Hu, Y., Wang, J., *et al.* (2019). New insights into oxidative stress and inflammation during diabetes mellitus-accelerated atherosclerosis. *Redox Biol.*, 20(August 2018), 247–260. <https://doi.org/10.1016/j.redox.2018.09.025>
- Zhang, M., Otake, K., Miyauchi, Y., Yagi, M., Yonei, Y., Miyakawa, T., *et al.* (2019). Comprehensive NMR analysis of two kinds of post-fermented tea and their anti-glycation activities in vitro. *Food Chem.*, 277(July 2018), 735–743. <https://doi.org/10.1016/j.foodchem.2018.11.028>
- Zhang, Q. W., Lin, L. G., & Ye, W. C. (2018). Techniques for extraction and isolation of natural products: A comprehensive review. *Chin. Med. (United Kingdom)*, 13(1), 1–26. <https://doi.org/10.1186/s13020-018-0177-x>
- Zhao, C., Yang, C., Liu, B., Lin, L., Sarker, S. D., Nahar, L., *et al.* (2018). Bioactive compounds from marine macroalgae and their hypoglycemic benefits. *Trends Food Sci. Technol.*, 72(December), 1–12. <https://doi.org/10.1016/j.tifs.2017.12.001>