UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Daftar Pustaka

Abbasi, S., Gharaghani, S., Benvidi, A., & Rezaeinasab, M. (2018). New insights into the efficiency of thymol synergistic effect with p-cymene in inhibiting advanced glycation end products: A multi-way analysis based on spectroscopic and electrochemical methods in combination with molecular docking study. *J Pharm Biomed Anal*, *150*, 436–451. https://doi.org/10.1016/j.jpba.2017.12.042

Adisakwattana, S. (2017). Cinnamic acid and its derivatives: Mechanisms for prevention and management of diabetes and its complications. *Nutrients*, *9*(2). https://doi.org/10.3390/nu9020163

Adisakwattana, S., Sompong, W., Meeprom, A., Ngamukote, S., & Yibchok-Anun, S. (2012). Cinnamic acid and its derivatives inhibit fructose-mediated protein glycation. *Int. J. Mol. Sci.*, *13*(2), 1778–1789. https://doi.org/10.3390/ijms13021778

Adnette, F. N., Ulbad, T. P., Magloire, N., Ruffine, F., Koutinhouin, & Akadiri, Y. (2019). Diabetes Mellitus: Classification, Epidemiology, Physiopathology, Immunology, Risk Factors, Prevention And Nutrition. *Int. J. of Adv Research Journal*, *7*(7), 855–863. https://doi.org/10.21474/IJAR01/ 9433

Ahmad S, Siddiqui Z, Rehman S, Khan MY, Khan H, Khanum S, *et al.*. (2017). A Glycation Angle to Look into the Diabetic Vasculopathy: Cause and Cure. *Curr Vasc Pharmacol*. 15(4):352-364. doi: 10.2174/1570161115666170327162639. PMID: 28356033.

Akash, M. S. H., Rehman, K., & Liaqat, A. (2018). Tumor Necrosis Factor-Alpha: Role in Development of Insulin Resistance and Pathogenesis of Type 2 Diabetes Mellitus. *J. Cell. Biochem.*, *119*(1), 105–110. https://doi.org/10.1002/jcb.26174

Alam, M. M., Ahmad, I., & Naseem, I. (2015). Inhibitory effect of quercetin in the formation of advanced glycation end products of human serum albumin: An in vitro and molecular interaction study. *Int. J. Biol. Macromol.*, *79*, 336–343. https://doi.org/10.1016/j.ijbiomac.2015.05.004

Al-Dalaeen AM, AL-Domi HA. (2018). Advanced Glycation End Products: Mechanisms in the Pathogenesis of Type 2 Diabetes and its Complications. *J Res Diabetes Metab*.4(1): 016-020

Allam-Ndoul, B., Guénard, F., Barbier, O., & Vohl, M. C. (2016). Effect of n-3 fatty acids on the expression of inflammatory genes in THP-1 macrophages. *Lipids Health Dis.*, *15*(1), 1–7. https://doi.org/10.1186/s12944-016-0241-4

Allen W, C., Yang, C., Zheleznova, N. N., Staruschenko, A., Kurth, T., Rein, L., *et al*. (2016). Evidence Of The Importance Of Nox4 In Production Of Hypertension In Dahl Salt-Sensitive Rats. *Hypertension*, *67*(2), 440–450. https://doi.org/10.1183/09031936.00063810.

Al-Qattan, K. K., Mansour, M. H., Thomson, M., & Ali, M. (2016). Garlic decreases liver and kidney receptor for advanced glycation end products expression in experimental diabetes. *Pathophysiology*, *23*(2), 135–145. https://doi.org/10.1016/j.pathophys.2016.02.003

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

American Diabetes Association. (2020). Standards of medical care in diabetes - 2020. *Diabetes Care*, *43*(SUPPL. 1). https://doi.org/10.2337/dc12-s011

American Association Diabetes. (2014). Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care*, *37*(Supplement 1, January 2014), 581. https://doi.org/10.2337/dc14-S081

Anderson, M. M., & Heinecke, J. W. (2003). Production of Nε-(carboxymethyl) Lysine Is Impaired in Mice Deficient in NADPH Oxidase. *Diabetes*, (19), 2137–2143.

Andreasen, A. S., Kelly, M., Berg, R. M. G., Møller, K., & Pedersen, B. K. (2011). Type 2 diabetes is associated with altered NF-κB DNA binding activity, JNK phosphorylation, and AMPK phosphorylation in skeletal muscle after LPS. *PLoS ONE*, *6*(9). https://doi.org/10.1371/journal.pone.0023999

Anguizola, J., Matsuda, R., Barnaby, O. S., Hoy, K. S., Wa, C., DeBolt, E., *et al.* (2013). Review: Glycation of human serum albumin. *Clin. Chim. Acta*, *425*, 64–76. https://doi.org/10.1016/j.cca.2013.07.013

Ansari, N. A., Moinuddin, & Ali, R. (2011). Glycated lysine residues: A marker for non-enzymatic protein glycation in age-related diseases. *Dis. Markers*, *30*(6), 317–324. https://doi.org/10.3233/DMA-2011-0791

Arifin, W. N., & Zahiruddin, W. M. (2017). Sample size calculation in animal studies using resource equation approach. *Malaysian J. of Med. Sciences*, *24*(5), 101–105. https://doi.org/10.21315/mjms2017.24.5.11

Asgharpour Dil F, Ranjkesh Z, Goodarzi MT. (2019). A systematic review of antiglycation medicinal plants. *Diabetes Metab Syndr*. Mar-Apr;13(2):1225-1229. doi: 10.1016/j.dsx.2019.01.053. Epub 2019 Jan 29. PMID: 31336468.

Asmat U, Abad K, Ismail K. (2016). Diabetes mellitus and oxidative stress-A concise review. *Saudi Pharm J*. Sep;24(5):547-553. doi: 10.1016/j.jsps.2015.03.013. Epub 2015 Mar 21. PMID: 27752226; PMCID: PMC5059829.

Awasthi, S., Murugan, N. A., & Saraswathi, N. T. (2015). Advanced glycation end products modulate structure and drug binding properties of albumin. *Mol. Pharmaceutics*, *12*(9), 3312–3322. https://doi.org/10.1021/acs.molpharmaceut.5b00318

Bains, Y., & Gugliucci, A. (2017). Ilex paraguariensis and its main component chlorogenic acid inhibit fructose formation of advanced glycation endproducts with amino acids at conditions compatible with those in the digestive system. *Fitoterapia*, *117*, 6–10. https://doi.org/10.1016/j.fitote.2016.12.006

Balaji, R., Duraisamy, R., & Santhosh Kumar, M. P. (2019). Complications of diabetes mellitus: A review. *Drug Invention Today*, *12*(1), 98–103.

Baraka-Vidot, J., Navarra, G., Leone, M., Bourdon, E., Militello, V., & Rondeau, P. (2014). Deciphering metal-induced oxidative damages on glycated albumin structure and function. *Biochim. Biophys. Acta, Gen. Subj.*, *1840*(6), 1712–1724. https://doi.org/10.1016/j.bbagen.2013.12.017

Bavkar, L. N., Patil, R. S., Rooge, S. B., Nalawade, M. L., & Arvindekar, A. U. (2019). Acceleration of protein glycation by oxidative stress and comparative role of antioxidant and protein glycation inhibitor. *Mol. Cell. Biochem.*, *459*(1–2), 61–

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

71. https://doi.org/10.1007/s11010-019-03550-7

Baynest, H. W. (2015). Classification, pathophysiology, diagnosis and management of diabetes mellitus. *J Diabetes Metab*, *06*(05), 1–10. https://doi.org/10.4172/2155-6156.1000541

Bedard, K., & Krause, K.-H. (2007). The NOX Family of ROS-Generating NADPH Oxidases: Physiology and Pathophysiology. *Physiol. Rev.*, *87*(1), 245–313. https://doi.org/10.1152/physrev.00044.2005

Bhattacharjee, N., Barma, S., Konwar, N., Dewanjee, S., & Manna, P. (2016). Mechanistic insight of diabetic nephropathy and its pharmacotherapeutic targets: An update. *Eur. J. Pharmacol.*, *791*, 8–24. https://doi.org/10.1016/j.ejphar.2016.08.022

Bhonsle, H. S., Korwar, A. M., Kote, S. S., Golegaonkar, S. B., Chougale, A. D., Shaik, M. L., *et al*. (2012). Low plasma albumin levels are associated with increased plasma protein glycation and HbA1c in diabetes. *J Proteome Res*, *11*(2), 1391–1396. https://doi.org/10.1021/pr201030m

Bigagli, E., & Lodovici, M. (2019). Circulating oxidative stress biomarkers in clinical studies on type 2 diabetes and its complications. *Oxid. Med. Cell. Longevity*, *2019*. https://doi.org/10.1155/2019/5953685

Blaser, H., Dostert, C., Mak, T. W., & Brenner, D. (2016). TNF and ROS Crosstalk in Inflammation. *Trends Cell Biol.*, *26*(4), 249–261. https://doi.org/10.1016/j.tcb.2015.12.002

Bohlooli, M., Moosavi-Movahedi, A. A., Taghavi, F., Saboury, A. A., Maghami, P., Seyedarabi, A., *et al.* (2014). Inhibition of fluorescent advanced glycation end products (AGEs) of human serum albumin upon incubation with 3-β-hydroxybutyrate. *Mol. Biol. Rep.*, *41*(6), 3705–3713. https://doi.org/10.1007/s11033-014-3235-1

Bongarzone, S., Savickas, V., Luzi, F., & Gee, A. D. (2017). Targeting the Receptor for Advanced Glycation Endproducts (RAGE): A Medicinal Chemistry Perspective. *J. Med. Chem.*, *60*(17), 7213–7232. https://doi.org/10.1021/acs.jmedchem.7b00058

Brings, S., Fleming, T., Freichel, M., Muckenthaler, M. U., Herzig, S., & Nawroth, P. P. (2017). Dicarbonyls and advanced glycation end-products in the development of diabetic complications and targets for intervention. *Int. J. Mol. Sci.*, *18*(5). https://doi.org/10.3390/ijms18050984

Byun, K., Yoo, Y. C., Son, M., Lee, J., Jeong, G. B., Park, Y. M., *et al.*. (2017). Advanced glycation end-products produced systemically and by macrophages: A common contributor to inflammation and degenerative diseases. *Pharmacol. Ther.*, *177*, 44–55. https://doi.org/10.1016/j.pharmthera.2017.02.030

Celikler, S., Tas, S., Vatan, O., Ziyanok-Ayvalik, S., Yildiz, G., & Bilaloglu, R. (2009). Anti-hyperglycemic and antigenotoxic potential of Ulva rigida ethanolic extract in the experimental diabetes mellitus. *Food Chem. Toxicol.*, *47*(8), 1837–1840. https://doi.org/10.1016/j.fct.2009.04.039

Charan, J., & Biswas, T. (2013). How to calculate sample size for different study

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

designs in medical research? *Indian J Psychol Med*., *35*(2), 121–126. https://doi.org/10.4103/0253-7176.116232

Chen, Y. H., Chen, Z. W., Li, H. M., Yan, X. F., & Feng, B. (2018). AGE/RAGE-induced EMP release via the NOX-derived ROS pathway. *J Diabetes Res*, Mar 18;2018:6823058. https://doi.org/10.1155/2018/6823058

Cho, N. H., Shaw, J. E., Karuranga, S., Huang, Y., da Rocha Fernandes, J. D., Ohlrogge, A. W., *et al.*. (2018). IDF Diabetes Atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract*, *138*, 271–281. https://doi.org/10.1016/j.diabres.2018.02.023

Chojnacka, K., Saeid, A., Witkowska, Z., & Tuhy, Ł. (2012). Biologically active compounds in seaweed extracts the prospects for the application. *The Open Conference Proceedings Journal*, (3), 20–28. https://doi.org/10.2174/1876326X01203020020

Ciaccio, M. (2019). Introduction of glycated albumin in clinical practice. *J of Lab and Precision Med*, *4*, 28–28. https://doi.org/10.21037/jlpm.2019.08.02

Coli, J. C., & Bowden, B. F. (1986). The application of vacuum liquid chromatography to the separation of terpene mixtures. *J. Nat. Prod.*, *49*(5), 934–936. https://doi.org/10.1021/np50047a033

Cyriac, B., & Eswaran, K. (2016). Anti- hyperglycemic effect of aqueous extract of kappaphycus alvarezii (Doty) doty ex. p. silva in alloxan-induced diabetic rats. *J. Appl. Phycol.*, *28*(4), 2507–2513. https://doi.org/10.1007/s10811-015-0762-7

D'Orazio, N., Gammone, M. A., Gemello, E., De Girolamo, M., Cusenza, S., & Riccioni, G. (2012). Marine bioactives: Pharmacological properties and potential applications against inflammatory diseases. *Mar. Drugs*, *10*(4), 812–833. https://doi.org/10.3390/md10040812

Daffu, G., del Pozo, C. H., O'Shea, K. M., Ananthakrishnan, R., Ramasamy, R., & Schmidt, A. M. (2013). Radical roles for RAGE in the pathogenesis of oxidative stress in cardiovascular diseases and beyond. *Int. J. Mol. Sci.*, *14*(10), 19891–19910. https://doi.org/10.3390/ijms141019891

Dal, S., & Sigrist, S. (2016). The Protective Effect of Antioxidants Consumption on Diabetes and Vascular Complications. *Diseases*, *4*(3), 24. https://doi.org/10.3390/diseases4030024

Dahiya, R. S., Kaur, N., Kishore, L., & Gupta, G. K. (2013). Management of diabetic complications : A Chemical constituents based. *J. Ethnopharmacol.*, *150*(October 2017), 51–70. https://doi.org/10.1016/j.jep.2013.08.051

Delgado-Andrade, C. (2016). Carboxymethyl-lysine: Thirty years of investigation in the field of AGE formation. *Food Funct*, *7*(1), 46–57. https://doi.org/10.1039/c5fo00918a

Desai K, Wu L. (2007). Methylglyoxal and advanced glycation endproducts: new therapeutic horizons? *Recent Pat Cardiovasc Drug Discov*. Jun;2(2):89-99. doi: 10.2174/157489007780832498. PMID: 18221107.

Di Marzo, N., Chisci, E., & Giovannoni, R. (2018). The role of hydrogen peroxide in redox-dependent signaling: homeostatic and pathological responses in

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

mammalian cells. *Cells*, *7*(10), 156. https://doi.org/10.3390/cells7100156

Ding, Q., & Keller, J. N. (2005). Evaluation of rage isoforms, ligands, and signaling in the brain. *Biochim. Biophys. Acta - Molecular Cell Research*, *1746*(1), 18–27. https://doi.org/10.1016/j.bbamcr.2005.08.006

Dludla, P. V., Joubert, E., Muller, C. J. F., Louw, J., & Johnson, R. (2017). Hyperglycemia-induced oxidative stress and heart disease-cardioprotective effects of rooibos flavonoids and phenylpyruvic acid-2-O-β-D-glucoside. *Nutr. Metab*, *14*(1), 1–18. https://doi.org/10.1186/s12986-017-0200-8

Dos Santos, J. M., Tewari, S., & Mendes, R. H. (2019). The role of oxidative stress in the development of diabetes mellitus and its complications. *J Diabetes Res.*, May 5;2019:4189813, 10–13. https://doi.org/10.1155/2019/4189813

Dugasani, S., Pichika, M. R., Nadarajah, V. D., Balijepalli, M. K., Tandra, S., & Korlakunta, J. N. (2010). Comparative antioxidant and anti-inflammatory effects of [6]-gingerol, [8]-gingerol, [10]-gingerol and [6]-shogaol. *J. Ethnopharmacol.*, *127*(2), 515–520. https://doi.org/10.1016/j.jep.2009.10.004

Fakhruddin, S., Alanazi, W., & Jackson, K. E. (2017). Diabetes-induced reactive oxygen species: mechanism of their generation and role in renal injury. *J Diabetes Res*, *2017*. https://doi.org/10.1155/2017/8379327

Festing, M. F. W., & Altman, D. G. (2002). Guidelines for the design and statistical analysis of experiments using laboratory animals. *ILAR J.*, *43*(4), 244–257. https://doi.org/10.1093/ilar.43.4.244

Filip-Ciubotaru, F., Manciuc, C., Stoleriu, G., & Foia, L. (2016). NADPH Oxidase: structure and activation mechanisms (review). Note I. *Rev Med Chir Soc Med Nat Iasi*. Jan-Mar;120(1):29-33 Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/27125069

Freitas, P. A. C., Ehlert, L. R., & Camargo, J. L. (2017). Glycated albumin: a potential biomarker in diabetes. *Arch Endocrinol Metab*, *61*(3), 296–304. https://doi.org/10.1590/2359-3997000000272

Ghasemi, A., Khalifi, S., & Jedi, S. (2014). Streptozotocin-nicotinamide-induced rat model of type 2 diabetes (review). *Acta Physiol Hung.*, *101*(4), 408–420. https://doi.org/10.1556/APhysiol.101.2014.4.2

Giacco, F., & Brownlee, M. (2010). Oxidative stress and diabetic complications. *Cir. Res.*, *107*(9), 1058–1070. https://doi.org/10.1161/CIRCRESAHA.110.223545

Gilmore, T. D., & Herscovitch, M. (2006). Inhibitors of NF- κB signaling : 785 and counting, *Oncogene*. 2006 Oct 30;25(51):6887-99. doi: 10.1038/sj.onc.1209982. PMID: 17072334.

Gkogkolou, P., & Böhm, M. (2012). Advanced glycation end products: Keyplayers in skin aging? *Dermatoendocrinol*, *4*(3), 259–270. https://doi.org/10.4161/derm.22028

Gokce, G., & Haznedaroglu, M. Z. (2008). Evaluation of antidiabetic, antioxidant and vasoprotective effects of Posidonia oceanica extract. *J. Ethnopharmacol.*, *115*(1), 122–130. https://doi.org/10.1016/j.jep.2007.09.016

Goldin, A., Beckman, J. A., Schmidt, A. M., & Creager, M. A. (2006a). Advanced

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

glycation end products: Sparking the development of diabetic vascular injury. *Circulation*, *114*(6), 597–605. https://doi.org/10.1161/CIRCULATIONAHA.106.621854

Gorin, Y., & Block, K. (2013). Nox as a target for diabetic complications. *Clin Sci (Lond)*, *125*(8), 361–382. https://doi.org/10.1042/cs20130065

Gorin, Y., & Block, K. (2013). Nox4 and diabetic nephropathy: With a friend like this, who needs enemies? *Free Radical Biol. Med.*, *61*(210), 130–142. https://doi.org/10.1016/j.freeradbiomed.2013.03.014

Gugliucci, A., & Teresita, M. (2014). The axis AGE-RAGE-soluble RAGE and oxidative stress in chronic kidney disease, *Adv Exp Med Biol*. 2014;824:191-208. https://doi.org/10.1007/978-3-319-07320-0

Guiry, M.D. & Guiry, G.M. (2020). AlgaeBase. World-wide electronic publication, National University of Ireland, Galway (taxonomic information republished from AlgaeBase with permission of M.D. Guiry). *Kappaphycus alvarezii* (Doty) Doty ex P.C.Silva, 1996. Accessed through: World Register of Marine Species at: http://www.marinespecies.org/aphia.php?p=taxdetails&id=371403 on 2020-11-06

Gupta, P., Jain, V., Pareek, A., Kumari, P., Singh, R., Agarwal, P., *et al.* (2017). Evaluation of effect of alcoholic extract of heartwood of Pterocarpus marsupium on in vitro antioxidant, anti-glycation, sorbitol accumulation and inhibition of aldose reductase activity. *J Tradit Complement Med.*, *7*(3), 307–314. https://doi.org/10.1016/j.jtcme.2016.11.001

Gutsol, A., Szyndralewiez, C., Kennedy, C. R. J., Page, P., Burns, K. D., Touyz, R. M., *et al.*. (2010). Critical role of Nox4-based NADPH oxidase in glucose-induced oxidative stress in the kidney: implications in type 2 diabetic nephropathy. *Am J Physiol Renal Physiol.*, *299*(6), F1348–F1358. https://doi.org/10.1152/ajprenal.00028.2010

Habibi, F., Soufi, F. G., Ghiasi, R., Khamaneh, A. M., & Alipour, M. R. (2016). Alteration in Inflammation-related miR-146a Expression in NF-KB signaling pathway in diabetic rat hippocampus alteration in inflammation-related mir-146a expression in NF-KB signaling pathway in diabetic rat hippocampus. *Adv Pharm Bull*, *6*(1), 99–103. https://doi.org/10.15171/apb.2016.015

Hafsa, J., Hammi, K. M., Khedher, M. R. Ben, Smach, M. A., Charfeddine, B., *et al.*. (2016). Inhibition of protein glycation, antioxidant and antiproliferative activities of Carpobrotus edulis extracts. *Biomed Pharmacother.*, *84*, 1496–1503. https://doi.org/10.1016/j.biopha.2016.11.046

Hamasaki, S., Kobori, T., Yamazaki, Y., Kitaura, A., Niwa, A., Nishinaka, T., *et al.* (2018). Effects of scavenger receptors-1 class A stimulation on macrophage morphology and highly modified advanced glycation end product-protein phagocytosis. *Sci. Rep.*, *8*(1), 1–6. https://doi.org/10.1038/s41598-018-24325-y

Harnedy, P. A., & FitzGerald, R. J. (2013). In vitro assessment of the cardioprotective, anti-diabetic and antioxidant potential of Palmaria palmata protein hydrolysates. *J. Appl. Phycol.*, *25*(6), 1793–1803. https://doi.org/10.1007/s10811-013-0017-4

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Hayashi, L., & Reis, R. P. (2012). Cultivation of the red algae Kappaphycus alvarezii in Brazil and its pharmacological potential. *Brazilian J. Pharmacogn.*, *22*(4), 748–752. https://doi.org/10.1590/S0102-695X2012005000055

Heo, S. J., Hwang, J. Y., Choi, J. I., Han, J. S., Kim, H. J., & Jeon, Y. J. (2009). Diphlorethohydroxycarmalol isolated from Ishige okamurae, a brown algae, a potent α-glucosidase and α-amylase inhibitor, alleviates postprandial hyperglycemia in diabetic mice. *Eur. J. Pharmacol.*, *615*(1–3), 252–256. https://doi.org/10.1016/j.ejphar.2009.05.017

Hocine, A., Belmokhtar, K., Bauley, K., Jaisson, S., Gaha, K., Oubaya, N., *et al.*. (2015). Serum and tissue accumulation of advanced glycation end-products correlates with vascular changes. *Perit. Dial. Int.*, *35*(5), 591–592. https://doi.org/10.3747/pdi.2014.00118

Hong, C. O., Nam, M. H., Oh, J. S., Lee, J. W., Kim, C. T., Park, K. W., *et al.*. (2015). Pheophorbide a from Capsosiphon fulvescens inhibits advanced glycation end products mediated endothelial dysfunction. *Planta Med.*, *82*(1–2), 46–57. https://doi.org/10.1055/s-0035-1557829

Hsia, S. M., Lee, W. H., Yen, G. C., & Wu, C. H. (2016). Capsaicin, an active ingredient from chilli peppers, attenuates glycative stress and restores sRAGE levels in diabetic rats. *J. Funct. Foods*, *21*, 406–417. https://doi.org/10.1016/j.jff.2015.11.043

Huang, S. M., Wu, C. S., Chiu, M. H., Yang, H. J., Chen, G. S., & Lan, C. C. E. (2017). High-glucose environment induced intracellular O-GlcNAc glycosylation and reduced galectin-7 expression in keratinocytes: Implications on impaired diabetic wound healing. *J Dermatol Sci*, *87*(2), 168–175. https://doi.org/10.1016/j.jdermsci.2017.04.014

Iwai, K. (2008). Antidiabetic and antioxidant effects of polyphenols in brown alga Ecklonia stolonifera in genetically diabetic KK-Ay mice. *Plant Foods Hum. Nutr.*, *63*(4), 163–169. https://doi.org/10.1007/s11130-008-0098-4

Iwamoto, M., Mizuiri, S., Arita, M., & Hemmi, H. (2005). Nuclear factor-κB activation in diabetic rat kidney: Evidence for involvement of P-selectin in diabetic nephropathy. *Tohoku J Exp Med.*, *206*(2), 163–171. https://doi.org/10.1620/tjem.206.163

Iwasaki, Y., Kambayashi, M., Asai, M., Yoshida, M., Nigawara, T., & Hashimoto, K. (2007). High glucose alone, as well as in combination with proinflammatory cytokines, stimulates nuclear factor kappa-B-mediated transcription in hepatocytes in vitro. *J Diabetes Complications*, *21*(1), 56–62. https://doi.org/10.1016/j.jdiacomp.2006.02.001

Jennifer, N., Kiruba, M., Pradeep, M. A., Jemima, S., & Juliana, B. (2015). Study of Phytoconstituents and antibacterial activity of Kappaphycus alvarezii. *Int.J.Curr.Microbiol.App.Sci*, *4*(5), 1209–1217.

Justino, A. B., Pereira, M. N., Vilela, D. D., Peixoto, L. G., Martins, M. M., Teixeira, R. R., *et al*. (2016). Peel of araticum fruit (Annona crassiflora Mart.) as a source of antioxidant compounds with α-amylase, α-glucosidase and glycation inhibitory

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

activities. *Bioorg. Chem.*, *69*, 167–182. https://doi.org/10.1016/j.bioorg.2016.11.001

Kaewseejan, N., & Siriamornpun, S. (2015). Bioactive components and properties of ethanolic extract and its fractions from Gynura procumbens leaves. *Ind. Crops Prod.*, *74*, 271–278. https://doi.org/10.1016/j.indcrop.2015.05.019

Kaku, K. (2010). Pathophysiology of type 2 diabetes and its treatment policy. *Japan Med. Assoc. J.*, *53*(1), 41–46.

Kang, C., Jin, Y. B., Lee, H., Cha, M., Sohn, E. Tae, Moon, J., *et al.*. (2010). Brown alga Ecklonia cava attenuates type 1 diabetes by activating AMPK and Akt signaling pathways. *Food Chem. Toxicol.*, *48*(2), 509–516. https://doi.org/10.1016/j.fct.2009.11.004

Kemenkes RI. (2018). Laporan Nasional Riskesdas 2018. *Badan Penelitian Dan Pengembangan Kesehatan*. Retrieved from http://labdata.litbang.kemkes.go.id/images/download/laporan/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf

Kerner, W., & Brückel, J. (2014). Definition, classification and diagnosis of diabetes mellitus. *Exp Clin Endocrinol Diabetes*, *122*, 384–386. https://doi.org/http://dx.doi.org/ 10.1055/s-0034-1366278

Khosravi, F., Kharazmi, F., Kamran, M., Malekzadeh, K., Talebi, A., & Soltani, N. (2018). The role of PPAR-γ and NFKB genes expression in muscle to improve hyperglycemia in STZ-induced diabetic rat following magnesium sulfate administration. *Int J Physiol Pathophysiol Pharmacol.*, *10*(3), 124–131. Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/30042813%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC6055085

Kim, K. J., & Lee, B. W. (2012). The roles of glycated albumin as intermediate glycation index and pathogenic protein. *Diabetes Metab J.*, *36*(2), 98–107. https://doi.org/10.4093/dmj.2012.36.2.98

Kim, K. Y., Nam, K. A., Kurihara, H., & Kim, S. M. (2008). Potent α-glucosidase inhibitors purified from the red alga Grateloupia elliptica. *Phytochem*, *69*(16), 2820–2825. https://doi.org/10.1016/j.phytochem.2008.09.007

Kim, Y. S., Jung, D. H., Lee, I. S., Pyun, B. J., & Kim, J. S. (2016). Osteomeles schwerinae extracts inhibits the binding to receptors of advanced glycation end products and TGF-β1 expression in mesangial cells under diabetic conditions. *Phytomedicine*, *23*(4), 388–397. https://doi.org/10.1016/j.phymed.2016.02.005

King, G. L., & Loeken, M. R. (2004). Hyperglycemia-induced oxidative stress in diabetic complications. *Histochem Cell Biol*, *122*(4), 333–338. https://doi.org/10.1007/s00418-004-0678-9

Koga, M. (2014). Glycated albumin; clinical usefulness. *Clin. Chim. Acta*, *433*, 96–104. https://doi.org/10.1016/j.cca.2014.03.001

Koo, Y. C., Pyo, M. C., Nam, M. H., Hong, C. O., Yang, S. Y., & Lee, K. W. (2016). Chebulic acid prevents hepatic fibrosis induced by advanced glycation end-products in LX-2 cell by modulating Nrf2 translocation via ERK pathway.

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Toxicol. in Vitro, *34*, 8–15. https://doi.org/10.1016/j.tiv.2016.03.013

Kuda, T., Eda, M., Kataoka, M., Nemoto, M., Kawahara, M., Oshio, S., *et al.*. (2016). Anti-glycation properties of the aqueous extract solutions of dried algae products and effect of lactic acid fermentation on the properties. *Food Chem.*, *192*, 1109–1115. https://doi.org/10.1016/j.foodchem.2015.07.073

Kuerban, A., Moselhy, S. S., Yaaser Q. A, Razvi, S.S, Hasan, N.M. Abulnaja, K.O. *et al*. (2017). Natural compounds that inhibit protein glycation: a review for recent findings. *Indo Am. J. Pharm. Sci*, *4*(11), 4027–4042. https://doi.org/10.5281/ZENODO.1045213

Kuo, C. T., Liu, T. H., Hsu, T. H., Lin, F. Y., & Chen, H. Y. (2015). Antioxidant and antiglycation properties of different solvent extracts from Chinese olive (Canarium album L.) fruit. *Asian Pac. J. Trop. Med*, *8*(12), 1013–1021. https://doi.org/10.1016/j.apjtm.2015.11.013

Lee, H. B., & Blaufox, M. D. (1985). Blood volume in the rat the estimation of blood volume is important in studies. *J Nucl Med*, *25*, 72–76.

Lee, J., Wang, J., & Teo, S. (2017). Antidiabetic effect of Kappaphycus alvarezii extracts on streptozotocin-induced Type II Diabetic ICR Mice. *Ann. Pharm. & Pharm. Sci.*, *2*(10), 1–5.

Lee, K. H., & Whang, W. K. (2018). Inhibitory effects of bioassay-guided isolation of anti-glycation components from taraxacum coreanum and simultaneous quantification. *Molecules*, *23*(9). https://doi.org/10.3390/molecules23092148

Lee, Y. H., Kown, M. H., Kim, K. J., Lee, E. Y., Kim, D., Lee, B. W., *et al*. (2014). Inverse association between glycated albumin and insulin secretory function may explain higher levels of glycated albumin in subjects with longer duration of diabetes. *PLoS ONE*, *9*(9). https://doi.org/10.1371/journal.pone.0108772

Liang, D., Li, F., Fu, Y., Cao, Y., Song, X., Wang, T., *et al*. (2014). Thymol inhibits LPS-stimulated inflammatory response via down-regulation of NF-κB and MAPK signaling pathways in mouse mammary epithelial cells. *Inflammation*, *37*(1), 214–222. https://doi.org/10.1007/s10753-013-9732-x

Lin, J. A., Wu, C. H., & Yen, G. C. (2017). Breadfruit flavonoid derivatives attenuate advanced glycation end products (AGEs)-enhanced colon malignancy in HCT116 cancer cells. *J. Funct. Foods*, *31*, 248–254. https://doi.org/10.1016/j.jff.2017.01.050

Ling, X., Nagai, R., Sakashita, N., Takeya, M., Takahashi, K., & Horiuchi, S. (2002). Immunohistochemical distribution and quantitative biochemical detection of advanced glycation end products in rats from fetal to adult life. *Int Congr Ser*, *1245*(C), 137–142. https://doi.org/10.1016/S0531-5131(02)00961-5

Liu, J. L, He, Y. L, Wang, S., He, Y., Wang, W. Y Li, Q. J, *et al.*. (2018). Ferulic acid inhibits advanced glycation end products (AGEs) formation and mitigates the AGEs-induced inflammatory response in HUVEC cells. *J. Funct. Foods*, *48*(June), 19–26. https://doi.org/10.1016/j.jff.2018.06.024

Liu, T., Zhang, L., Joo, D., & Sun, S. C. (2017). NF-κB signaling in inflammation. *Signal Transduction Targeted Ther.*, *2*(April).

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

https://doi.org/10.1038/sigtrans.2017.23

Liu, Z., Duan, Y. L., Ge, S. L., Zhang, C. X., Gong, W. H., & Xu, J. J. (2019). Effect of estrogen on right ventricular remodeling of monocrotaline-induced pulmonary arterial hypertension in rats and its mechanism. *Eur. Rev. Med. Pharmacol. Sci.*, *23*(4), 1742–1750. https://doi.org/10.26355/eurrev_201902_17136

Livak, K. J., & Schmittgen, T. D. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the 2-ΔΔCT method. *Methods*, *25*(4), 402–408. https://doi.org/10.1006/meth.2001.1262

Lotfy, M., Adeghate, J., Kalasz, H., Singh, J., & Adeghate, E. (2016). Chronic complications of diabetes mellitus: a mini review. *Curr Diabetes Rev.*, *13*(1), 3–10. https://doi.org/10.2174/1573399812666151016101622

Mardiah, Zakaria, F. R., Prangdimurti, E., & Damanik, R. (2015). Anti-inflammatory of purple roselle extract in diabetic rats induced by streptozotocin. *Procedia Food Sci.*, *3*(December), 182–189. https://doi.org/10.1016/j.profoo.2015.01.020

Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillaire-Buys, D., *et al*. (1998). Experimental NIDDM: Development of a new model in adult rats administered streptozotocin and nicotinamide. *Diabetes*, *47*(2), 224–229. https://doi.org/10.2337/diab.47.2.224

Matough, F. A., Budin, S. B., Hamid, Z. A., Alwahaibi, N., & Mohamed, J. (2012). The role of oxidative stress and antioxidants in diabetic complications, *Sultan Qaboos Univ Med J*. 2012 Feb;12(1):5-18. doi: 10.12816/0003082 PMID: 22375253; PMCID: PMC3286717.

Matsui, T., Higashimoto, Y., Nishino, Y., Nakamura, N., Fukami, K., & Yamagishi, S. I. (2017). RAGE-aptamer blocks the development and progression of experimental diabetic nephropathy. *Diabetes*, *66*(6), 1683–1695. https://doi.org/10.2337/db16-1281

Meenatchi, P., Purushothaman, A., & Maneemegalai, S. (2017). Antioxidant, antiglycation and insulinotrophic properties of Coccinia grandis (L.) in vitro: Possible role in prevention of diabetic complications. *J. Tradit. Complement. Med*, *7*(1), 54–64. https://doi.org/10.1016/j.jtcme.2016.01.002

Meng Ji Wu. 1998. Extraction process of pheophorbide A from algae. China No CN1230546A

Mestry, S. N., Dhodi, J. B., Kumbhar, S. B., & Juvekar, A. R. (2017). Attenuation of diabetic nephropathy in streptozotocin-induced diabetic rats by Punica granatum Linn. leaves extract. *J. Tradit. Complement. Med*, *7*(3), 273–280. https://doi.org/10.1016/j.jtcme.2016.06.008

Mildner-Szkudlarz, S., Siger, A., Szwengiel, A., & Bajerska, J. (2015). Natural compounds from grape by-products enhance nutritive value and reduce formation of CML in model muffins. *Food Chem.*, *172*, 78–85. https://doi.org/10.1016/j.foodchem.2014.09.036

Moselhy, S. S., Razvi, S. S., Al shibili, F. A., Kuerban, A., Hasan, M. N., Balamash, K. S., *et al*. (2018). m-Coumaric acid attenuates non-catalytic protein glycosylation in the retinas of diabetic rats. *J Pestic Sci*, *43*(3), 180–185.

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

https://doi.org/10.1584/jpestics.D17-091

Motawi, T. M. K., Abou-Seif, M. A., Bader, A. M. A., & Mahmoud, M. O. (2013). Effect of glycemic control on soluble RAGE and oxidative stress in type 2 diabetic patients. *BMC Endocr. Disord.*, *13*(1), 1. https://doi.org/10.1186/1472-6823-13-32

Mukai, Y., Shibata, H., Nakamura, T., Yoshioka, Y., Abe, Y., Nomura, T., *et al.* (2009). Structure-function relationship of tumor necrosis factor (TNF) and its receptor interaction based on 3D structural analysis of a fully active TNFR1-selective TNF mutant. *J. Mol. Biol.*, *385*(4), 1221–1229. https://doi.org/10.1016/j.jmb.2008.11.053

Muramatsu, D., Uchiyama, H., Kida, H., & Iwai, A. (2019). Cell cytotoxity and anti-glycation activity of taxifolin-rich extract from Japanese larch, Larix kaempferi. *Heliyon*, *5*(7), e02047. https://doi.org/10.1016/j.heliyon.2019.e02047

Murini, T., Wahyuningsih, M. S. H., Satoto, T. B. T., Fudholi, A., & Hanafi, M. (2018). Isolation and identification of naturally occurring larvicidal compound isolated from zingiber zerumbet (L).J.E. smith. *Asian J. Pharm. Clin. Res.*, *11*(2), 189–193. https://doi.org/10.22159/ajpcr.2018.v11i2.21703

Nagarani, N., & Kumaraguru, A. K. (2013). Evaluation of anti-inflammatory, antidiabetic, cytotoxic activity of Kappaphycus alvarezii. *Int J Pharm Bio Sci*, *4(1)*(January 2013), 921–929.

Negre-Salvayre, A., Salvayre, R., Augé, N., Pamplona, R., & Portero-Otín, M. (2009). Hyperglycemia and glycation in diabetic complications. *Antioxid. Redox Signaling*, *11*(12), 3071–3109. https://doi.org/10.1089/ars.2009.2484

Nevin, C., McNeil, L., Ahmed, N., Murgatroyd, C., Brison, D., & Carroll, M. (2018). Investigating the glycating effects of glucose, glyoxal and methylglyoxal on human sperm. *Sci. Rep.*, *8*(1), 1–12. https://doi.org/10.1038/s41598-018-27108-7

Niedowics, D. M., & Daleke, D. L (2005). The role of oxidative stress in diabetic complications the role of oxidative stress. *Cell Biochem Biophys*, *43*(October 2015), 289–290. https://doi.org/10.1385/CBB

Nowotny, K., Jung, T., Höhn, A., Weber, D., & Grune, T. (2015). Advanced glycation end products and oxidative stress in type 2 diabetes mellitus. *Biomolecules*, *5*(1), 194–222. https://doi.org/10.3390/biom5010194

Nurilmala, M., Anwar, E., & Luthfiyana, N. (2018). Identification of bioactive compounds of seaweed Sargassum sp . and Eucheuma cottonii Doty as a raw sunscreen cream, *Proceedings of the Pakistan Academy of Sciences 54*(February), 311–318.

Nwosu, F., Morris, J., Lund, V. A., Stewart, D., Ross, H. A., & McDougall, G. J. (2011). Anti-proliferative and potential anti-diabetic effects of phenolic-rich extracts from edible marine algae. *Food Chem.*, *126*(3), 1006–1012. https://doi.org/10.1016/j.foodchem.2010.11.111

Oliveira-Marques, V., Marinho, H. S., Cyrne, L., & Antunes, F. (2009). Role of hydrogen peroxide in NF-κB activation: From inducer to modulator. *Antioxid Redox Signal*, *11*(9), 2223–2243. https://doi.org/10.1089/ars.2009.2601

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Ott, C., Jacobs, K., Haucke, E., Navarrete Santos, A., Grune, T., & Simm, A. (2014). Role of advanced glycation end products in cellular signaling. *Redox Biol.*, *2*(1), 411–429. https://doi.org/10.1016/j.redox.2013.12.016

Oyedemi, S. O., Oyedemi, B. O., Coopoosamy, R. M., Prieto, J. M., Stapleton, P., & Gibbons, S. (2017). Antibacterial and norfloxacin potentiation activities of Ocimum americanum L. against methicillin resistant Staphylococcus aureus. *S. Afr. J. Bot.*, *109*, 308–314. https://doi.org/10.1016/j.sajb.2016.12.025

Ozougwu, J. C., Obimba, K. C., Belonwu, C. D., & Unakalamba, C. B. (2013). The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *J. Physiol. Pathophysiol.*, *4*(4), 46–57. https://doi.org/10.5897/jpap2013.0001

Papagrigoraki, A., Maurelli, M., Del Giglio, M., Gisondi, P., & Girolomoni, G. (2017). Advanced glycation end products in the pathogenesis of psoriasis. *Int. J. Mol. Sci.*, *18*(11), 1–9. https://doi.org/10.3390/ijms18112471

Papatheodorou, K., Banach, M., Bekiari, E., Rizzo, M., & Edmonds, M. (2018). Complications of diabetes 2017. *J Diabetes Res.*, Mar 11, 10–13. https://doi.org/10.1155/2018/3086167

Papatheodorou, K., Papanas, N., Banach, M., Papazoglou, D., & Edmonds, M. (2016). Complications of diabetes 2016. *J Diabetes Res*, *2016*(September). https://doi.org/10.1155/2016/6989453

Parameswaran, N., & Patial, S. (2010). Tumor Necrosis Factor-α signalling in macrophages. *Crit Rev Eukaryot Gene Expr.*, *20*(2), 87–103. https://doi.org/10.1016/j.dcn.2011.01.002.

Park, K. H., Kim, S. J., Kim, J. Y., Min, K. J., Kim, J. R., & Cho, K. H. (2016). Cold-water extract of Korean Red Ginseng exhibits potent inhibitory effects against cholesteryl ester transfer protein activity and fructose-mediated glycation along with lipid-lowering activity in hyperlipidemic zebrafish. *J. Ginseng Res.*, 1–12. https://doi.org/10.1016/j.jgr.2017.01.005

Patel, S., & Santani, D. (2009). Role of NF-κB in the pathogenesis of diabetes and its associated complications. *Pharmacol. Rep.*, *61*(4), 595–603. https://doi.org/10.1016/S1734-1140(09)70111-2

Peng, X., Ma, J., Chen, F., & Wang, M. (2011). Naturally occurring inhibitors against the formation of advanced glycation end-products. *Food Funct.*, *2*(6), 289–301. https://doi.org/10.1039/c1fo10034c

Perez Gutierrez, R. M., & Velazquez, E. G. (2020). Glucopyranoside flavonoids isolated from leaves of Spinacia oleracea (spinach) inhibit the formation of advanced glycation end products (AGEs) and aldose reductase activity (RLAR). *Biomed. Pharmacother*, *128*(April), 110299. https://doi.org/10.1016/j.biopha.2020.110299

Perrone, A., Giovino, A., Benny, J., & Martinelli, F. (2020). Advanced Glycation End Products (AGEs): biochemistry, signaling, analytical methods, and epigenetic Effects. *Oxid. Med. Cell. Longevity*, *2020*(Cml). https://doi.org/10.1155/2020/3818196

Pia de la Maza, M., Garrido, F., Escalante, N., Leiva, L., Barrera, G., Schnitzler, S., *et*

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

al. (2012). Fluorescent advanced glycation end-products (ages) detected by spectro-photofluorimetry, as a screening tool to detect diabetic microvascular complications. *J. Diabetes Mellit.*, *02*(02), 221–226. https://doi.org/10.4236/jdm.2012.22035

Picot, M. C. N., Bender, O., Atalay, A., Zengin, G., Loffredo, L., Hadji-Minaglou, F., *et al.*. (2017). Multiple pharmacological targets, cytotoxicity, and phytochemical profile of Aphloia theiformis (Vahl.) Benn. *Biomed Pharmacother.*, *89*, 342–350. https://doi.org/10.1016/j.biopha.2017.02.031

Picu, A., Petcu, L., Stefan, S., Mitu, M., Lixandru, D., Ionescu-Tîrgoviste, C., *et al.*. (2017). Markers of oxidative stress and antioxidant defense in romanian patients with type 2 diabetes mellitus and obesity. *Molecules*, *22*(5), 1–14. https://doi.org/10.3390/molecules22050714

Popov, A. M., & Krivoshapko, O. N. (2013). Protective effects of polar lipids and redox-active compounds from marine organisms at modeling of hyperlipidemia and diabetes, *J. Biomed. Sci. Eng. 2013*(May), 543–550. https://doi.org/10.4236/jbise.2013.65069

Popova, E. A., Mironova, R. S., & Odjakova, M. K. (2010). Non-enzymatic glycosylation and deglycating enzymes. *Biotechnol. Biotechnol. Equip.*, *24*(3), 1928–1935. https://doi.org/10.2478/V10133-010-0066-7

Prasad, K., & Mishra, M. (2018). AGE–RAGE stress, stressors, and antistressors in health and disease. *Int J Angiol*, *27*(1), 1–12. https://doi.org/10.1055/s-0037-1613678

Prieto-bermejo, R., & Hern, A. (2017). The Importance of NADPH Oxidases and redox signaling in angiogenesis. *Antioxidants*, *6*(2), 32. https://doi.org/10.3390/antiox6020032

Qais, F. A., Alam, M. M., Naseem, I., & Ahmad, I. (2016). Understanding the mechanism of non-enzymatic glycation inhibition by cinnamic acid: An in vitro interaction and molecular modelling study. *RSC Advances*, *6*(70). https://doi.org/10.1039/c6ra12321j

Quehenberger, P., Bierhaus, A., Fasching, P., Muellner, C., Klevesath, M., Hong, M., *et al*. (2000). Endothelin 1 transcription is controlled by nuclear factor-κB in AGE-stimulated cultured endothelial cells. *Diabetes*, *49*(9), 1561–1570. https://doi.org/10.2337/diabetes.49.9.1561

Radoi, V., Lixandru, D., Mohora, M., & Virgolici, B. (2012). Advanced glycation end products in diabetes mellitus: mechanism of action and focused treatment. *Proc Rom Acad Series B*, *1*, 9–19.

Rahimi-Madiseh, M., Malekpour-Tehrani, A., Bahmani, M., & Rafieian-Kopaei, M. (2016). The research and development on the antioxidants in prevention of diabetic complications. *Asian Pac J Trop Med*, *9*(9), 825–831. https://doi.org/10.1016/j.apjtm.2016.07.001

Rajasulochana P., K. P. and D. R. (2012). Potential application of Kappaphycus alvarezii in agricultural and pharmaceutical industry. *J Chem Pharm Res*, *4*(1), 33–37.

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Ramana, K. V., Friedrich, B., Srivastava, S., Bhatnagar, A., & Srivastava, S. K. (2004). Activation of nuclear factor-κB by hyperglycemia in vascular smooth muscle cells is regulated by aldose reductase. *Diabetes*, *53*(11), 2910–2920. https://doi.org/10.2337/diabetes.53.11.2910

Ramdhan, Y. M. (2018). Pengelolaan budidaya rumput laut berbasis daya dukung lingkungan perairan di pesisir kabupaten Dompu, Provinsi Nusa Tenggara Barat. *Seminar Nasional Geomatika*, *2*(February), 01. https://doi.org/10.24895/SNG.2017.2-0.391

Ramkissoon, J. S., Mahomoodally, M. F., Subratty, A. H., & Ahmed, N. (2016). Inhibition of glucose- and fructose-mediated protein glycation by infusions and ethanolic extracts of ten culinary herbs and spices. *Asian Pac J Trop Biomed*, *6*(6), 492–500. https://doi.org/10.1016/j.apjtb.2016.01.016

Raupbach, J., Ott, C., Koenig, J., & Grune, T. (2020). Proteasomal degradation of glycated proteins depends on substrate unfolding: Preferred degradation of moderately modified myoglobin. *Free Radical Biology and Medicine*, *152*(July 2019), 516–524. https://doi.org/10.1016/j.freeradbiomed.2019.11.024

Rhee, S. Y., & Kim, Y. S. (2018). The role of advanced glycation end products in diabetic vascular complications. *Diabetes Metab J*, *42*(3), 188–195. https://doi.org/10.4093/dmj.2017.0105

Rimando, A. M., Olofsdotter, M., Dayan, F. E., & Duke, S. O. (2001). Searching for rice allelochemicals. *Agron. J.*, *93*(1), 16–20. https://doi.org/10.2134/agronj2001.93116x

Sadowska-Bartosz, I., & Bartosz, G. (2015). Prevention of protein glycation by natural compounds. *Molecules*, *20*(2), 3309–3334. https://doi.org/10.3390/molecules20023309

Saide, A., Lauritano, C., & Ianora, A. (2020). Pheophorbide A: State of the art. *Mar. Drugs*, *18*(5), 1–12. https://doi.org/10.3390/md18050257

Salami, M., Rahimmalek, M., & Ehtemam, M. H. (2016). Inhibitory effect of different fennel (Foeniculum vulgare) samples and their phenolic compounds on formation of advanced glycation products and comparison of antimicrobial and antioxidant activities. *Food Chem.*, *213*, 196–205. https://doi.org/10.1016/j.foodchem.2016.06.070

Sampath, C., Rashid, M. R., Sang, S., & Ahmedna, M. (2017). Green tea epigallocatechin 3-gallate alleviates hyperglycemia and reduces advanced glycation end products via nrf2 pathway in mice with high fat diet-induced obesity. *Biomed. Pharmacother.*, *87*, 73–81. https://doi.org/10.1016/j.biopha.2016.12.082

Sanz, A. B., Sanchez-Niño, M. D., Ramos, A. M., Moreno, J. A., Santamaria, B., Ruiz-Ortega, M., *et al*. (2010). NF-κB in renal inflammation. *J. Am. Soc. Nephrol.*, *21*(8), 1254–1262. https://doi.org/10.1681/ASN.2010020218

Sasidharan, S., Chen, Y., Saravanan, D., Sundram, K. M., & Latha, L. Y. (2011). Extraction, isolation and characterization of bioactive compounds from plants' extracts. *Afr J Tradit Complement Altera Med*, *8*(1), 1–10.

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

https://doi.org/10.4314/ajtcam.v8i1.60483

Schinzela, R., Münchb, G., Heidlandc, A., & Sebekova, K. (2001). Advanced glycation end products in end-stage renal disease and their removal. *Nephron*, *87*(4), 295–303. https://doi.org/10.1159/000045934

Schmidt, A. M., Yan, S. Du, Yan, S. F., & Stern, D. M. (2000). The biology of the receptor for advanced glycation end products and its ligands. *Biochim. Biophys. Acta, Mol. Cell Res.*, *1498*(2–3), 99–111. https://doi.org/10.1016/S0167-4889(00)00087-2

Schröder, K. (2015). NADPH oxidases in bone homeostasis and osteoporosis. *Cell. Mol. Life Sci.*, *72*(1), 25–38. https://doi.org/10.1007/s00018-014-1712-2

Selvan, K. B., Piriya, S. P., & Vennison, J. S. (2014). Macro algae (Eucheuma cottoni and Sargassum Sp.) are reservoirs of biodiesel and bioactive compounds. *J. Chem. Pharm. Sci JCHPS Special Issue*, *2*(2), 974–2115.

Semba, R. D., Nicklett, E. J., & Ferrucci, L. (2010). Does accumulation of advanced glycation end products contribute to the aging phenotype? J *Gerontol A Biol Sci Med Sci*, *65 A*(9), 963–975. https://doi.org/10.1093/gerona/glq074

Serasanambati, M., & Chilakapati, S. R. (2016). Function of Nuclear Factor Kappa B (NF-kB) in human diseases-a review. *South Indian J. Biol. Sci*, *2*(4), 368. https://doi.org/10.22205/sijbs/2016/v2/i4/103443

Serban, A. I., Stanca, L., Geicu, O. I., & Dinischiotu, A. (2015). Ages-induced IL-6 synthesis precedes RAGE up-regulation in HEK 293 cells: An alternative inflammatory mechanism? *Int. J. Mol. Sci.*, *16*(9), 20100–20117. https://doi.org/10.3390/ijms160920100

Serdiati, N., & Widiastuti, I. M. (2010). Pertumbuhan dan produksi rumput laut *Eucheuma cottonii* pada kedalaman penanaman yang berbeda. *Media Litbang Sulteng*, *III*(1), 21–26.

Shah, S., Baez, E. A., Felipe, D. L., Maynard, J. D., Hempe, J. M., & Chalew, S. A. (2013). Advanced glycation endproducts in children with diabetes. *J. Pediatr.*, *163*(5), 1427–1431. https://doi.org/10.1016/j.jpeds.2013.06.044

Shi, D., Guo, S., Jiang, B., Guo, C., Wang, T., Zhang, L., *et al.*. (2013). HPN, a synthetic analogue of bromophenol from red alga Rhodomela confervoides: Synthesis and anti-diabetic effects in C57BL/KsJ-db/db mice. *Mar. Drugs*, *11*(2), 350–362. https://doi.org/10.3390/md11020350

Siddiqui, H., Bashir, M. A., Javaid, K., Nizamani, A., Bano, H., Yousuf, S., *et al*. (2016). Ultrasonic synthesis of tyramine derivatives as novel inhibitors of α-glucosidase in vitro. *J. Enzyme Inhib. Med. Chem.*, *31*(6), 1392–1403. https://doi.org/10.3109/14756366.2016.1142983

Sifuentes-Franco, S., Padilla-Tejeda, D. E., Carrillo-Ibarra, S., & Miranda-Díaz, A. G. (2018). Oxidative stress, apoptosis, and mitochondrial function in diabetic nephropathy. *Int. J. Endocrinol.*, *2018*. https://doi.org/10.1155/2018/1875870

Singh, V. P., Bali, A., Singh, N., & Jaggi, A. S. (2014a). Advanced glycation end products and diabetic complications. *Korean J Physiol Pharmacol*, *18*(1), 1–14. https://doi.org/10.4196/kjpp.2014.18.1.1

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Singh, M. P., & Pathak, K. (2015). Animal models for biological screening of anti-diabetic drugs: An overview. *Eur. J. Exp. Biol.*, *5*(5), 37–48. Retrieved from http://www.imedpub.com/articles/animal-models-for-biological-screening-of-antidiabetic-drugs-an-overview.pdf

Silván, J. M., Assar, S. H., Srey, C., Dolores Del Castillo, M., & Ames, J. M. (2011). Control of the Maillard reaction by ferulic acid. *Food Chem.*, *128*(1), 208–213. https://doi.org/10.1016/j.foodchem.2011.03.047

Siriamornpun, S., Weerapreeyakul, N., & Barusrux, S. (2015). Bioactive compounds and health implications are better for green jujube fruit than for ripe fruit. *J. Funct. Foods*, *12*, 246–255. https://doi.org/10.1016/j.jff.2014.11.016

Soenardjo, N. (2011). Aplikasi budidaya rumput laut *Eucheuma cottonii* (Weber van Bosse) dengan Metode Jaring Lepas Dasar (Net Bag) Model Cidaun. *Buletin Oseanografi Marina*, *1*(1), 36–44. Retrieved from https://ejournal.undip.ac.id/index.php/buloma/article/view/2970

Stirban, A., Gawlowski, T., & Roden, M. (2014). Vascular effects of advanced glycation endproducts : Clinical effects and molecular mechanisms. *Mol. Metab.*, *3*(2), 94–108. https://doi.org/10.1016/j.molmet.2013.11.006

Sudirman, S., Hsu, Y., He, J., & Kong, Z. (2018). Red seaweed *Eucheuma cottonii* ethanol extract regulates inflammatory mediators production and prevents colonic injury on acute colitis disease in mice, (June), 1–11. https://doi.org/10.20944/preprints201806.0120.v1

Suganya, A. M., Sanjivkumar, M., Chandran, M. N., Palavesam, A., & Immanuel, G. (2016). Pharmacological importance of sulphated polysaccharide carrageenan from red seaweed Kappaphycus alvarezii in comparison with commercial carrageenan. *Biomed. Pharmacother.*, *84*, 1300–1312. https://doi.org/10.1016/j.biopha.2016.10.067

Sumayya, S., & Murugan, K. (2017). Phytochemical screening, RP-HPLC and FTIR analysis of Kappaphycus alvarezii (Doty) Doty EX P.C Silva: Macro red algae. *J. Pharmacogn. Phytochem.*, *6*(1), 325–330.

Sun, Q., Li, J., & Gao, F. (2014). New insights into insulin: The anti-inflammatory effect and its clinical relevance. *World Journal of Diabetes*, *5*(2), 89. https://doi.org/10.4239/wjd.v5.i2.89

Suryavanshi, S. V., & Kulkarni, Y. A. (2017). NF-κβ: A potential target in the management of vascular complications of diabetes. *Front Pharmacol*, *8*(NOV), 1–12. https://doi.org/10.3389/fphar.2017.00798

Swaroop, J. J., Rajarajeswari, D., & Naidu, J. N. (2012). Association of TNF-α with insulin resistance in type 2 diabetes mellitus. *Indian J Med Res*, *135*(1), 127–130. https://doi.org/10.4103/0971-5916.93435

Syarif, R. A., Wahyuningsih, M. S. H., Mustofa, M., & Ngatidjan, N. (2018). Antiplasmodial and onset speed of growth inhibitory activities of *Tithonia diversifolia* (Hemsley) A gray leaf fractions against *Plasmodium falciparum*. *Trop J Pharm Res*, *17*(11), 2213–2218. https://doi.org/10.4314/tjpr.v17i11.15

Szymczak, I., Śliwińska, A., & Drzewoski, J. (2014). DNA damage and efficacy of

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

DNA repair in patients with type 2 diabetes and coexisting colorectal cancer. *Pol Arch Intern Med*, *124*(7–8), 352–358. https://doi.org/10.20452/pamw.2337

Takahashi, S., Uchino, H., Shimizu, T., Kanazawa, A., Tamura, Y., Sakai, K., *et al*. (2007). Comparison of Glycated Albumin (GA) and Glycated Hemoglobin (HbA1c) in type 2 diabetic patients: Usefulness of GA for evaluation of short-term changes in glycemic control. *Endocr. J.*, *54*(1), 139–144. https://doi.org/10.1507/endocrj.K06-103

Tangko, A. M. (2009). Budidaya Rumput Laut Di Provinsi Sulawesi Selatan.

Tanji, N., Markowitz, G. S., Fu, C., Kislinger, T., Taguchi, A., Pischetsrieder, M., *et al*. (2000). Expression of advanced glycation end products and their cellular receptor RAGE diabetic nephropathy and nondiabetic renal disease. *J. Am. Soc. Nephrol.*, *11*(9), 1656–1666.

Thornalley, P. J., Häussinger, D., Lüthen, R., Heidland, A., Schinzel, R., Ahmed, N., *et al*. (2005). Increased protein glycation in cirrhosis and therapeutic strategies to prevent it. *Ann. N. Y. Acad. Sci.*, *1043*(1), 718–724. https://doi.org/10.1196/annals.1333.083

Tiwari, B. K., Pandey, K. B., Abidi, A. B., & Rizvi, S. I. (2013). Markers of oxidative stress during diabetes mellitus. *J. Biomark.*, *2013*, 1–8. https://doi.org/10.1155/2013/378790

Tomobe, Y. I., Morizawa, K., Tsuchida, M., Hibino, H., Nakano, Y., & Tanaka, Y. (2000). Dietary docosahexaenoic acid suppresses inflammation and immunoresponses in contact hypersensitivity reaction in mice. *Lipids*, *35*(1), 61–69. https://doi.org/10.1007/s11745-000-0495-0

Trask, O. J., Jr (2012). Nuclear Factor Kappa B (NF-κB) Translocation assay development and validation for high content screening. In S. Markossian (Eds.) et. al., *Assay Guidance Manual*. Eli Lilly & Company and the National Center for Advancing Translational Sciences. PMID: 23035273. P 554

Tripathi, B. K., & Srivastava, A. K. (2006). Diabetes mellitus: Complications and therapeutics. *Med. Sci. Monit.*, *12*(7), RA130–RA147. https://doi.org/4187 [pii]

Tupe, R. S., Kulkarni, A., Adeshara, K., Shaikh, S., Shah, N., & Jadhav, A. (2015). Syzygium jambolanum and Cephalandra indica homeopathic preparations inhibit albumin glycation and protect erythrocytes: An invitro study. *Homeopathy*, *104*(3), 197–204. https://doi.org/10.1016/j.homp.2015.02.009

Venskutonis, P. R., Miliauskas, G., & Sivik, B. (2009). Extraction and fractionation of bioactive compounds. *J. Liq. Chromatogr.*, *25*, 317. https://doi.org/10.1186/1471-2105-14-370

Vlassara, H., & Uribarri, J. (2014). Advanced Glycation End Products (AGE) and Diabetes: cause, effect, or both? *Curr Diab Rep*, *14*(1), 453. https://doi.org/10.1007/s11892-013-0453-1.Advanced

Volpe, C. M. O., Villar-Delfino, P. H., Dos Anjos, P. M. F., & Nogueira-Machado, J. A. (2018). Cellular death, reactive oxygen species (ROS) and diabetic complications review-Article. *Cell Death Dis.*, *9*(2). https://doi.org/10.1038/s41419-017-0135-z

UNIVERSITAS GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Wahyuningsih, M. S. H., Mubarika, S., Ganjar, I. G., Wahyuono, S., & Takeya, T. (2017). 5α-Oleandrin reduce Bcl-2 protein and increase Bax protein expression on Hela cervical cancer cell. *UnivMed*, *36*(2), 102. https://doi.org/10.18051/univmed.2017.v36.102-109

Wahyuningsih, M. S. H., Wijayanti, M. A., Budiyanto, A., & Hanafi, M. (2015). Isolation and identification of potential cytotoxic compound from kembang bulan [tithonia diversifolia (Hemsley) a gray] leaves. *Int J Pharm Pharm Sci*, *7*(6), 298–301.

Wang, W., Okada, Y., Shi, H., Wang, Y., & Okuyama, T. (2005). Structures and aldose reductase inhibitory effects of bromophenols from the red alga Symphyocladia latiuscula. *J. Nat. Prod.*, *68*(4), 620–622. https://doi.org/10.1021/np040199j

Weiss, M. F., Erhard, P., Kader-Attia, F. A., Wu, Y. C., Deoreo, P. B., Araki, A., *et al*. (2000). Mechanisms for the formation of glycoxidation products in end-stage renal disease. *Kidney Int*, *57*(6), 2571–2585. https://doi.org/10.1046/j.1523-1755.2000.00117.x

Wu, L., Ma, L., Nicholson, L. F. B., & Black, P. N. (2011). Advanced glycation end products and its receptor (RAGE) are increased in patients with COPD. *Respir. Med.*, *105*(3), 329–336. https://doi.org/10.1016/j.rmed.2010.11.001

Xue, J., Rai, V., Frolov, S., Singer, D., Chabierski, S., Xie, J., *et al*. (2012). Advanced glycation end product (AGE) recognition by the receptor for AGEs (RAGE), *Structure 19*(5), 722–732. https://doi.org/10.1016/j.str.2011.02.013.

Yamada, H., Umemoto, T., Kakei, M., Momomura, S. I, Kawakami, M., Ishikawa, S. e., *et al.* (2017). Eicosapentaenoic acid shows anti-inflammatory effect via GPR120 in 3T3-L1 adipocytes and attenuates adipose tissue inflammation in diet-induced obese mice. *Nutr. Metab.*, *14*(1), 1–11. https://doi.org/10.1186/s12986-017-0188-0

Yao, D., & Brownlee, M. (2010). Hyperglycemia-induced reactive oxygen species increase expression of the receptor for advanced glycation end products (RAGE) and RAGE ligands. *Diabetes*, *59*(1), 249–255. https://doi.org/10.2337/db09-0801

Yazdanpanah, S., Rabiee, M., Tahriri, M., Abdolrahim, M., Rajab, A., Jazayeri, H. E., *et al*. (2017). Evaluation of glycated albumin (GA) and GA/HbA1c ratio for diagnosis of diabetes and glycemic control: A comprehensive review. *Crit. Rev. Clin. Lab. Sci.*, *54*(4), 219–232. https://doi.org/10.1080/10408363.2017.1299684

Yang, P., Feng, J., Peng, Q., Liu, X., Fan, Z., & Luca, M. (2019). Advanced Glycation End Products: potential mechanism and therapeutic target in cardiovascular complications under diabetes. *Oxid. Med. Cell. Longev.*, *2019*. https://doi.org/10.1155/2019/9570616

Yehualashet, A. S. (2020). Toll-like receptors as a potential drug target for diabetes mellitus and diabetes-associated complications. *Diabetes Metab Syndr Obes.: Targets and Therapy*, *13*, 4763–4777. https://doi.org/10.2147/DMSO.S274844

Yoon, H. J., Lee, Y. H., Kim, K. J., Kim, S. R., Kang, E. S., Cha, B. S., *et al.*. (2015). Glycated albumin levels in patients with type 2 diabetes increase relative to HbA1c with time. *Biomed Res. Int*, *2015*. https://doi.org/10.1155/2015/576306

UNIVERSITAS
GADJAH MADA
Efek Fraksi Aktif Alga Merah Kappaphycus alvarezii (Doty) Doty ex P.C Silva terhadap Homeostasis Glukosa Darah pada Tikus Model Diabetik. Kajian : Kadar Glycated Albumin, Ne-(carboxymethyl) lysine, H 2O2 , TNF alpha, Ekspresi gen RAGE (Ager), NADPH oksidase 4 (NOX4) dan NFkappaB
EVY YULIANTI, Prof. Dr. Dra. Mae Sri Hartati W. M.Si., Apt.; Prof. Dr. Sunarti, M.Kes
Universitas Gadjah Mada, 2021 | Diunduh dari http://etd.repository.ugm.ac.id/

Yoshiuchi, K., Matsuhisa, M., Katakami, N., Nakatani, Y., Sakamoto, K., Matsuoka, T., Hori, M. (2008). Glycated albumin is a better indicator for glucose excursion than glycated hemoglobin in type 1 and type 2 diabetes. *Endocr. J*, *55*(3), 503–507. https://doi.org/10.1507/endocrj.K07E-089

Younus, H., & Anwar, S. (2016). Prevention of non-enzymatic glycosylation (glycation): Implication in the treatment of diabetic complication. *Int. J. Health Sci.*, *10*(2). Retrieved from http://ijhs.org.sa/index.php/journal/article/view/1310

Yu, P., Xu, X. B., & Yu, S. J. (2017). Inhibitory effect of sugarcane molasses extract on the formation of Nε-(carboxymethyl)lysine and Nε-(carboxyethyl)lysine. *Food Chem.*, *221*, 1145–1150. https://doi.org/10.1016/j.foodchem.2016.11.045

Yuan, T., Yang, T., Chen, H., Fu, D., Hu, Y., Wang, J., *et al*. (2019). New insights into oxidative stress and inflammation during diabetes mellitus-accelerated atherosclerosis. *Redox Biol.*, *20*(August 2018), 247–260. https://doi.org/10.1016/j.redox.2018.09.025

Zhang, M., Otake, K., Miyauchi, Y., Yagi, M., Yonei, Y., Miyakawa, T., *et al*. (2019). Comprehensive NMR analysis of two kinds of post-fermented tea and their anti-glycation activities in vitro. *Food Chem.*, *277*(July 2018), 735–743. https://doi.org/10.1016/j.foodchem.2018.11.028

Zhang, Q. W., Lin, L. G., & Ye, W. C. (2018). Techniques for extraction and isolation of natural products: A comprehensive review. *Chin. Med. (United Kingdom)*, *13*(1), 1–26. https://doi.org/10.1186/s13020-018-0177-x

Zhao, C., Yang, C., Liu, B., Lin, L., Sarker, S. D., Nahar, L., *et al*. (2018). Bioactive compounds from marine macroalgae and their hypoglycemic benefits. *Trends Food Sci. Technol.*, *72*(December), 1–12. https://doi.org/10.1016/j.tifs.2017.12.001