

## DAFTAR PUSTAKA

- Aba, L. *et al.* (2014) 'Sensitivity improvement of ammonia gas sensor based on poly(3,4-ethylenedioxythiophene):poly(styrenesulfonate) by employing doping of bromocresol green', *Journal of Nanotechnology*, 2014, pp. 1–5. doi: 10.1155/2014/864274.
- Adaramoye, O. A. *et al.* (2011) 'Antiproliferative action of *Xylopia aethiopica* fruit extract on human cervical cancer cells', *Phytotherapy Research*, 25(10), pp. 1558–1563. doi: 10.1002/ptr.3551.
- Ademiluyi, A. O. *et al.* (2013) 'Anthocyanin - rich red dye of *Hibiscus sabdariffa* calyx modulates cisplatin-induced nephrotoxicity and oxidative stress in rats', *International Journal of Biomedical Science*, 9(4), pp. 243–248.
- Ademiluyi, Adedayo O. and Oboh, G. (2013) 'Aqueous extracts of Roselle (*Hibiscus sabdariffa* Linn.) varieties inhibit  $\alpha$ -amylase and  $\alpha$ -glucosidase activities in vitro', *Journal of Medicinal Food*, 16(1), pp. 88–93. doi: 10.1089/jmf.2012.0004.
- Aderibigbe, S. A. *et al.* (2017) 'Knowledge and practice of sedentary lifestyle among bankers in Abuja, North-Central Ngeria', *Research Journal of Health Sciences*, 5(3), p. 167. doi: 10.4314/rejhs.v5i3.6.
- Adeyemi, D. O. *et al.* (2014) 'Anti-hepatotoxic activities of *Hibiscus sabdariffa* L. in animal model of streptozotocin diabetes-induced liver damage', *BMC Complementary and Alternative Medicine*, 14(1), p. 227. doi: 10.1186/1472-6882-14-277.
- Affourtit, C. and Brand, M. D. (2008) 'On the role of uncoupling protein-2 in pancreatic beta cells', *Biochimica et Biophysica Acta - Bioenergetics*, 1777(7–8), pp. 973–979. doi: 10.1016/j.bbabi.2008.03.022.
- Agrawal, N. *et al.* (2017) 'Correlation between body mass index and blood glucose levels in Jharkhand population', *International Journal of Contemporary Medical Research*, 4(8), p. 1633.
- Ajay, M. *et al.* (2007) 'Mechanisms of the anti-hypertensive effect of *Hibiscus sabdariffa* L. calyces', *Journal of Ethnopharmacology*, 109(3), pp. 388–393. doi: 10.1016/j.jep.2006.08.005.
- Akbarzadeh, A. *et al.* (2007) 'Induction of diabetes by Streptozotocin in rats', *Indian Journal of Clinical Biochemistry*, 22(2), pp. 60–64. doi: 10.1007/BF02913315.

- Akindahunsi, A. A. and Olaleye, M. T. (2003) 'Toxicological investigation of aqueous-methanolic extract of the calyces of *Hibiscus sabdariffa* L.', *Journal of Ethnopharmacology*, 89(1), pp. 161–164. doi: 10.1016/S0378-8741(03)00276-9.
- Akram, S. *et al.* (2015) 'Determination of heavy metal contents by atomic absorption spectroscopy (AAS) in some medicinal plants from Pakistani and Malaysian origin', *Pakistan Journal of Pharmaceutical Sciences*, 28(5), pp. 1781–1787.
- Alarcon-Aguilar, F. J., Zamilpa, A., Perez-Garcia, M. D., Almanza-Perez, J. C., Romero-Nuñez, E., Campos-Sepulveda, E. A., Vazquez-Carrillo, L. I., & Roman-Ramos, R. (2007). Effect of *Hibiscus sabdariffa* on obesity in MSG mice. *Journal of ethnopharmacology*, 114(1), 66–71. <https://doi.org/10.1016/j.jep.2007.07.020>
- Albareda, M. *et al.* (2000) 'Assessment of insulin sensitivity and beta-cell function from measurements in the fasting state and during an oral glucose tolerance test', *Diabetologia*, 43(12), pp. 1507–1511. doi: 10.1007/s001250051561.
- Ali, B. H., Al Wabel, N. and Blunden, G. (2005) 'Phytochemical, pharmacological and toxicological aspects of *Hibiscus sabdariffa* L.: A review', *Phytotherapy Research*, 19(5), pp. 369–375. doi: 10.1002/ptr.1628.
- Allorerung, D. and Sekeon, S. (2016) 'Hubungan antara umur, jenis kelamin dan tingkat pendidikan dengan kejadian Diabetes Melitus Tipe 2 Di Puskesmas Ranotana Weru Kota Manado Tahun 2016', *FKM Sam Ratulangi*, 8(3), pp. 1–8.
- American Diabetes Association (2018) 'Standards of medical care in diabetes', *The Journal of Clinical and Applied Research and Education*, 41(January). doi: <https://doi.org/10.2337/dc18-Sint01>.
- Amin, R. *et al.* (2014) 'Study and impact evaluation of particle size distribution on physicochemical properties of three different tablet formulations through sieve technology', *International Journal for Pharmaceutical Research Scholars (IJPRS)*, 3(January), pp. 448–563.
- Andersen, G. *et al.* (2013) 'The frequent UCP2 –866G/A polymorphism protects against insulin resistance and is associated with obesity: a study of obesity and related metabolic traits among 17 636 Danes', *International Journal of Obesity*, 37(2), pp. 175–181. doi: 10.1038/ijo.2012.22.
- Andrainsi, T. and Yolanda, S. (2014) 'Prevention of insulin resistance with *Hibiscus sabdariffa* Linn. extract in high-fructose fed rat', *Medical Journal of Indonesia*, 23(4), pp. 192–196. doi: 10.13181/mji.v23i4.848.

- Ansel. H.C. (2008). *Pengantar bentuk sediaan farmasi*. 4th edn. Jakarta: UI Press.
- Ardalani, Hamidreza. (2016) ‘The ethnopharmacological review of medicinal plants in uses hypertension treatment in Iranian complementary medicine, *Journal of Herbal Drug*’, 7(1), pp. 9–13.
- Asiimwe, D., Mauti, G. O. and Kiconco, R. (2020) ‘Prevalence and risk factors associated with type 2 diabetes in elderly patients aged 45-80 years at Kanungu District’, *Journal of Diabetes Research*, 2020. doi: 10.1155/2020/5152146.
- Asman, M. (2014) ‘Insulin resistance as a predictor of worsening of glucose tolerance in type 2 diabetes mellitus’, *Medicinus*, 27(2), pp. 3–8.
- Atiqoh, H., Wardani, R. S. and Wulandari, M. (2011) ‘Uji antidiabetik infusa kelopak bunga Rosella (*Hibiscus sabdariffa* Linn.) Pada tikus putih jantan galur wistar yang diinduksi glukosa’, *Jurnal.Unimus.Ac.Id* 4, 7(1), pp. 43–50.
- Awa, W. L. *et al.* (2012) ‘Type 2 diabetes from pediatric to geriatric age: Analysis of gender and obesity among 120 183 patients from the German/Austrian DPV database’, *European Journal of Endocrinology*, 167(2), pp. 245–254. doi: 10.1530/EJE-12-0143.
- Aynalem, S. B. and Zeleke, A. J. (2018) ‘Prevalence of diabetes mellitus and its risk factors among individuals aged 15 years and above in Mizan-Aman Town, Southwest Ethiopia, 2016: a cross sectional study’, *International Journal of Endocrinology*, 2018. doi: 10.1155/2018/9317987.
- Babic, N. *et al.* (2019) ‘The triglyceride/HDL ratio and triglyceride glucose index as predictors of glycemic control in patients with diabetes mellitus type 2’, *Medical archives (Sarajevo, Bosnia and Herzegovina)*, 73(3), pp. 163–168. doi: 10.5455/medarh.2019.73.163-168.
- Badan Penelitian dan Pengembangan Kesehatan. (2013) ‘Riset Kesehatan Dasar (RISKESDAS) 2013’, *Laporan Nasional 2013*, pp. 1–384. doi: 1 Desember 2013.
- Badan Pengawas Obat dan Makanan. (2014) ‘Peraturan Kepala Badan Pengawas Obat dan Makanan Republik Indonesia Nomor 12 Tahun 2014 Tentang Pedoman Uji Klinik Obat Herbal’.
- Belwal, T. *et al.* (2017) ‘Dietary anthocyanins and insulin resistance: When food becomes a medicine’, *Nutrients*, 9(10). doi: 10.3390/nu9101111.
- Bhawani, S. A. *et al.* (2010) ‘Thin-layer chromatographic analysis of steroids: A review’, *Tropical Journal of Pharmaceutical Research*, 9(3), pp. 301–313. doi: 10.4314/tjpr.v9i3.56293.

- Brambilla, Paolo. Valle, EL. Falbo, Rosanna. Limonta, Giuseppe. Signorini, Stefano. Cappellini, Fabrizio. Mocarelli, P. (2011) 'Normal fasting plasma glucose and risk of type 2 diabetes', *Diabetes Care*, 34, pp. 1372–1374. doi: 10.2337/dc10-2263.
- Buchholz, T. and Melzig, M. F. (2016) 'Medicinal plants traditionally used for treatment of obesity and diabetes mellitus - screening for pancreatic lipase and  $\alpha$ -amylase inhibition', *Phytotherapy Research*, 30(2), pp. 260–266. doi: 10.1002/ptr.5525.
- Bulotta, A., Ludovico, O. and Coco, A. (2005) 'The common -866G/A polymorphism in the promoter region of the UCP-2 gene is associated with reduced risk of type 2 diabetes in Caucasians from Italy.', *Journal of Clinical Endocrinology and Metabolism*, 90(2), pp. 1176–1180.
- Bunbupha, S. *et al.* (2012) 'Antihyperinsulinemia effect of *Hibiscus sabdariffa* extract in high fructose diet induced insulin resistance rats methods objective proceeding', *Srinagarind Med J*, 27, pp. 4–7.
- Buraerah, H. (2010) 'Analisis faktor risiko diabetes melitus tipe 2 di Puskesmas Tanrutedong', *Medika: Jurnal Kedokteran Indoensia*, 35(4), p. 228.
- Cadenaro, M. *et al.* (2009) 'Effects of residual ethanol on the rate and degree of conversion of five experimental resins', *Dental Materials*, 25(5), pp. 621–628. doi: 10.1016/j.dental.2008.11.005.
- Carvajal-Zarrabal, O. *et al.* (2005) 'The consumption of *Hibiscus sabdariffa* dried calyx ethanolic extract reduced lipid profile in rats', *Plant Foods for Human Nutrition*, 60(4), pp. 153–159. doi: 10.1007/s11130-005-9023-x.
- Carvajal-Zarrabal, O. *et al.* (2009) 'Effect of *Hibiscus sabdariffa* L. dried calyx ethanol extract on fat absorption-excretion, and body weight implication in rats', *Journal of Biomedicine and Biotechnology*, 2009. doi: 10.1155/2009/394592.
- Cersosimo, E. *et al.* (2014) 'Assessment of pancreatic beta-cell function: review of methods and clinical applications', *Current Diabetes Reviews*, 10, pp. 2–42. doi: 10.2174/1573399810666140214093600.
- Cha, M. H. *et al.* (2007) 'Association of UCP2 and UCP3 gene polymorphisms with serum high-density lipoprotein cholesterol among Korean women', *Metabolism: Clinical and Experimental*, 56(6), pp. 806–813. doi: 10.1016/j.metabol.2007.01.023.
- Chan, C. B. and Harper, M.-E. (2006) 'Uncoupling proteins: role in insulin resistance and insulin insufficiency', *Current Diabetes Reviews*, 2(3), pp. 271–283. doi: 10.2174/157339906777950660.

- Chang-che Chew (2003) ‘*Hibiscus sabdariffa* Linn extract inhibits the development atherosclerosis in cholesterol-fed rabbit’, *Journal of Agricultural and Food Chemistry*, 51(18), p. 5472.
- Chang, H. C., Peng, C. H., Yeh, D. M., Kao, E. S., & Wang, C. J. (2014). *Hibiscus sabdariffa* extract inhibits obesity and fat accumulation, and improves liver steatosis in humans. *Food & function*, 5(4), 734–739. <https://doi.org/10.1039/c3fo60495k>
- Chen, C. C. *et al.* (2003) ‘*Hibiscus sabdariffa* extract inhibits the development of atherosclerosis in cholesterol-fed rabbits’, *Journal of Agricultural and Food Chemistry*, 51(18), pp. 5472–5477. doi: 10.1021/jf030065w.
- Chen, C. C. *et al.* (2004) ‘Inhibitory effects of *Hibiscus sabdariffa* L extract on low-density lipoprotein oxidation and anti-hyperlipidemia in fructose-fed and cholesterol-fed rats’, *Journal of the Science of Food and Agriculture*, 84(15), pp. 1989–1996. doi: 10.1002/jsfa.1872.
- Cheurfa, N. *et al.* (2008) ‘The common-866 G>A variant in the promoter of UCP2 is associated with decreased risk of coronary artery disease in type 2 diabetic men’, *Diabetes*, 57(4), pp. 1063–1068. doi: 10.2337/db07-1292.
- Chuenta, S. *et al.* (2011) ‘The effect of Roselle extract on insulin resistance , adiponectin secretion and fat tissue weight in high fructose-high fat diet induced insulin resistant rats methods proceeding results proceeding’, in *Methods Proceeding Results Proceeding*, pp. 25–28.
- Cissé, M. *et al.* (2009) ‘The bissap (*Hibiscus sabdariffa* L.): composition and principal uses’, *Fruits*, 64(3), pp. 179–193.
- D’Adamo, M. *et al.* (2004) ‘The -866 G/A genotype in the promoter of the human uncoupling protein 2 gene is associated with insulin resistance and increased risk of type 2 diabetes’, *Diabetes*, 53(7), pp. 1905–1910. doi: 10.2337/diabetes.53.7.1905.
- Da-Costa-Rocha, I. *et al.* (2014) ‘*Hibiscus sabdariffa* L-a phytochemical and pharmacological review’, *Food Chemistry*, 165, pp. 424–443. doi: 10.1016/j.foodchem.2014.05.002.
- Dalgaard, L. T. *et al.* (1999) ‘An untranslated insertion variant in the uncoupling protein 2 gene is not related to body mass index and changes in body weight during a 26-year follow-up in Danish Caucasian men’, *Diabetologia*, 42(12), pp. 1413–1416. doi: 10.1007/s001250051312.
- Dalgaard, L. T. *et al.* (2003) ‘Mutational analysis of the UCP2 core promoter and relationships of variants with obesity’, *Obesity Research*, 11(11), pp. 1420–1427. doi: 10.1038/oby.2003.191.

- Dalgaard, L. T. (2011) 'Genetik variance in uncoupling protein 2 in relation to obesity, type 2 diabetes, and related metabolic traits: Focus on the functional -866G>A promoter variant (rs659366)', *Journal of Obesity*, 2011. doi: 10.1155/2011/340241.
- Dalgaard, L. T. and Pedersen, O. (2001) 'Uncoupling proteins: functional characteristics and role in the pathogenesis of obesity and type II diabetes', *Diabetologia*, 44(8), pp. 946–965. doi: 10.1007/s001250100596.
- Davey, P. (2014). *Medicine at a Glance*. 4th edn. Oxford: John Wiley and Sons.
- Decroli, E. (2019) *Diabetes Mellitus tipe 2*. Padang: Pusat Penerbitan Penyakit Dalam. Available at: <http://aulanni.lecture.ub.ac.id/files/2012/04/MAKALAH-DIABETES-MELITUS-TIPE-I.pdf>.
- Depkes RI (1995). *Farmakope Indonesia*.
- Depkes RI (2005). *Pharmaceutical care untuk penyakit diabetes mellitus*. Jakarta: Direktorat Bina Farmasi Komunitas dan Klinik Direktorat Jenderal Bina Kefarmasian dan Alat Kesehatan.
- Depkes RI (2009) *Profil Kesehatan Indonesia*.
- Dhamrait, S. S. *et al.* (2004) 'FT Cardiovascular risk in healthy men and markers of oxidative stress in diabetic men are associated with common variation in the gene for uncoupling protein 2', *European Heart Journal*, 25(6), pp. 468–475. doi: 10.1016/j.ehj.2004.01.007.
- Diabetes UK (2010). *Key Statistic on Diabetes*.
- Drew, B. G. *et al.* (2009) 'High-density lipoprotein modulates glucose metabolism in patients with type 2 diabetes mellitus', *Circulation*, 119(15), pp. 2103–2111. doi: 10.1161/CIRCULATIONAHA.108.843219.
- Duangjai, A., Ingkaninan, K. and Limpeanchob, N. (2011) 'Potential mechanisms of hypocholesterolaemic effect of Thai spices/dietary extracts', *Natural Product Research*, 25(4), pp. 341–352. doi: 10.1080/14786411003754249.
- Duke, J. A. (1983) *Hanbook of energy crops*.
- Essa, M. M. and Subramanian, P. (2007) 'Hibiscus sabdariffa affects ammonium chloride-induced hyperammonemic rats', *Evidence-based Complementary and Alternative Medicine*, 4(3), pp. 321–325. doi: 10.1093/ecam/nel087.
- Esteghamati, A. *et al.* (2010) 'Optimal cut-off of homeostasis model assessment of insulin resistance (HOMA-IR) for the diagnosis of metabolic syndrome: Third national surveillance of risk factors of non-communicable diseases in Iran (SuRFNCD-2007)', *Nutrition and Metabolism*, 7. doi: 10.1186/1743-7075-7-26.

- Esterbauer, H. *et al.* (2001) 'A common polymorphism in the promoter of UCP2 is associated with decreased risk of obesity in middle-aged humans', *Nature Genetiks*, 28(2), pp. 178–183. doi: 10.1038/88911.
- Farombi, E. O. and Ige, O. O. (2007) 'Hypolipidemic and antioxidant effects of ethanolic extract from dried calyx of *Hibiscus sabdariffa* in alloxan-induced diabetic rats', *Fundamental and Clinical Pharmacology*, 21(6), pp. 601–609. doi: 10.1111/j.1472-8206.2007.00525.x.
- Farooq, M. *et al.* (2017) 'Correlation between serum glucose and BMI in undergraduate students', *Pakistan Journal of Medical and Health Sciences*, 11(3), pp. 887–889.
- Fatimah, R. N. (2015) 'Diabetes Melitus Tipe 2', *Fakultas Kedokteran Universitas Lampung*, 4, pp. 93–101. doi: 10.2337/dc12-0698.
- Fatmawati, N., Lyrawati, D. and Yurina, V. (2013) 'Effect of alpha lipoic acid on oxidative stress in hearth of male wistar rat with type II diabetes mellitus model induced by Streptozotocin.', in *6th Asian Association of Schools of Pharmacy Conference Program and Abstracts Book*. Singapore, p. 183.
- Feller, S., Boeing, H. and Pischon, T. (2010) 'Body mass index, waist circumference, and the risk of type 2 diabetes mellitus', *Deutsches Arzteblatt Online*, 107(26), pp. 470–476. doi: 10.3238/arztebl.2010.0470.
- Fischer, A., Fisher, E., Möhlig, M., Schulze, M., Hoffmann, K., Weickert, M. O., Schueler, R., Osterhoff, M., Pfeiffer, A. F., Boeing, H., & Spranger, J. (2008). KCNJ11 E23K affects diabetes risk and is associated with the disposition index: results of two independent German cohorts. *Diabetes care*, 31(1), 87–89. <https://doi.org/10.2337/dc07-1157>
- Florez, J. C., Burtt, N., de Bakker, P. I., Almgren, P., Tuomi, T., Holmkvist, J., Gaudet, D., Hudson, T. J., Schaffner, S. F., Daly, M. J., Hirschhorn, J. N., Groop, L., & Altshuler, D. (2004). Haplotype structure and genotype-phenotype correlations of the sulfonylurea receptor and the islet ATP-sensitive potassium channel gene region. *Diabetes*, 53(5), 1360–1368. <https://doi.org/10.2337/diabetes.53.5.1360>
- Le Fur, S. *et al.* (2004) 'The common - 866 G/A polymorphism in the promoter of uncoupling protein 2 is associated with increased carbohydrate and decreased lipid oxidation in juvenile obesity', *Diabetes*, 53(1), pp. 235–239. doi: 10.2337/diabetes.53.1.235.
- Gable, D. R. *et al.* (2006) 'Variation in the UCP2-UCP3 gene cluster predicts the development of type 2 diabetes in healthy middle-aged men', *Diabetes*, 55(5), pp. 1504–1511. doi: 10.2337/db05-1645.
- Gafoor, K. A. and Ashraf, M. (2006) 'Academic self efficacy scale'.

- Gagnier JJ, Boonc H, Rochona P, *et al.* (2006). Recommendations for reporting randomized controlled trials of herbal interventions: explanation and elaboration. *J Clin Epidemiol*, 59:1134-1149
- Gayoso-Diz, P. *et al.* (2013) 'Insulin resistance (HOMA-IR) cut-off values and the metabolic syndrome in a general adult population: Effect of gender and age: EPIRCE cross-sectional study', *BMC Endocrine Disorders*, 13. doi: 10.1186/1472-6823-13-47.
- Gloyn, A. and McCarthy, M. (2001) 'The genetiks of type 2 diabetes', *Best Practice dan Research Clinical Endocrinology dan Metabolism*, 15(3), pp. 293–308.
- Gordis L. (2014). *Epidemiology*, 5<sup>th</sup> ed. Philadelphia: Elsevier Saunders
- Gozel, N. and Dalkilic, S. (2017) 'UCP2 866 G/A gene (rs659366) polymorphism associated with diabetes type 2 in Turkish population', *Progress in Nutrition*, 19(4), pp. 430–435.
- Gulam, A. H., CF Otieno, F. and Oyoo, G. O. (2017) 'Prevalence of psychological insulin resistance among patients with type 2 diabetes at Kenyatta National Hospital, Kenya', *Health Science Journal*, 11(3), pp. 1–8. doi: 10.21767/1791-809x.1000508.
- Gurrola-Díaz, C. M. *et al.* (2010) 'Effects of *Hibiscus sabdariffa* extract powder and preventive treatment (diet) on the lipid profiles of patients with metabolic syndrome (MeSy)', *Phytomedicine*, 17(7), pp. 500–505. doi: 10.1016/j.phymed.2009.10.014.
- H. Syamsuni. (2006). *Farmasetika dasar dan hitungan farmasi*. 1st edn. Edited by W. R. Syarief. Jakarta: Penerbit Buku Kedokteran EGC Jakarta.
- Halton, T. L. *et al.* (2008) 'Low-carbohydrate-diet score and risk of type 2 diabetes in women', *American Journal of Clinical Nutrition*, 87(2), pp. 339–346. doi: 10.1093/ajcn/87.2.339.
- Hanachi, P., Moghadam, R. H. and Latiffah, A. L. (2009) 'Investigation of lipid profiles and lipid peroxidation in patients with type 2 diabetes', *European Journal of Scientific Research*, 28(1), pp. 6–13.
- Harborne, J. . (1978). *Metode fitokimia. penuntun cara modern menganalisis tumbuhan*. Alih bahasa Kosasih Padmawinata., Bandung : ITB.
- Hardinsyah and Supariasa, I. D. N. (2016). *Ilmu Gizi : Teori dan aplikasi*. Jakarta: EGC.
- Hartono, A. (2004). *Terapi gizi dan diet rumah sakit*. Jakarta: Buku Kedokteran EGC.

- Hartwig, S. *et al.* (2015) ‘Association of change of anthropometric measurements with incident type 2 diabetes mellitus: a pooled analysis of the prospective population-based CARLA and SHIP Cohort studies’, *Medicine*, 94(34). doi: 10.1097/MD.0000000000001394.
- Haryadi, N. K. (2011). *Kelor herbal multikhasiat ampuh melawan diabetes mellitus, kolesterol tinggi dan penyakit lainnya*. Surakarta: Delta Medika.
- Herranz-López, M. *et al.* (2017) ‘Multi-targeted molecular effects of *Hibiscus sabdariffa* polyphenols: An opportunity for a global approach to obesity’, *Nutrients*, 9(8). doi: 10.3390/nu9080907.
- Hirunpanich, V. *et al.* (2005) ‘Antioxidant effects of aqueous extracts from dried calyx of *hibiscus sabdariffa* Linn. (roselle) in vitro using rat low-density lipoprotein (LDL)’, *Biological and Pharmaceutical Bulletin*, 28(3), pp. 481–484. doi: 10.1248/bpb.28.481.
- Hong, M. *et al.* (2019) ‘Contribution and interaction of the low-density lipoprotein cholesterol to high-density lipoprotein cholesterol ratio and triglyceride to diabetes in hypertensive patients: A cross-sectional study’, *Journal of Diabetes Investigation*, 10(1), pp. 131–138. doi: 10.1111/jdi.12856.
- Hong, S. *et al.* (2010) ‘Development and validation of a semi-quantitative food frequency questionnaire to assess diets of Korean type 2 diabetic patients’, *Korean Diabetes Journal*, 34(1), p. 32. doi: 10.4093/kdj.2010.34.1.32.
- Hopkins, A. L. *et al.* (2013) ‘*Hibiscus sabdariffa* L. in the treatment of hypertension and hyperlipidemia: A comprehensive review of animal and human studies’, *Fitoterapia*, 85(1), pp. 84–94. doi: 10.1016/j.fitote.2013.01.003.
- Hou, G. *et al.* (2020) ‘UCP2–866G/A polymorphism is associated with prediabetes and type 2 diabetes’, *Archives of Medical Research*, 51(6), pp. 556–563. doi: 10.1016/j.arcmed.2020.06.001.
- Hsu, Y. *et al.* (2008) ‘Genetik variants in the UCP2-UCP3 gene cluster and risk of diabetes in the Women’s Health Initiative Observational Study’, 57(April). doi: 10.2337/db07-1269.Additional.
- Hsu, Y. H. *et al.* (2008) ‘Genetik variants in the UCP2-UCP3 gene cluster and risk of diabetes in the women’s health initiative observational study’, *Diabetes*, 57(4), pp. 1101–1107. doi: 10.2337/db07-1269.
- Huang, C.-N. *et al.* (2016) ‘*Hibiscus sabdariffa* polyphenols prevent palmitate-induced renal epithelial mesenchymal transition by alleviating dipeptidyl peptidase-4-mediated insulin resistance’, *Food Funct.*, 7(1), pp. 475–482. doi: 10.1039/C5FO00464K.

- Huang, C. N. *et al.* (2016) ‘Hibiscus sabdariffa polyphenols prevent palmitate-induced renal epithelial mesenchymal transition by alleviating dipeptidyl peptidase-4-mediated insulin resistance’, *Food and Function*, 7(1), pp. 475–482. doi: 10.1039/c5fo00464k.
- Huaysrichan, W. *et al.* (2017) ‘Hibiscus sabdariffa linn. calyx extraction and gallic acid improving cardiac diastolic dysfunction in high fat diet-STZ-induced type 2 diabetic rats’, *Asia-Pacific Journal of Science and Technology*, 22(2), pp. 1–10
- Huriyati, E. *et al.* (2016) ‘Dyslipidemia, insulin resistance and dietary fat intake in obese and normal weight adolescents: The role of uncoupling protein 2 - 866G/A gene polymorphism’, *International Journal of Molecular Epidemiology and Genetics*, 7(1), pp. 67–73.
- Husna, E. and Ashra, F. (2014) ‘Kadar gula darah sewaktu pada penderita diabetes mellitus tipe 2 di wilayah kerja Puskesmas Air Tabit Kota Payakumbuh’, *Jurnal Kesehatan STIKes Prima Nusantara Bukittinggi*, Vol.6, 6(1), pp. 50–57.
- Husna, E. and Ashra, F. (2015) ‘Pengaruh pemberian teh Rosella terhadap penurunan kadar gula darah pada penderita diabetes mellitus tipe 2’, *Jurnal Kesehatan STIKes Prima Nusantara Bukittinggi*, Vol.6, 6(1). Available at: <http://perawat-home-care.blogspot.com/2010/11/pengaruh-pemberian-teh-Rosella-terhadap.html?m=1pdf>.
- IDAI. (2015). *Konsensus pengelolaan diabetes mellitus tipe 2*.
- Ike Sitoresmi M Purbowati, Khaswar Syamsu, Endang Warsiki, H. S. R. (2015) ‘Evaluasi toksisitas, aktivitas antibakteri dan antioksidan komponen bioaktif Rosela dengan variasi jenis pelarut’, *Journal of Agroindustrial Technology*, 25(2), pp. 182–189.
- Ilyas E. (2011). *Olahraga bagi diabetisi*. Jakarta: FK UI.
- Inuwa, I. *et al.* (2012) ‘Long-term ingestion of Hibiscus sabdariffa calyx extract enhances myocardial capillarization in the spontaneously hypertensive rat’, *Experimental Biology and Medicine*, 237(5), pp. 563–569. doi: 10.1258/ebm.2012.011357.
- IPAQ. (2005) ‘Guidelines For data processing and analysis of The International Physical Activity Questionnaire (IPAQ) Short and Long Forms’.
- Jain, H. R. *et al.* (2016) ‘A study of lipid profile in diabetes mellitus’, *International Journal of Scientific Study*, 4(4), pp. 56–61.

- Jeenduang, N. *et al.* (2017) 'APOE and CETP TaqIB polymorphisms influence metabolic responses to *Hibiscus sabdariffa* L. and *Gynostemma pentaphyllum* Makino tea consumption in hypercholesterolemic subjects', *Asia Pacific Journal of Clinical Nutrition*, 26(2), pp. 368–378. doi: 10.6133/apjcn.122015.04.
- Juliani, H. R. *et al.* (2009) 'Chemistry and quality of hibiscus (*hibiscus sabdariffa*) for developing the natural-product industry in senegal', *Journal of Food Science*, 74(2). doi: 10.1111/j.1750-3841.2009.01076.x.
- Kahn, B. B. and Flier, J. S. (2000) 'Obesity and insulin resistance', *Journal of Clinical Investigation*, 106(4), pp. 473–481. doi: 10.1007/978-1-59259-716-1\_4.
- Kang, H. M. and Kim, D. J. (2012) 'Body mass index and waist circumference according to glucose tolerance status in Korea: The 2005 Korean health and nutrition examination survey', *Journal of Korean Medical Science*, 27(5), pp. 518–524. doi: 10.3346/jkms.2012.27.5.518.
- Kao, E. S., Yang, M. Y., Hung, C. H., Huang, C. N., & Wang, C. J. (2016). Polyphenolic extract from *Hibiscus sabdariffa* reduces body fat by inhibiting hepatic lipogenesis and preadipocyte adipogenesis. *Food & function*, 7(1), 171–182. <https://doi.org/10.1039/c5fo00714c>
- Kariadi, S. H. (2009). *Panduan lengkap untuk diabetesi, keluarganya dan profesional medis*. Bandung: Qonita PT Mizan Pustaka.
- Kartinah, N. T. *et al.* (2019) 'The potential of *Hibiscus sabdariffa* Linn. for treatment of obesity: focus on FGF21 in liver and adipose tissue', *e-Jurnal Kedokteran Indonesia Universitas Indonesia*, 7(2), pp. 131–136.
- Kaul, N. and Ali, S. (2016) 'Genes , genetics , and environment in type 2 diabetes : Implication in personalized medicine genes , genetics , and environment in type 2 diabetes ':, (October 2015). doi: 10.1089/dna.2015.2883.
- Kautzky-Willer, A., Harreiter, J. and Pacini, G. (2016) 'Sex and gender differences in risk, pathophysiology and complications of type 2 diabetes mellitus', *Endocrine Reviews*, 37(3), pp. 278–316. doi: 10.1210/er.2015-1137.
- Kemenkes (2018) *Riset Kesehatan Dasar 2018*. Jakarta.
- Keskin, M. *et al.* (2005) 'Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents', *Pediatrics*, 115(4). doi: 10.1542/peds.2004-1921.
- Kharal, P. M., Prasad, P. N. and Acharya, R. P. (2013) 'Gross correlation between waist hip ratio and blood sugar level in a village', *Journal of the Nepal Medical Association*, 52(6), pp. 361–364.

- El Khoury, D. and Hwalla, N. (2010) 'Metabolic and appetite hormone responses of hyperinsulinemic normoglycemic males to meals with varied macronutrient compositions', *Annals of Nutrition and Metabolism*, 57(1), pp. 59–67. doi: 10.1159/000317343.
- Kim, J. D. and Lee, W. Y. (2016) 'Insulin secretory capacity and insulin resistance in Korean type 2 diabetes mellitus patients', *Endocrinology and Metabolism*, 31(3), pp. 354–360. doi: 10.3803/EnM.2016.31.3.354.
- Kim, J. K. *et al.* (2007) 'Hibiscus sabdariffa L. water extract inhibits the adipocyte differentiation through the PI3-K and MAPK pathway', *Journal of Ethnopharmacology*, 114(2), pp. 260–267. doi: 10.1016/j.jep.2007.08.028.
- Kovacs, P. *et al.* (2005) 'Genetik variation in UCP2 (uncoupling protein-2) is associated with energy metabolism in Pima Indians', *Diabetologia*, 48(11), pp. 2292–2295. doi: 10.1007/s00125-005-1934-9.
- Krempler, F. *et al.* (2002) 'A functional polymorphism in the promoter of UCP2 enhances obesity risk but reduces type 2 diabetes risk in obese middle-aged humans', *Diabetes*, 51(11), pp. 3331–3335. doi: 10.2337/diabetes.51.11.3331.
- Kumar, A., Tewari, P. and Sahoo, S. S. (2005) 'Type-2 diabetes mellitus patients: a prospective study in North', *Indian Journal of Clinical Biochemistry*, 20(2), pp. 10–17.
- Labayen, I. *et al.* (2009) 'Association of common variants of UCP2 gene with low-grade inflammation in Swedish children and adolescents', *The European Youth Heart Study, Pediatric Research*, 66(3), pp. 350–354.
- Lapice, E. *et al.* (2010) 'Uncoupling protein 2 G(-866)A polymorphism: A new gene polymorphism associated with C-reactive protein in type 2 diabetic patients C-reactive protein in type 2 diabetic patients', *Cardiovascular Diabetology*, 9. doi: 10.1186/1475-2840-9-68.
- Lee, C. H. *et al.* (2016) 'Optimal cut-offs of homeostasis model assessment of insulin resistance (HOMA-IR) to identify dysglycemia and type 2 diabetes mellitus: A 15-year prospective study in Chinese', *PLoS ONE*, 11(9). doi: 10.1371/journal.pone.0163424.
- Lee, C. W. *et al.* (2012) 'Effects of hot air and freeze drying methods on physicochemical properties of citrus "hallabong" powders', *Food Science and Biotechnology*, 21(6), pp. 1633–1639. doi: 10.1007/s10068-012-0217-8.
- Lee, J., Durst, R. W. and Wrolstad, R. E. (2005) 'Determination of total monomeric anthocyanin pigment content of fruit juices, beverages, natural colorants, and wines by the pH differential method: Collaborative study', *Journal of AOAC International*, 88(5), pp. 1269–1278. doi: 10.1093/jaoac/88.5.1269.

- Lee, W. C. *et al.* (2009) ‘Polyphenol extracts from hibiscus sabdariffa linnaeus attenuate nephropathy in experimental type 1 diabetes’, *Journal of Agricultural and Food Chemistry*, 57(6), pp. 2206–2210. doi: 10.1021/jf802993s.
- Lembaga Ilmu Pengetahuan Indonesia (2004). *Widyakarya Nasional Pangan dan Gizi VIII*.
- Li, J. *et al.* (2016) ‘Dietary protein intake and type 2 diabetes among women and men in Northeast China’, *Scientific Reports*, 6. doi: 10.1038/srep37604.
- Lin, D. *et al.* (2016) ‘An overview of plant phenolic compounds and their importance in human nutrition and management of type 2 diabetes’, *Molecules*, 21(10). doi: 10.3390/molecules21101374.
- Lin, H. H. *et al.* (2007) ‘Chemopreventive properties of Hibiscus sabdariffa L. on human gastric carcinoma cells through apoptosis induction and JNK/p38 MAPK signaling activation’, *Chemico-Biological Interactions*, 165(1), pp. 59–75. doi: 10.1016/j.cbi.2006.10.011.
- Linn, T. *et al.* (2000) ‘Effect of long-term dietary protein intake on glucose metabolism in humans’, *Diabetologia*, 43(10), pp. 1257–1265. doi: 10.1007/s001250051521.
- Liu, J. *et al.* (2013) ‘The role of uncoupling proteins in diabetes mellitus’, *Journal of Diabetes Research*, vol 2013, pp.1-8. <https://www.hindawi.com/journals/jdr>  
<https://doi.org/10.1155/2013/585897>
- Lo, C. W. *et al.* (2007) ‘Effect of Hibiscus anthocyanins-rich extract induces apoptosis of proliferating smooth muscle cell via activation of P38 MARK and p53 pathway’, *Molecular Nutrition and Food Research*, 51(12), pp. 1452–1460. doi: 10.1002/mnfr.200700151.
- López-Miranda, J. *et al.* (2007) ‘Dietary fat, genes and insulin sensitivity’, *Journal of Molecular Medicine*, 85(3), pp. 209–222. doi: 10.1007/s00109-006-0138-1.
- Lyssenko *et al.* (2008) ‘Clinical risk factors, DNA variants, and the development of type 2 diabetes’, *The New England Journal of Medicine*, 2008(359), pp. 2220–2232.
- Mahadik, S. *et al.* (2012) ‘Uncoupling protein-2 (UCP2) gene expression in subcutaneous and omental adipose tissue of Asian Indians: Relationship to adiponectin and parameters of metabolic syndrome’, *Adipocyte*, 1(2), pp. 101–107. doi: 10.4161/adip.19671.
- Mahan, K., Escott-Stump, S. and Raymond, J. (2012). *Krause’s food and the nutrition care process*. Philadelphia: Saunders.

- McCarthy M. I. (2010). Genomics, type 2 diabetes, and obesity. *The New England journal of medicine*, 363(24), 2339–2350. <https://doi.org/10.1056/NEJMra0906948>
- Mardiah *et al.* (2015) ‘Anti-inflammatory of purple Roselle extract in diabetic rats induced by streptozotocin’, *Procedia Food Science*, 3, pp. 182–189. doi: 10.1016/j.profoo.2015.01.020.
- Mardiah, S., H., R. W. A. and Rahayu, A. (2009). *Budi daya dan pengolahan Rosella si merah segudang manfaat*. Jakarta: Agromedia Pustaka.
- Maryani, H. (2005) *Khasiat dan manfaat Rosella*. Jakarta: Agromedia Pustaka.
- Maturin, M. and Peeler, J. . (2001) ‘Chapter 3. Aerobic Plate Count’, in *Food and Drug Administration (FDA), bacteriological analytical manual online*. 8th edn. Berlin: Silver Spring.
- Mayasari, E., Nosi, H. and Zainal, S. (2014) ‘Faktor yang berhubungan dengan kepatuhan klien diabetes melitus dalam mengontrol gula darah di Poliklinik Interna RSUD Labuang Baji Makassar’, *Jurnal Ilmiah Kesehatan*, 5(5), pp. 568–573.
- Mayasari, N. R. *et al.* (2018) ‘Antidiabetic effect of Rosella-Stevia Tea on Prediabetic women in Yogyakarta, Indonesia’, *Journal of the American College of Nutrition*, 37(5), pp. 373–379. doi: 10.1080/07315724.2017.1400927.
- McKay, D. L. *et al.* (2010) ‘Hibiscus Sabdariffa L. tea (tisane) lowers blood pressure in prehypertensive and mildly hypertensive adults’, *Journal of Nutrition*, 140(2), pp. 298–303. doi: 10.3945/jn.109.115097.
- Md Idris, M. H. *et al.* (2012) ‘Protective role of Hibiscus sabdariffa calyx extract against streptozotocin induced sperm damage in diabetic rats’, *EXCLI Journal*, 11, pp. 659–669. doi: 10.17877/DE290R-5145.
- Meigs, J. B. (2003) ‘Epidemiology of the insulin resistance syndrome’, *Current Diabetes Reports*, 3(1), pp. 73–79. doi: 10.1002/0470846429.ch20.
- Merentek, E. (2006) Resistensi insulin pada diabetes melitus tipe 2, *Cermin Dunia Kedokteran*. Jakarta: Cernin Dunia Kedokteran.
- Mizokami-Stout, K., Cree-Green, M., & Nadeau, K. J. (2012). Insulin resistance in type 2 diabetic youth. *Current opinion in endocrinology, diabetes, and obesity*, 19(4), 255–262. <https://doi.org/10.1097/MED.0b013e3283557cd5>
- Misra, A., & Khurana, L. (2008). Obesity and the metabolic syndrome in developing countries. *The Journal of clinical endocrinology and metabolism*, 93(11 Suppl 1), S9–S30. <https://doi.org/10.1210/jc.2008-1595>

- Mohamed, B. B., Sulaiman, A. A. and Dahab, A. A. (2012) 'Roselle ( *Hibiscus sabdariffa* L.) in Sudan , cultivation and Their uses', *Bull. Environ. Pharmacol. Life Sci*, 1(6), pp. 48–54.
- Mohamed, J. *et al.* (2013) 'The protective effect of aqueous extracts of roselle (*Hibiscus sabdariffa* L. UKMR-2) against red blood cell membrane oxidative stress in rats with streptozotocin-induced diabetes', *Clinics*, 68(10), pp. 1358–1363. doi: 10.6061/clinics/2013(10)11.
- Mohd-Esa, N. *et al.* (2010) 'Antioxidant activity in different parts of roselle (*Hibiscus sabdariffa* L.) extracts and potential exploitation of the seeds', *Food Chemistry*, 122(4), pp. 1055–1060. doi: 10.1016/j.foodchem.2010.03.074.
- Morten, M. and Gail Vance, C. (1999). *Sensory evaluation technique*. 3rd edn. Raton Florida: CRC Press Boca.
- Morton, J. F. (1987). *Fruits of warm climates. Roselle Hibiscus sabdariffa L., Purdue University*. Miami Florida. Available at: <https://hort.purdue.edu/newcrop/morton/roselle.html>.
- Mozaffari-Khosravi, H. *et al.* (2009) 'Effects of sour tea (*Hibiscus sabdariffa*) on lipid profile and lipoproteins in patients with type II diabetes', *Journal of Alternative and Complementary Medicine*, 15(8), pp. 899–903. doi: 10.1089/acm.2008.0540.
- Mozaffari-Khosravi, H., Ahadi, Z. and Fallah Tafti, M. (2014) 'The effect of Green Tea versus Sour Tea on insulin resistance, lipids profiles and oxidative stress in patients with type 2 diabetes mellitus: A randomized clinical trial', *Iranian J Med Sci*, 39(5), pp. 424–432. doi: 10.3109/19390211.2013.790333.
- Nafizah, A. H. N. *et al.* (2017) 'Aqueous calyces extract of Roselle or *Hibiscus sabdariffa* Linn supplementation improves liver morphology in streptozotocin induced diabetic rats', *Arab Journal of Gastroenterology*, 18(1).
- Ndraha, Suzanna. (2014). Diabetes melitus tipe 2 dan tata laksana terkini. *Scientific Journal of Pharmaceutical Development and Medical Application*, Vol. 27, No.2, Agustus 2014. MEDICINUS 1
- Nerdy (2015) 'In silico docking of chemical compounds from roselle calyces (*Hibiscus sabdariffa* L.) as antidiabetic', *International Journal of ChemTech Research*, 7(1), pp. 148–152.
- Nnam, N. M. and Onyeke, N. G. (2003) 'Chemical composition of two varieties of sorrel (*Hibiscus sabdariffa* L.), calyces and the drinks made from them', *Plant Foods for Human Nutrition*, 58(3), pp. 1–7. doi: 10.1023/B:QUAL.0000040310.80938.53.

- Nnamonu EI, Ejere VC, Ejim AO, Echi PC, Egbuji JV, Eze TR, Eyo JE. (2013). Effects of *Hibiscus sabdariffa* calyces aqueous extract on serum cholesterol, body weight and liver biomarkers of *Rattus norvegicus*. *International Journal of Indigenous Medicinal Plants* 46 (4):1405-1411
- Nurcahyo, H. and Febriani, R. (2019) 'Potensi bunga Rosela (*Hibiscus sabdariffa* Linn.) sebagai bahan pangan fungsional, sumber pigmen dan antioksidan alami', *Seminar Nasional INAHCO (Indonesian Anemia dan Health Conference) 2019 ISBN*, pp. 192–196.
- Nurkhasanah (2015) 'The effect of Rosella (*Hibiscus Sabdariffa* L) treatment on IL-10 and IL-14 secretion on dimethylbenz (A) anthracene (DMBA) induced rat', *International Journal of Pharmacy and Pharmaceutical Sciences*, 7(4), pp. 402–404.
- Odigie, I. P., Ettarh, R. R. and Adigun, S. A. (2003) 'Chronic administration of aqueous extract of *Hibiscus sabdariffa* attenuates hypertension and reverses cardiac hypertrophy in 2K-1C hypertensive rats', *Journal of Ethnopharmacology*, 86(2–3), pp. 181–185. doi: 10.1016/S0378-8741(03)00078-3.
- Ojeda, D. *et al.* (2010) 'Inhibition of angiotensin convertin enzyme (ACE) activity by the anthocyanins delphinidin- and cyanidin-3-O-sambubiosides from *Hibiscus sabdariffa*', *Journal of Ethnopharmacology*, 127(1), pp. 7–10. doi: 10.1016/j.jep.2009.09.059.
- Ojewumi, A. W. and Kadiri, M. (2013) 'Physiological evaluation of the anti-diabetic properties of *Hibiscus sabdariffa* on rats', *Journal of Natural Science, Engineering and Technology*, 12, pp. 50-61
- Ojulari, O. V., Lee, S. G., & Nam, J. O. (2019). Beneficial effects of natural bioactive compounds from *Hibiscus sabdariffa* L. on Obesity. *Molecules (Basel, Switzerland)*, 24(1), 210. <https://doi.org/10.3390/molecules24010210>
- Okereke, C. N., Iroka, F. and Chukwuma, M. (2015) 'Phytochemical analysis and medicinal uses of *Hibiscus sabdariffa*', *International Journal of Herbal Medicine*, 2(6), pp. 16–19. Available at: <https://www.researchgate.net/publication/304157698>.
- Olaleye, M. T. (2007) 'Cytotoxicity and antibacterial activity of Methanolic extract of *Hibiscus sabdariffa*', *Journal of Medicinal Plants Research*, 1(1), pp. 009–013.
- Omiya, K. *et al.* (2015) 'Impaired  $\beta$ -cell function attenuates training effects by reducing the increase in heart rate reserve in patients with myocardial infarction', *Journal of Cardiology*, 65(2), pp. 128–133. doi: 10.1016/j.jjcc.2014.04.012.

- Onyenekwe, P. C. *et al.* (1999) 'Antihypertensive effect of Roselle (*Hibiscus sabdariffa*) calyx infusion in spontaneously hypertensive rats and a comparison of its toxicity with that in Wistar rats', *Cell Biochemistry and Function*, 17(3), pp. 199–206. doi: 10.1002/(SICI)1099-0844(199909)17:3<199::AID-CBF829>3.0.CO;2-2.
- Palmer, B. R. *et al.* (2009) 'The common G-866A polymorphism of the UCP2 gene and survival in diabetic patients following myocardial infarction', *Cardiovascular Diabetology*, 8. doi: 10.1186/1475-2840-8-31.
- Paramitha, G. M. (2014) 'Hubungan aktivitas fisik dengan kadar gula darah pada pasien dm tipe 2 di Rumah Sakit Umum Daerah Karanganyar', *Universitas Muhammadiyah Surakarta*, pp. 1–14.
- Peng, C. H. *et al.* (2011) 'Hibiscus sabdariffa polyphenolic extract inhibits hyperglycemia, hyperlipidemia, and glycation-oxidative stress while improving insulin resistance', *Journal of Agricultural and Food Chemistry*, 59(18), pp. 9901–9909. doi: 10.1021/jf2022379.
- PERKENI. (2006). *Konsensus pengelolaan dan pencegahan diabetes mellitus tipe 2 di Indonesia*. Jakarta: PB Perkeni.
- PERKENI. (2015). *Konsensus pengelolaan dan pencegahan diabetes mellitus tipe 2 di indonesia pengurus besar endokrinologi indonesia*, *Jurnal Konsensus*. Jakarta: PB Perkeni.
- Peter, E. *et al.* (2014) 'Iron and ascorbic acid content in *Hibiscus sabdariffa* calyces in Tanzania: modeling and optimization of extraction conditions', *International Journal of Food Science and Nutrition Engineering*, 4(2), pp. 27–35. doi: 10.5923/j.food.20140402.01.
- Pierik, R. (1988). *In vitro culture of higher plants*, *Photosynthesis Research*. London: Martinus Nijhoff Publisher. doi: 10.1007/bf00035455.
- Pocock (2008). *Clinical trial. A practical approach*. New York: A Willey Medical Publication.
- Pratiwi, A. T. (2018) 'Pengaruh ekstrak Rosella (*Hibiscus sabdariffa* Linn) dalam menurunkan kadar gula darah untuk diabetes melitus', *Jurnal Ilmiah Mahasiswa Kedokteran Indonesia*, 6(2).
- Quispe, R. *et al.* (2015) 'Relationship of the triglyceride to high-density lipoprotein cholesterol (TG/HDL-C) ratio to the remainder of the lipid profile: The very large database of lipids-4 (VLDL-4) study', *Atherosclerosis*, 242(1), pp. 243–250. doi: 10.1016/j.atherosclerosis.2015.06.057.

- Quispe, R., Martin, S. S. and Jones, S. R. (2016) 'Triglycerides to high-density lipoprotein-cholesterol ratio, glycemic control and cardiovascular risk in obese patients with type 2 diabetes', *Current Opinion in Endocrinology, Diabetes and Obesity*, 23(2), pp. 150–156. doi: 10.1097/MED.0000000000000241.
- Rai, E. *et al.* (2007) 'Interaction between the UCP2-866G/A, mtDNA 10398G/A and PGC1 $\alpha$  p.Thr394Thr and p.Gly482Ser polymorphisms in type 2 diabetes susceptibility in North Indian population', *Human Genetiks*, 122(5), pp. 535–540. doi: 10.1007/s00439-007-0421-4.
- Reynolds, K. and He, J. (2005) 'Epidemiology of metabolic syndrome', *The American Journal of the Medical Science*, 330, pp. 273–9. doi: 10.1002/9780470779750.ch4.
- Rietman, A. *et al.* (2014) 'High dietary protein intake, reducing or eliciting insulin resistance?', *European Journal of Clinical Nutrition*, 68(9), pp. 973–979. doi: 10.1038/ejcn.2014.123.
- Rosemary, Rosidah and Haro, G. (2014) 'Antidiabetic effect of Roselle calyces extract (*Hibiscus Sabdariffa* L.) in streptozotocin induced mice', *International Journal of PharmTech Research*, 6(5), pp. 1703–1711.
- Ross, I. A. (2003). *Medicinal plants of the world*. 2nd edn, New Jersey: Humana Press. doi: 10.1007/978-1-59259-365-1.
- Rudofsky, G. (2006) 'Functional polymorphisms of UCP2 and UCP3 are associated with a reduced prevalence of diabetic neuropathy in patients with type 1 diabetes', *Diabetes Care*, 29(1), pp. 89–94. doi: 10.2337/diacare.29.1.89.
- Rütti, S. *et al.* (2009) 'Low- and high-density lipoproteins modulate function, apoptosis, and proliferation of primary human and murine pancreatic  $\beta$ -cells', *Endocrinology*, 150(10), pp. 4521–4530. doi: 10.1210/en.2009-0252.
- Sabin, M. A. *et al.* (2018) 'Insulin and BMI as predictors of adult type 2 diabetes mellitus', *Pediatrics*, 135(1). doi: 10.1542/peds.2014-1534.
- Salopuro, T. *et al.* (2009) 'Variation in the UCP2 and UCP3 genes associates with abdominal obesity and serum lipids: The Finnish diabetes prevention study', *BMC Medical Genetiks*, 10, p. 94. doi: 10.1186/1471-2350-10-94.
- Sari (2016) *Toksistas ekut ekstrak etanol kelopak Rosella (Hibiscus Sabdariffa L.) dan pengaruhnya terhadap fungsi hepar*. Universitas Ahmad Dahlan Yogyakarta.
- Sasahara, M. *et al.* (2004) 'Uncoupling protein 2 promoter polymorphism - 866G/A affects its expression in  $\beta$ -cells and modulates clinical profiles of Japanese type 2 diabetic patients', *Diabetes*, 53(2), pp. 482–485. doi: 10.2337/diabetes.53.2.482.

- Sastroasmoro, S. and Ismael, S. (2002). *Dasar-Dasar Metodologi Penelitian Klinis*. 2nd edn. Jakarta: Sagung Seto.
- Savjani Ketan, T., Gajjar Anuradha, K. and Savjani Jignasa, K. (2012) 'Drug solubility: importance and enhancement techniques', *ISRN pharmaceuticals*, 2012. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3399483/#B23>.
- Schumann, G. *et al.* (2005) 'IFCC primary reference procedures for the measurement of catalytic activity concentrations of enzymes at 37°C. part 5. reference procedure for the measurement of catalytic concentration of aspartate aminotransferase', *Clinical Chemistry and Laboratory Medicine*, 40(7). doi: 10.1515/cclm.2002.125.
- Serban, C. *et al.* (2015) 'Effect of sour tea (*Hibiscus sabdariffa* L.) on arterial hypertension', *Journal of Hypertension*, 33(6), pp. 1119–1127. doi: 10.1097/hjh.0000000000000585.
- Shadhan, R. *et al.* (2019) 'Hibiscus sabdariffa linn fruits methanolic extract and fractions mediated glucose uptake stimulation and glucose transporter 4 regulation', *Journal of Pharmaceutical Negative Results*, 10(1), pp. 25–31. doi: 10.4103/jpnr.JPNR\_14\_18.
- Shadhan, R. M. and Bohari, S. P. M. (2017) 'Effects of Hibiscus sabdariffa Linn. fruit extracts on  $\alpha$ -glucosidase enzyme, glucose diffusion and wound healing activities', *Asian Pacific Journal of Tropical Biomedicine*, 7(5), pp. 466–472. doi: 10.1016/j.apjtb.2017.01.023.
- Shah, N., Amanullah and Atallah, S. (2008) 'Prevalence of insulin resistance in type-2 diabetic patients', *JPMI - Journal of Postgraduate Medical Institute*, 22(1), pp. 52–56.
- Shen, X. (2007) 'Relationships of uncoupling protein 2 gene -866G/A polymorphism with metabolic syndrome in Chinese Han'.
- Shori, A. B. (2015) 'Screening of antidiabetic and antioxidant activities of medicinal plants', *Journal of Integrative Medicine*, 13(5), pp. 297–305. doi: 10.1016/S2095-4964(15)60193-5.
- Siddiqui, A. A. *et al.* (2013a) 'Diabetes: mechanism, pathophysiology and management-A review', *International Journal of Drug Development and Research*, 5(2), pp. 1–23.
- Siddiqui, A. A. *et al.* (2013b) 'Diabetes: mechanism, pathophysiology and management-A review', *International Journal of Drug Development and Research*, 5(2), pp. 1–23. doi: 10.2337/dc11-S011.

- Sindi, H. A., Marshall, L. J. and Morgan, M. R. A. (2014) 'Comparative chemical and biochemical analysis of extracts of *Hibiscus sabdariffa*', *Food Chemistry*, 164, pp. 23–29. doi: 10.1016/j.foodchem.2014.04.097.
- Singh, R. and Pannangpetch, P. (2017a) 'Roselle (*Hibiscus sabdariffa*) calyce ethanolic extract decreases insulin resistance in high carbohydrate diet fed cats', *Asia-Pacific Journal of Science and Technology*, 22(4), pp. 1–7.
- Singh, R. and Pannangpetch, P. (2017b) 'Roselle (*Hibiscus sabdariffa*) calyce ethanolic extract decreases insulin resistance in high carbohydrate diet fed cats', *Asia-Pacific Journal of Science and Technology*, 22(4), pp. 1–7.
- Singh, Y. *et al.* (2013) 'A study of insulin resistance by HOMA-IR and its cut-off value to identify metabolic syndrome in urban Indian adolescents', *JCRPE Journal of Clinical Research in Pediatric Endocrinology*, 5(4), pp. 245–251. doi: 10.4274/Jcrpe.1127.
- Singleton V L dan Rossi J A Jr (1985) 'Colorunetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents', *American Journal of Enology and Viticulture*, 16, pp. 144–158.
- Sireeratawong, S. *et al.* (2013) 'Toxicity studies of the water extract from the calyces of *Hibiscus sabdariffa* L. in rats', *African journal of traditional, complementary, and alternative medicines : AJTCAM*, 10(4), pp. 122–127. doi: 10.4314/ajtcam.v10i4.20.
- Soetiarto, F. and Suhardi, R. (2010) 'Hubungan indeks massa tubuh dan lingkar pinggang: Data Riskesdas 2007', *Buletin Penelitian Kesehatan*, 38, pp. 36–42.
- Souza, B. M. de *et al.* (2011) 'The role of uncoupling protein 2 (UCP2) on the development of type 2 diabetes mellitus and its chronic complications', *Arq. Bras Endocrinol Metab*, 55(4), pp. 239–48.
- Srihardyastutie, A. *et al.* (2014) 'The relationship between HbA1c, insulin resistance and changes of insulin secretion in Indonesian type 2 diabetic subjects', *Adv. in Nat. Appl. Sci.*, 8(8), pp. 25–30.
- Stephens, J. W. *et al.* (2008) 'Interaction between the uncoupling protein 2 - 866G>A gene variant and cigarette smoking to increase oxidative stress in subjects with diabetes', *Nutrition, Metabolism and Cardiovascular Diseases*, 18(1), pp. 7–14. doi: 10.1016/j.numecd.2007.01.010.
- Sudargo, T. *et al.* (2016) 'The relationship between fried food consumption and physical activity with diabetes mellitus in Yogyakarta, Indonesia', *International Journal Of Community Medicine And Public Health*, 4(1), p. 38. doi: 10.18203/2394-6040.ijcmph20164709.

- Sudaryanto, Agus. (2014) 'Hubungan antara pola makan, genetik dan kebiasaan olahraga terhadap kejadian diabetes melitus tipe II DI Wilayah Kerja Puskesmas Nusukan, Banjarsari', in *Universitas Muhammadiyah Surakarta*, pp. 19–24.
- Suiraka. (2012) *Penyebab penyakit degeneratif*. Yogyakarta: Nuha Medika.
- Sumitro, R. (2011) *Pengaruh pemberian seduhan bunga Rosella pada perubahan tekanan darah sistolik dan diastolik*, Universitas Airlangga: Surabaya. Surabaya: Universitas Airlangga.
- Surapon Tangvarasittichai (2015) 'DM (oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus)', *World J Diabetes*, Apr 15, 20, pp. 456–480. doi: <http://dx.doi.org/10.4239/wjd.v6.i3.456>.
- T.H., T. (1997) 'Protective effect of dried flower extract of *Hibiscus Sabdariffa* Linn againsts oxidative stress in rat primery hepatocytes', *Food Chemical Toxicology*, 35(12), pp. 1159–1164.
- Tandra, H. (2013) *Life healthy with diabetes mellitus*. Yogyakarta: Rapha Publishing.
- Tangvarasittichai, S. (2015) 'Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus', *World Journal of Diabetes*, 6(3), p. 456. doi: [10.4239/wjd.v6.i3.456](http://dx.doi.org/10.4239/wjd.v6.i3.456).
- Timóteo, A. T. *et al.* (2014) 'Optimal cut-off value for homeostasis model assessment (HOMA) index of insulin-resistance in a population of patients admitted electively in a Portuguese cardiology ward', *Acta Medica Portuguesa*, 27(4), pp. 473–479. doi: [10.20344/amp.5180](https://doi.org/10.20344/amp.5180).
- Trisnawati, S. (2013) 'Faktor risiko kejadian diabetes melitus tipe II Di Puskesmas Kecamatan Cengkareng Jakarta Barat', *Jurnal Ilmiah Kesehatan*, 5(1), pp. 6–11.
- Villalpando-Arteaga, E. V., Mendieta-Condado, E., Esquivel-Solís, H., Canales-Aguirre, A. A., Gálvez-Gastélum, F. J., Mateos-Díaz, J. C., Rodríguez-González, J. A., & Márquez-Aguirre, A. L. (2013). *Hibiscus sabdariffa* L. aqueous extract attenuates hepatic steatosis through down-regulation of PPAR- $\gamma$  and SREBP-1c in diet-induced obese mice. *Food & function*, 4(4), 618–626. <https://doi.org/10.1039/c3fo30270a>
- Urhammer, S. A. *et al.* (1997) 'Mutational analysis of the coding region of the uncoupling protein 2 gene in obese NIDDM patients: Impact of a common amino acid polymorphism on juvenile and maturity onset forms of obesity and insulin resistance', *Diabetologia*, 40(10), pp. 1227–1230. doi: [10.1007/s001250050811](https://doi.org/10.1007/s001250050811).

- Utari, N. M. *et al.* (2014) 'Astaxanthin menurunkan kadar vascular endothelial growth factor , level on mild nonproliferative diabetic retinopathy ', *Medicina Journal*, 45(1), pp. 31–37.
- Vimaleswaran, K. S. *et al.* (2011) 'Uncoupling protein 2 and 3 gene polymorphisms and their association with type 2 diabetes in Asian Indians', *Diabetes Technology and Therapeutics*, 13(1), pp. 19–25. doi: 10.1089/dia.2010.0091.
- Wang, H. *et al.* (2004) 'Uncoupling protein-2 polymorphisms in type 2 diabetes, obesity, and insulin secretion', *American Journal of Physiology - Endocrinology and Metabolism*, 286(1 49-1). doi: 10.1152/ajpendo.00231.2003.
- Wang, J. (2000). *Analytical chemistry*. New York: Willey-VCH.
- Wang, M. L. *et al.* (2012) 'Assessment of oil content and fatty acid composition variability in two economically important Hibiscus species', *Journal of Agricultural and Food Chemistry*, 60(26), pp. 6620–6626. doi: 10.1021/jf301654y.
- Wang, P. W. *et al.* (2009) 'Mitochondrial DNA variants in the pathogenesis of type 2 diabetes - relevance of Asian population studies', *Review of Diabetic Studies*, 6(4), pp. 237–246. doi: 10.1900/RDS.2009.6.237.
- Werdani, A. R. and Triyanti, T. (2014) 'Asupan karbohidrat sebagai faktor dominan yang berhubungan dengan kadar gula darah puasa', *Kesmas: National Public Health Journal*, 9(1), p. 71. doi: 10.21109/kesmas.v9i1.459.
- Whiting, D. R. *et al.* (2011) 'IDF Diabetes Atlas: Global estimates of the prevalence of diabetes for 2011 and 2030', *Diabetes Research and Clinical Practice*, 94(3), pp. 311–321. doi: 10.1016/j.diabres.2011.10.029.
- WHO (2011). *Diabetes Mellitus*.
- WHO (2016). *Global Report on Diabetes*.
- Widya S, S., Budi P, L. and Purba, M. B. (2016) 'Konseling gizi mempengaruhi kualitas diet pasien diabetes mellitus tipe 2 di RSUP Dr. Sardjito Yogyakarta', *Jurnal Gizi dan Dietetik Indonesia (Indonesian Journal of Nutrition and Dietetics)*, 3(1), p. 31. doi: 10.21927/ijnd.2015.3(1).31-40.
- Widyastuti, D. A. (2013) 'Profil darah tikus putih wistar pada kondisi subkronis pemberian natrium nitrit', *Jurnal Sain Veteriner*, 31(201–215).
- Winarti, S. (2010). *Makanan Fungsional*, Kanisius. Yogyakarta: Graha Ilmu.
- Wu, Y. *et al.* (2014) 'Risk factors contributing to type 2 diabetes and recent advantages in the treatment and prevention', *International Journal of Medical Science*, 11(11), pp. 1185–1200.

- Xu, K. *et al.* (2011) ‘UCP2 -866G/A and Ala55Val, and UCP3 -55C/T polymorphisms in association with type 2 diabetes susceptibility: A meta-analysis study’, *Diabetologia*, 54(9), pp. 2315–2324. doi: 10.1007/s00125-011-2245-y.
- Yabe, D. *et al.* (2015) ‘ $\beta$  cell dysfunction versus insulin resistance in the pathogenesis of type 2 diabetes in East Asians’, *Current Diabetes Reports*, 15(6). doi: 10.1007/s11892-015-0602-9.
- Yusni, Y. and Meutia, F. (2020) ‘Action mechanism of Rosella (*Hibiscus sabdariffa* L.) used to treat metabolic syndrome in elderly women’, *Evidence-based Complementary and Alternative Medicine*, 2020. doi: 10.1155/2020/5351318.
- Zakaria, F. R., Prangdimurti, E. and Damanik, R. (2014) ‘The effect of Roselle extract (*Hibiscus sabdariffa* Linn.) on blood glucose level and total antioxidant level on diabetic rat induced by streptozotocin’, *IOSR Journal of Pharmacy (IOSRPHR)*, 4(10), pp. 08–16. doi: 10.9790/3013-0401008016.
- Zangeneh, F., Arora, P. S., Dyck, P. J., Bekris, L., Lernmark, A., Achenbach, S. J., Oberg, A. L., & Rizza, R. A. (2006). Effects of duration of type 2 diabetes mellitus on insulin secretion. *Endocrine practice: official journal of the American College of Endocrinology and the American Association of Clinical Endocrinologists*, 12(4), 388–393. <https://doi.org/10.4158/EP.12.4.388>
- Zhang, C. Y. *et al.* (2001) ‘Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity,  $\beta$  cell dysfunction, and type 2 diabetes’, *Cell*, 105(6), pp. 745–755. doi: 10.1016/S0092-8674(01)00378-6.