



References

- Alkaff TR. (2006). Hubungan Variasi Musim Dengan Kejadian Preeklamsia Di Rsup. Dr. Sardjito Yogyakarta. Fakultas Kedokteran Universitas Gadjah Mada Yogyakarta.
- Badan Perencanaan Pembangunan Nasional. (2013). Riset Kesehatan Dasar (RISKESDAS) 2013. Badan Perencanaan Pembangunan Nasional, Jakarta.
- Benirschke K, Kaufmann P. (1995). Pathology of the Human Placenta. 3rd ed. New York, NY: Springer-Verlag.
- Boghossian, NS, Yeung E., Mendola P., Hinkle SN, Laughon, SK, Zhang C., Albert PS. (2014). Risk factors differ between recurrent and incident preeclampsia: a hospital-based cohort study. Ann Epidemiol, Vol. 24, No. 12, pp. 871-877.
- Duley L. (2009). The global impact of pre-eclampsia and eclampsia. Semin Perinatol, 33:130–137.
- Brown MA, Mackenzie C, Dunsmuir W, Roberts L, Ikin K, Matthews J, Mangos G, Davis G. (2007). Can we predict recurrence of pre-eclampsia or gestational hypertension?. The British Journal of Obstetrics and Gynaecology, 114 (8): 984-993.
- Catov JM, Ness RB, Kip KE, Olse J. (2007). Risk of early or severe pre-eclampsia related to pre-existing conditions. Int J Epidemiol, 36 (2): 412-419.
- Cunningham FG, Leveno KJ, Bloom SL, Spong CY, Dashe JS, Hoffman B, Casey BM, Sheffield JS. (2014). Williams Obstetric, 24th ed. McGraw-Hill, United States of America.
- Fisher S, Roberts JM. 2014. The placenta in normal pregnancy and preeclampsia. Chesley's Hypertensive Disorders in Pregnancy, 4th ed. Amsterdam, Academic Press.
- Gupta T, Gupta N, Jain J, Gupta S, Bhatia P, Bagla J. (2013). Maternal and Perinatal Outcome in Patients with Severe Preeclampsia/Eclampsia with and without HELLP Syndrome. Delhi, India: Journal of Universal College of Medical Sciences, 1 (4): 7-12.
- Harahap, DH. (2013). Perbandingan Luaran Perinatal Antara Persalinan Vaginal dan Bedan Besar Pada Pasien Preeklamsia di RSUP Dr. Sardjito Tahun 2009-2011. Fakultas Kedokteran Universitas Gadjah Mada, Yogyakarta.



Haram K, Svendsen E, Abildgaard U. (2009). The HELLP syndrome : Clinical issues and management . A Review. BMC Pregnancy and Chidlbirth, 9 (8): 1-15.

Harris BM, Kuczkowski KM. (2005). Diagnostic dilemma: hepatic rupture due to HELLP syndrome vs. trauma. Arch Gynecol Obstet; 272: 176-178

Hawa, LR. (2009). Faktor Resiko Kematian Maternal Pada Preeklamsia Berat Dan Eklamsia Di Rumah Sakit Sardjito Tahun 2004-2007. Fakultas Kedokteran Universitas Gadjah Mada, Yogyakarta.

Hjartardottir S, Leifsson BG, Geirsson RT, Steinhorstdottir V. (2006). American Journal of Obstetrics and Gynecology, 194: 916-920.

Hogan MC, Foreman KJ, Naghavi M, Ahn SY, Wang M, Makela SM, Lopez AD, Lozano R, Murray CJ. (2010). Maternal mortality for 181 countries, 1980–2008: A systematic analysis of progress towards Millennium Development Goal 5. The Lancet, 375(9726), pp. 1609-1623.

Jabeen M, Gul F. (2004). Abruptio Placentae Risk facots and perinatal outcome. J Postgrade Med Inst, 18: 669-76.

Jeyabalan A. (2014). Epidemiology of preeclampsia: Impact of obesity. National Institutes of Health, 71: 1-4.

Kementrian Kesehatan Republik Indonesia. 2014. Situasi Kesehatan Ibu, Pusat Data dan Informasi Kementrian Kesehatan Republik Indonesia, Jakarta.

Kıñay T, Küçük C, Kayıkçıoğlu F, Karakaya J. (2015). Severe Preeclampsia versus HELLP Syndrome: Maternal and Perinatal Outcomes at <34 and \geq 34 Weeks' Gestation. Balkan Med J, 32 (4): 359-363.

Kurniawardhani DR. (2013). Incident Of Preeclampsia Effects On Maternal And Perinatal In Dr.sardjito Hospital 2009-2011. Fakultas Kedokteran Universitas Gadjah Mada Yogyakarta.

Liu CM, Chang SD, Cheng PJ, Chao AS. (2006). Comparisons of maternal and perinatal outcomes in Taiwanese women with complete and partial HELLP syndrome and women with severe preeclampsia without HELLP. J Obstet Gynaecol Res, 32 (6): 550–558.

Mallick IH, Syed SA, Kar AK. (2007). Liver rupture following delivery: HELLP needed. Emerg Med J, 24(5):372.

McDonald SD, Best C, Lam K. (2009). The recurrence risk of severe de novo preeclampsia in singleton pregnancies: a population-based cohort. The British Journal of Obstetrics and Gynaecology, 116 (12): 1578



Mihu D, Costin N, Mihu CM, Seicean A, Ciortea R. (2007). HELLP Syndrome- a Multisystemic Disorder. *J Gastrointestin Liver Dis*, 16 (4): 419-424.

NSW Ministry of Health. (2010). Maternity - Prevention, Early Recognition & Management of Postpartum Haemorrhage (PPH). Sydney.

O'Brien TE, Ray JG, Chan WS. (2003). Maternal Body Mass Index and the Risk of Preeclampsia: A Systematic Overview. *Epidemiology*, 14 (3): 368-374.

Polley LS. (2003). Anesthetic management of hypertension in pregnancy. *Clin Obstet Gynecol*; 46: 688-699.

Prawirohardjo S. (2014). Ilmu Kebidanan, edisi keempat. PT Bina Pustaka Sarwono Prawirohardjo, Jakarta.

Rahmayanti S. (2013). HELLP Syndrome and Partial HELLP Syndrome in Severe Preeklampsia: The Perinatal Outcomes. Fakultas Kedokteran Universitas Gadjah Mada Yogyakarta.

Rath W, Faridi A, Dudenhausen J W. (2000). HELLP syndrome. *J Perinat Med*, 28:249–260.

Redman, C. W., Bonnar, J., & Beilin, L. (1978). Early platelet consumption in pre-eclampsia. *British Medical Journal*, 1(6111), 467–469.

Redman CW, Sargent IL. 2005. Latest advances in understanding preeclampsia. *Science*, 308: 1592–1594.

Sibai BM. (2004). Diagnosis, Controversies, and Management of the Syndrome of Hemolysis, Elevated Liver Enzymes, and Low Platelet Count. American College of Obstetricians and Gynecologists, 103 (5): 981-990.

Sibai, BM. (1990). The HELLP syndrome (hemolysis, elevated liver enzyme, and low platelets): much ado about nothing?. *Am J Obstet Gynecol*, 162 (2): 311-316.

Sibai BM, Dekker G, Kupferminc M. (2005). Pre-Eclampsia. *The Lancet*, 365: 785-799.

Smith JR, Brennan BG. (2004). Postpartum Hemorrhage, <http://www.emedicine.com>.

Smulian J, Shen-Schwarz S, Scorza W, Kinzler W, Vintzileos A. (2004). A clinicohistopathologic comparison between HELLP syndrome and severe preeclampsia. *J Matern Fetal Neonatal Med*, 16:287–293.



Steegers EAP, Dadelszen PV, Duvekot JJ, Pijenborg R. (2010). Pre-eclampsia. *The Lancet*, 376: 631–644.

Suzuki T, Kaneda T. (2007). Anesthesia in three women with HELLP syndrome. *Masui*; 56: 838-841

The American College of Obstetricians and Gynecologists. (2013). Hypertension in Pregnancy, American College of Obstetricians and Gynecologists, Washington.

Thomas T, Jophy R, Mhaskar A, Misguith D. (2005). Are we increasing serious maternal morbidity by postponing termination of pregnancy in severe pre-eclampsia/eclampsia?. *J Obstet Gynaecol*, 25: 347-351.

Turgut A, Demirci O, Demirci E, Uludoğan M. (2010). Comparison of maternal and neonatal outcomes in women with HELLP syndrome and women with severe preeclampsia without HELLP syndrome. *J Prenat Med*, 4 (3): 51–58.

Vinnars MT, Wijnaendts LCD, Westgren M, Bolte A., Papadogiannakis N. (2008). Severe preeclampsia with and without HELLP differ with regard to placental pathology. *Hypertension*, 55: 1295–1299.

Wagner LK. (2004). Diagnosis and Management of Preeclampsia. *American Association of Family Physician*, 70 (12): 2317-2319.

World Health Organization. (2014). WHO Recommendations For Prevention and Treatment of Pre-eclampsia and Eclampsia, World Health Organization, Geneva.