

### Daftar Pustaka

- Balconi, E., Pennati, A., Crobu, D., Pandini, V., Cerutti, R., Zanetti, G., et al., 2009. The ferredoxin-NADP<sup>+</sup> reductase/ferredoxin electron transfer system of *Plasmodium falciparum*. *FEBS Journal*, vol.276, no.14, pp. 4249-4260.
- Basco, L.K., 2007. Field application of in vitro assays for the sensitivity of human malaria parasites to antimalarial drugs . *WHO Publications*
- Batista, R., de Jesus Silva Júnior, A. & de Oliveira, A.B., 2009. Plant-Derived Antimalarial Agents: New Leads and Efficient Phytomedicines. Part II. Non-Alkaloidal Natural Products. *Molecules*, vol. 14, no.8, pp.3037–3072
- Brunton, L. L., Chabner, B. A. & Knollmann, B. C., 2011. Goodman & Gilman's The Pharmacological Basis of Therapeutics. 12th ed. New York:McGraw-Hill.
- Bhattacharya, A., Mishra, L. C., Sharma, M., Awasthi, S. K., Bhasin, V. K., 2009. Antimalarial pharmacodynamics of chalcone derivatives in combination with artemisinin against *Plasmodium falciparum* in vitro. *European Journal of Medicinal Chemistry*, vol.44, no.9, pp.3388–3393.
- Chen, M., Christensen, S. B. & Zhai, L., 1997. The Novel Oxygenated Chalcone, The novel oxygenated chalcone, 2,4-dimethoxy-4'-butoxychalcone, exhibits potent activity against human malaria parasite *Plasmodium falciparum* in vitro and rodent parasites *Plasmodium berghei* and *Plasmodium yoelii* in vivo. *The Journal of Infectious Diseases*, Volume 176, pp. 1327-1333.

- Chen, M., Theander, T. G., Christensen, S. B., Hviid, L., Zhai, L., Kharazmi, A.,  
1994. Licochalcone A, a new antimalarial agent, inhibits in vitro growth of  
the human malaria parasite *Plasmodium falciparum* and protects mice from  
*P. yoelii* infection. *Antimicrobial Agents and Chemotherapy*, vol.38, no.7,  
pp. 1470-1475.
- Cruz, L. R., Spangenberg, T., Lacerda, M. V. G. & Wells, T. N. C., 2013. Malaria  
in South America: a drug discovery perspective.. *Malaria journal*, 12(1), p.  
168.
- Domínguez, J.N. et al., 2001. Synthesis of quinolinyl chalcones and evaluation of  
their antimalarial activity. *European Journal of Medicinal Chemistry*,  
vol.36, pp.555–560.
- Egan, T.J., 2008. Haemozoin formation. *Molecular and Biochemical  
Parasitology*, vol.157, no.2, pp.127–136.
- Farrar, J., Hotez, P. J., Junghanss, T. & Kang, G., Lallo, D., White, N.J., 2014.  
*Manson's Tropical Disease*. 23th ed. s.l.:Elsevier.
- Frölich, S., Schubert, C., Bienzle, U. & Jenett-Siems, K., 2005. In vitro  
antiplasmodial activity of prenylated chalcone derivatives of hops (*Humulus  
lupulus*) and their interaction with haemin. *Journal of Antimicrobial  
Chemotherapy*, vol.55, no.6, pp. 883-887.

- Humardewayanti, R. & Nugroho, A., 2012. Malaria. In: R. Humardewayanti & A. Nugroho, eds. *Buku Ajar Pendidikan Dokter Ilmu Penyakit Dalam*. Yogyakarta: Bagian Ilmu Penyakit Dalam FK UGM, pp. 332-367.
- Hyde, J.E., 2009. Exploring the folate pathway in *Plasmodium falciparum*. , vo.94, no.3, pp.191–206.
- Kalanon, M. & McFadden, G. I., 2010. Malaria, *Plasmodium falciparum* and its apicoplast.. *Biochemical Society transactions*, vol.38, no.3, pp. 775-782.
- Kimata-Ariga, Y. Saitoh, T., Ikegami, T., Horii, T., Hase, T., et al., 2007. Molecular interaction of ferredoxin and ferredoxin-NADP<sup>+</sup> reductase from human malaria parasite. *Journal of Biochemistry*, vol.142, no.6, pp. 715-720.
- Kumar, R., Mohanakrishnan, D., Sharma, A., Kaushik, N. K., Kalia, K., Sinha, A. K., et al., 2010. Reinvestigation of structure-activity relationship of methoxylated chalcones as antimalarials: Synthesis and evaluation of 2,4,5-trimethoxy substituted patterns as lead candidates derived from abundantly available natural  $\beta$ -asarone. *European Journal of Medicinal Chemistry*, vol. 45, no.11, pp.5292–5301.
- Matthews, H., Ali, M., Carter, V., Underhill, A., Hunt, J., Szor, H., et al., 2012. Variation in apoptosis mechanisms employed by malaria parasites: the roles of inducers, dose dependence and parasite stages. *Malaria Journal*, 11(1), p.297. Available at: Malaria Journal.

- Milani, M., Balconi, E., Aliverti, A. & Mastrangelo, E., 2007. Ferredoxin-NADP+ Reductase from *Plasmodium falciparum* Undergoes NADP+-dependent Dimerization and Inactivation: Functional and Crystallographic Analysis. *Journal of Molecular Biology*, vol.367, no.2, pp. 501-513.
- Moll, K., Ljungström, I., Perlmann, H., Scherf, A., Wahlgren, M., 2008. *Methods in Malaria Research*.5<sup>th</sup> ed. MR4/ATCC, Manassas, Virginia.
- Petersen, E., 1987. In vitro susceptibility of *Plasmodium falciparum* malaria to pyrimethamine, sulfadoxine, trimethoprim and sulfamethoxazole, singly and in combination. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, vol.81, pp.238–241.
- Pingaew, R., Saekee, A., Mandi, P. & Nantasenamat, C., 2014. Synthesis, biological evaluation and molecular docking of novel. *European Journal of Medicinal Chemistry*, vol.85, pp. 65-76.
- Ralph, S., van Dooren, G. G., Waller, R. F., Crawford, M. J., Fraunholz, M. J., Foth, B. J., *et al.*, 2004. Tropical infectious diseases: metabolic maps and functions of the *Plasmodium falciparum* apicoplast. *Nature reviews. Microbiology*, vol. 2, no. 3, pp.203–216.
- Sholikhah, E. N., 2010. Antiplasmodium Baru Turunan N-Alkil dan N-Benzil 1,10-Fenantrolin: Kajian Aktivitas In Vitro, Sitotoksitas, Sifat Fisika Kimia, Dan Profil Farmakokinetika, Yogyakarta: Fakultas Kedokteran Universitas Gadjah Mada.

- Sinha, S., Medhi, B. & Sehgal, R., 2013. Chalcones as an Emerging Lead Molecule for Antimalarial Therapy : A Review. *Journal of Modern Medicinal Chemistry*, vol.1, pp. 64-77.
- Smit, F. J. & N'Da, D. D., 2014. Synthesis, in vitro antimalarial activity and cytotoxicity of novel 4-aminoquinoliny-chalcone amides. *Bioorganic and Medicinal Chemistry*, vol.22, no.3, pp. 1128-1138.
- Smit, F. J., van Biljon, R. a., Birkholtz, L.-M. & N'Da, D. D., 2015. Synthesis and in vitro biological evaluation of dihydroartemisinin-chalcone esters. *European Journal of Medicinal Chemistry*, vol.90, pp. 33-44.
- Suwito, H., Kristanti, A. N., Nyoman, N., Puspaningsih, T., 2014<sup>a</sup>. Review Article Chalcones : Synthesis , structure diversity and pharmacological aspects. *Journal of Chemical and Pharmaceutical Research*, vol.6, no.5, pp.1076–1088.
- Suwito, H. Jumina, Mustofa, Pudjiastuti, P., Fanani, M., Kimata-Arigo, Y., et al., 2014<sup>b</sup>. Design and Synthesis of Chalcone Derivatives as Inhibitors of the Ferredoxin — Ferredoxin-NADP<sup>+</sup> Reductase Interaction of *Plasmodium falciparum*: Pursuing New Antimalarial Agents. *Molecules*, vol.19, no.12, pp. 21473-21488.
- Valla, A., Valla, B., Cartier, D., Le, R., Labia, R., Florent, L., Charneau, S., Schrevel, J., Potier, P., et al., 2006. New syntheses and potential

antimalarial activities of new “ retinoid-like chalcones .” , vol. 41, pp.142–146.

Wang, P., Sims, P.F. & Hyde, J.E., 1997<sup>a</sup>. A modified in vitro sulfadoxine susceptibility assay for Plasmodium falciparum suitable for investigating Fansidar resistance. *Parasitology*, vol. 115, pp.223–30

Wang, P., Read, M., Sims, P. F., Hyde, J. E., et al., 1997<sup>b</sup>. Sulfadoxine resistance in the human malaria parasite Plasmodium falciparum is determined by mutations in dihydropteroate synthetase and an additional factor associated with folate utilization. *Molecular microbiology*, vol.23, no.5, pp.979–986.

WHO, 2010. Guidelines for the treatment of malaria. 2nd ed. Geneva,Switzerland: World Health Organization.

WHO, 2015. Guidelines for the treatment of malaria. 3rd ed. Geneva,Switzerland: World Health Organization.

WHO, 2014. *World Malaria Report 2014*, Geneva,Switzerland: WHO.