

## REFERENCES

- Allen, C.M.C., 1984. Predicting the outcome of acute stroke : a prognostic score. *J Neurol Neurosurgery Psychiatry* ; 47 : 475-480
- Aminoff, M., Boller, F. and Swaab, D. (2008). *Handbook of Clinical Neurology : Neuropsychology and Behavioral neurology*. 3rd ed. Amsterdam: Elsevier.
- Beydoun, M., Beydoun, H. and Dore, G. (2015). Total serum cholesterol, atherogenic indices and their longitudinal association with depressive symptoms among US adults. *Translational Psychiatry*, 5(3).
- Carlsson, C., Nondahl, D. and Klein, B. (2009). Increased Atherogenic Lipoproteins Are Associated with Cognitive Impairment: Effects of Statins and Subclinical Atherosclerosis. *National Institutes of Health*, 23(1), pp.11-17.
- Dobiasova M, Frohlich J, Sedova M, Cheung MC, Brown BG. Cholesterol Esterification and Atherogenic Index of Plasma correlate with Lipoprotein size and

findings on Coronary Angiography. *J Lipid Res.*2011;52 (3) :566-71

Folstein, M.F., Folstein, S.E., McHugh, P.R. 1975. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician". *Journal of Psychiatric Research* 12 (3): 189-98

Gorelick PB, Wong KS, Bae HJ, Pandey DK. Large artery intracranial occlusive disease: a large worldwide burden but a relatively neglected frontier. *Stroke.*2008;39 (8) :2396-2399

Gutiérrez Pérez C, Savborg M, Pahlman U, et al. High frequency of cognitive dysfunction before stroke among older people. *Int J Geriatr Psychiatry* 2011;26:622-9

Holmstedt CA, Turan TN, Chimowitz MI. Atherosclerotic intracranial arterial stenosis: Risk factors, diagnosis and treatment. *Lancet Neurol.* 2013;12:1106-14

Ihle-Hansen H, Thommessen B, Wyller TB, et al. Incidence and subtypes of MCI and dementia 1 year after first-ever stroke in patients without

pre-existing cognitive impairment. *Dement Geriatr Cogn Disord* 2011;32:401-7

Jacquin A, Binquet C, Rouaud O, et al. Post-stroke cognitive impairment: high prevalence and determining factors in a cohort of mild stroke. *J Alzheimers Dis* 2014;40:1029-38.

Kumar, V. and Abbas, (2007). *Robins : Basic Pathology*. 8th ed. Chicago: Saunders

Kanthe, P., Bheemshetty, P. and Shrilaxmi, B. (2015). Atherogenic Index as a Predictor of Cardiovascular Risk among Women with Different Grades of Obesity. *International Journal of Collaborative Research on Internal Medicine & Public Health Vol.*, 4(10), pp.1768-1770.

Moussouttas M, Aguilar L, Fuentes K, et al. Cerebrovascular disease among patients from the Indian subcontinent. *Neurology*. 2006;67(5):894-896.

Onat, A. and Can, G. (2010). "Atherogenic index of plasma" (log<sub>10</sub> triglyceride/high-density lipoprotein-cholesterol) predicts high blood pressure, diabetes, and vascular event. *Journal Of Clinical Lipidology*, 4(2), pp.89-

- Ovbiagele B, Nguyen-Huynh MN. Stroke epidemiology: Advancing our understanding of disease mechanism and therapy. *Neurotherapeutics*.2011;8:319-29
- Rasquin SM, Verhey FR, van Oostenbrugge RJ, et al. Demographic and CT scan features related to cognitive impairment in the first year after stroke. *J Neurol Neurosurg Psychiatry* 2004;75:1562-7.
- Rist PM, Chalmers J, Arima H, et al. Baseline cognitive function, recurrent stroke, and risk of dementia in patients with stroke. *Stroke* 2013;44:1790-5.
- Ropper, A. and Samuels, M. (2009). *Adams and Victor's : Principle of Neurology*. 9th ed. United States: Mc Graw Hill.
- Schiffer, R., Rao, S. and Fogel, B. (2003). *Neuropsychiatry*. 2nd ed. Philadelphia: Lippincot Williams & WilSchiffer
- Setyopranoto, I. (2012). *Odem Otak : pada pasien stroke iskemik akut*. Yogyakarta: Badan Penerbit Fakultas Kedokteran UGM
- Russel, R. (1976). *Cerebral Arterial Disease*. New York: Churchill Livingstone.

Wang Y, Zhao X, Liu L, Soo YO, Pu Y, Pan Y, et al. ,  
CICAS Study Group. Prevalence and outcomes of  
symptomatic intracranial large artery stenoses  
and occlusions in China: The Chinese Intracranial  
Atherosclerosis (CICAS)  
Study. *Stroke*. 2014;45:663-9

Zambón, D., Quintana, M., Mata, P., Alonso, R.,  
Benavent, J., Cruz-Sánchez, F., Gich, J., Pocoví,  
M., Civeira, F., Capurro, S., Bachman, D.,  
Sambamurti, K., Nicholas, J. and Pappolla, M.  
(2010). Higher Incidence of Mild Cognitive  
Impairment in Familial Hypercholesterolemia. *The  
American Journal of Medicine*, 123(3), pp.267-274.