

## DAFTAR PUSTAKA

- Abed, A.A., Ayesh, B.M., Hamdona, O.M., 2013. Single nucleotide polymorphism E23K of KCNJ11 gene and other risk factors associated with type-2 diabetes mellitus in Gaza. *Int. J. Chem. Life Sci. 2*, 1146–1152.
- Alami, F.M., Ahmadi, M., Bazrafshan, H., Tabarraei, A., Khosravi, A., Tabatabaiefar, M.A., Samaei, N.M., 2012. Association of the TCF7L2 rs12255372 (G/T) variant with type 2 diabetes mellitus in an Iranian population. *Genet. Mol. Biol. 35*, 413–417. doi:10.1590/S1415-47572012005000029
- Alsmadi, O., Al-Rubean, K., Mohamed, G., Alkayal, F., Al Saud, H., Al Saud, N.A., AAl Daghri, N., Mohammad, S., Meyer, B. F. Weak or no association of TCF7L2 variants with Type 2 diabetes risk in an Arab population. *BMC Medical Genetics 2008,9:72* doi:10.1186/1471-2350-9-72
- Asayama, K., Dobashi, K., Hayashibe, H., Koderu, K., Uchida, N., Nakane, T., Araki, T., Nakazawa, S., 2002. Threshold values of visceral fat measures and their anthropometric alternatives for metabolic derangement in Japanese obese boys. *Publ. Online 30 January 2002* Doi101038/sjijo0801865 26. doi:10.1038/sj/ijo/0801865
- Ball GD, Huang TT, Gower BA, Cruz ML, Shaibi GO, Weigensberg MJ et al. (2006). Longitudinal changes in insulin sensitivity, insulin secretion, and beta-cell function during puberty. *J Pediatr; 148: 16–22*.
- Bano, G., 2013. Glucose homeostasis, obesity and diabetes. *Best Pract. Res. Clin. Obstet. Gynaecol. 27*, 715–726. doi:10.1016/j.bpobgyn.2013.02.007
- Barros, C.M.A.R., Araújo-Neto, A.P., Lopes, T.R., Barros, M.A.L., Motta, F.J.N., Canalle, R., Nunes, L.C.C., Rey, J.A., Burbano, R.R., Lima-Barros, M.A., Yoshioka, F.K.N., Pinto, G.R., 2014. Association of the rs7903146 and rs12255372 polymorphisms in the TCF7L2 gene with type 2 diabetes in a population from northeastern Brazil. *Genet. Mol. Res. 13*, 7889–7898. doi:10.4238/2014.September.29.1
- Bauman, AE. (2004). updating the evidence that physical activity is good for health: an epidemiological review 2000-2003. *Journal of Science and Medicine in Sport 7 (1): Supplement: 6-19*.
- Blair SN, Nichaman MZ. (2002). the public health problem of increasing prevalence rates of obesity and what should be done about it. *Mayo Clin Proc 77(2)*, 109-13.
- Boden G (1998). Free fatty acids (FFA), a link between obesity and insulin resistance. *Front Biosci 47:d169 – d175*

- Bollheimer LC, Skelly RH, Chester MW, McGarry JD, Rhodes CJ. (1998). Chronic exposure to free fatty acid reduces pancreatic  $\beta$  cell insulin content by increasing basal insulin secretion that is not compensated for by a corresponding increase in proinsulin biosynthesis translation. *J Clin Invest* 101:1094–1101
- Botto, Lorenzo D. and Khoury, Muin J. Commentary: Facing the Challenge of Gene-Environment Interaction: The Two-by-Four Table and Beyond. *American Journal of Epidemiology* Vol. 153 No.10 2001
- Bouchard C. (1991). Current understanding of Aetiology of Obesity Genetic and Non-Genetic Factors. *Am J Clin Nutr*; 53:1561-1565.
- Bournat, J.C., Brown, C.W., 2010. Mitochondrial dysfunction in obesity. *Curr. Opin. Endocrinol. Diabetes Obes.* 17, 446–452. doi:10.1097/MED.0b013e32833c3026
- Britz B, Siegfried W, Ziegler A, Lamertz C, Herpertz-Dahlmann BM, Remschmidt H et al. (2000). Rates of psychiatric disorders in a clinical study group of adolescents with extreme obesity and in obese adolescents ascertained via a population based study. *Int J Obes Relat Metab Disord*; 24: 1707–1714.
- Brufani, C., Ciampalini, P., Grossi, A., Fiori, R., Fintini, D., Tozzi, A., Cappa, M., Barbetti, F., 2010. Glucose tolerance status in 510 children and adolescents attending an obesity clinic in Central Italy. *Pediatr. Diabetes* 11, 47–54. doi:10.1111/j.1399-5448.2009.00527.x
- Butler AE, Janson J, Bonner-Weirs S, Ritzel R, Rizza RA, Butler PC. (2003).  $\beta$ -Cell Deficit And Increased  $\beta$ -Cell Apoptosis In Humans With Type 2 Diabetes. *Diabetes* vol 52, 102-110
- Chandak, G.R., Janipalli, C.S., Bhaskar, S., Kulkarni, S.R., Mohankrishna, P., Hattersley, A.T., Frayling, T.M., Yajnik, C.S., 2006. Common variants in the TCF7L2 gene are strongly associated with type 2 diabetes mellitus in the Indian population. *Diabetologia* 50, 63–67. doi:10.1007/s00125-006-0502-2
- Chen, C.-M., Lou, M.-F., Gau, B.-S., 2014. Prevalence of impaired fasting glucose and analysis of related factors in Taiwanese adolescents. *Pediatr. Diabetes* 15, 220–228. doi:10.1111/pedi.12081
- Clement K, Garner C, Harger J, Philippi A, LeDuc C, Carcy A, Harris TJ, Jury C, Cardon LR, Basdevant A, Demenais F, Guy-Grand B, North M, Froguel P. (1996). Indication for Linkage of The Human Ob Gene Region With Extreme Obesity. *Diabetes* 45:687-690.
- Coll AP, Farooqi IS & O’Rahilly S. (2007). The hormonal control of food intake. *Cell* 129, 251-262

- Comuzzie, Anthony G. (2003). The Genetics of Obesity in Mexican Americans: The Evidence from Genome Scanning Efforts in the San Antonio Family Heart Study. *Human Biology* 75(5): 635-646.
- Corbala n MS, Marti A, Forga L, Mart´inez-Gonza´lez MA & Mart´inez JA. (2002) Beta2-adrenoceptor polymorphism and abdominal obesity risk: effect modification by gender and HDL-cholesterol levels. *Eur J Nutr* 41, 114–118.
- Corbala n MS, MargaretoJ, Martinez JA & Marti A. (1999). High fat feeding reduced muscle uncoupling protein 3 expression in rats. *J Physiol Biochem* 55, 67–73
- Cornelis, M.C., Qi, L., Kraft, P., Hu, F.B., 2009. TCF7L2, dietary carbohydrate, and risk of type 2 diabetes in US women. *Am. J. Clin. Nutr.* 89, 1256–1262. doi:10.3945/ajcn.2008.27058
- Daalgaard, L., & Pedersen, O. (2001). Uncoupling proteins: functional characteristics and role in the pathogenesis of obesity and type II diabetes. *Diabetologia*, 44(8), 946–65.
- Daniels SR, Loggie JM, Khoury P, Kimball TR. (1998). Left ventricular geometry and severe left ventricular hypertrophy in children and adolescents with essential hypertension. *Circulation*; 97: 1907–1911.
- Daniels, S. R. (2009). Complications of obesity in children and adolescents, *International Journal of Obesity* (2005), 33 Suppl 1, S60–S65. doi:10.1038/ijo.2009.20
- de Onis, M. dan Blossner, M. (2000). Prevalence and trends of overweight among preschool children in developing countries. *Am J Clin Nutr*, 72, 1032-1039.
- de Pádua Cintra, I., Passos, M.A.Z., Santos, L.C. dos, da Costa Machado, H., Fisberg, M., 2014. Waist-to-Height Ratio Percentiles and Cutoffs for Obesity: A Cross-sectional Study in Brazilian Adolescents. *J. Health Popul. Nutr.* 32, 411.
- de Simone G, Devereur RB, Daniels SR, Koren MJ, Meyer RA, Laragh JH. (1995) Effect of growth on variability of left ventricular mass: assessment of allometric signals in adults and children and their capacity to predict cardiovascular risk. *J Am Coll Cardiol* 25 1056–1062.
- Deng, Y., Scherer, P.E., 2010. Adipokines as novel biomarkers and regulators of the metabolic syndrome. *Ann. N. Y. Acad. Sci.* 1212, E1–E19. doi:10.1111/j.1749-6632.2010.05875.x
- Denton, JS & Jacobson, DA. (2012). Channeling dysglycemia: ion channel variations perturbing glucose homeostasis. *Trends Endocrinol Metab*; 23(1): 41-48.

- Derosa, G., Ragonesi, P. D., Carbone, A., Fogari, E., Bianchi, L., Bonaventura, A., Maffioli, P. (2012). Vildagliptin added to metformin on  $\beta$ -cell function after a euglycemic hyperinsulinemic and hyperglycemic clamp in type 2 diabetes patients. *Diabetes Technology & Therapeutics*, 14(6), 475–84. doi:10.1089/dia.2011.0278
- Deurenberg-Yap M, Chew SK, Lin VF, Tan BY, van Staveren W A, Deurenberg P.(1999). Relationships between indices of obesity and its comorbidities in multi-ethnic Singapore. *Int J Obes Relat Metab Disord*; 23:1136-42.
- Donath MY, Gross DJ, Cerasi E, Kaiser N. (1999). Hyperglycemia- induced  $\beta$ -cell apoptosis in pancreatic islets of *Psammomys obesus* during development of diabetes. *Diabetes* 48:738 –744
- Donath MY, Halban PA. (2004). Decreased  $\beta$ -cell mass in diabetes: significance, mechanism and therapeutic implication. *Diabetologia* 47:581-589
- Donohoue, P.A. (2004). *Obesity dalam Nelson Textbook of Pediatric 17th Edition. Philadelphia: Saunders.*
- Drewnowski A, Popkin BM. (1997). The nutrition transition: new trends in the global diet. *Nutr Rev* 55, 31–43.
- Duan, Q.L., Dubé, M.-P., Frasure-Smith, N., Barhdadi, A., Lesperance, F., Thérout, P., St-Onge, J., Rouleau, G.A., McCaffery, J.M., 2007. Additive Effects of Obesity and TCF7L2 Variants on Risk for Type 2 Diabetes Among Cardiac Patients. *Diabetes Care* 30, 1621–1623. doi:10.2337/dc06-2421
- Dulloo AG, Seydoux J & Jacquet J (2004) Adaptive thermogenesis and uncoupling proteins: a reappraisal of their roles in fat metabolism and energy balance. *Physiol Behav* 83, 587–602.
- Dulskiene, V., Kuciene, R., Medzioniene, J., Benetis, R., 2014. Association between obesity and high blood pressure among Lithuanian adolescents: a cross-sectional study. *Ital. J. Pediatr.* 40, 60–77. doi:10.1186/s13052-014-0102-6
- Ello-Martin JA, Ledikwe JH, Rolls BJ. (2005). the influence of food portion size and energy density on energy intake: implications for weight management. *Am J Clin Nutr*; 82: 236S–241S.
- Erlichman J, Kerbey AL, James WPT. (2002). Physical activity and its impact on health outcomes. Paper 2: prevention of unhealthy weight gain and obesity by physical activity: an analysis of the evidence. *Obes Rev* 3, 273-87.
- Evans J.L., Ira D Goldfine, Betty A Maddux, Gerold M Grodsky. (2002). *Endocr. Rev.* 23: 599-622

- Farmer, S.R. (2006). 'Transcriptional Control of Adipocyte Formation'. *Cell Metab.* 4: 263–273.
- Ferranti S., Dariush Mozaffarian. (2008). The Perfect storm: Obesity, Adipocyte Dysfunction, and Metabolic Consequences. *Clinical Chemistry* 54:6 945-955
- Ferretti, R.L., Cintra, I.P., Passos, M.A., Ferrari, G.L., Fisberg, M., 2015. Elevated neck circumference and associated factors in adolescents. *BMC Public Health* 15, 208. doi:10.1186/s12889-015-1517-8
- Flynn JT, Alderman MH. (2005). Characteristics of children with primary hypertension seen at a referral center. *Pediatr Nephrol*; 20: 961–966.
- Food and Agriculture Organization. (1998). *Obesity in developing countries: causes and implications*. Rome, Italy: Food and Agriculture Organization.
- Froguel P and Boutin P. (2001). Genetic of Pathways Regulating Body Weight in The Development of Obesity in Humans. *Exp Biol Med* 226(11):991-996.
- Fujisawa T, Ikegami H, Kawaguchi Y & Ogihara T (1998) Meta-analysis of the association of Trp64Arg polymorphism of beta 3-adrenergic receptor gene with body mass index. *J Clin Endocrinol Metab* 83, 2441–2444.
- Fur, S.L., Fradin, D., Boileau, P., Bougnères, P., 2005. Association of Kir6.2 and INS VNTR variants with glucose homeostasis in young obese. *Physiol. Genomics* 22, 398–401. doi:10.1152/physiolgenomics.00090.2005
- Fur, S.L., Stunff, C.L., Santos, C.D., Bougnères, P., 2004. The Common –866 G/A Polymorphism in the Promoter of Uncoupling Protein 2 Is Associated With Increased Carbohydrate and Decreased Lipid Oxidation in Juvenile Obesity. *Diabetes* 53, 235–239. doi:10.2337/diabetes.53.1.235
- Garg MA, Dutta MK, Namita Mahalle. (2011). Study of beta cell function (by HOMA model) in Metabolic Syndrome. *Indian J. Endocrinol metab.* July 2011; 15 (Suppl1): S44-S49
- Gloyn, A. L., Diatloff-Zito, C., Edghill, E. L., Bellanné-Chantelot, C., Nivot, S., Coutant, R. Robert, J. J. (2006). KCNJ11 activating mutations are associated with developmental delay, epilepsy and neonatal diabetes syndrome and other neurological features. *European Journal of Human Genetics : EJHG*, 14(7), 824–30. doi:10.1038/sj.ejhg.5201629
- Gloyn, A.L., Siddiqui, A.J. & Ellard, S., (2006). Mutations in the Genes Encoding the Pancreatic Beta-Cell K ATP Channel Subunits Kir6.2 (KCNJ11) and SUR1 (ABCC) in Diabetes Mellitus and Hyperinsulinism. , 27(January), pp.220–231.

- Gong, B., Yu, J., Li, H., Li, W., Tong, X., 2011. The Effect of KCNJ11 Polymorphism on the Risk of Type 2 Diabetes: A Global Meta-Analysis Based on 49 Case-Control Studies. *DNA Cell Biol.* 31, 801–810. doi:10.1089/dna.2011.1445
- Govaerts C, Srinivasan S, Shapiro A, Zhang S, Picard F, Clement K, Lubrano-Berthelie C & Vaisse C. (2005). Obesity-associated mutation in the melanocortin 4 receptor provide novel insight into its function. *Peptides* 26, 1909-1919
- Grant SF, et al. (2006). Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat Genet*; 38(3): 320-3.
- Grant, S.F.A., Hakonarson, H. & Schwartz, S., (2010). Can the genetics of type 1 and type 2 diabetes shed light on the genetics of latent autoimmune diabetes in adults? *Endocrine reviews*, 31, pp.183–193.
- Gregoire, F.M. (2001). 'Adipocyte Differentiation: From Fibroblast to Endocrine Cell'. *Exp. Biol. Med.*, 226: 997–1002.
- Guo, X., Zheng, L., Li, Y., Yu, S., Zhou, X., Wang, R., Zhang, X., Sun, Z., Sun, Y., 2012. Gender-specific prevalence and associated risk factors of prehypertension among rural children and adolescents in Northeast China: a cross-sectional study. *Eur. J. Pediatr.* 172, 223–230. doi:10.1007/s00431-012-1873-7
- Hamet, P. and Tremblay, J. (2005) Genetics and genomics of depression. *Metabolism*, 54, 10–15.
- Hapsari, P.O. (2008). *Hubungan antara pola asupan energi dengan resistensi insulin pada remaja putri obes di Kota Yogyakarta*. Skripsi: Universitas Gadjah Mada.
- Harding HP, Ron D. (2002). Endoplasmic reticulum stress and the development of diabetes. A review. *Diabetes* 51(Suppl 3):S455-461
- Hartanti M, (2006). *Persepsi orang tua terhadap postur tubuh dan porsi makan siswa sekolah dasar yang obes dan tidak obes di Kota Yogyakarta*, Skripsi, Fakultas Kedokteran UGM, Yogyakarta
- Hatipoglu, N., Mazicioglu, M.M., Kurtoglu, S., Kendirci, M., 2009. Neck circumference: an additional tool of screening overweight and obesity in childhood. *Eur. J. Pediatr.* 169, 733–739. doi:10.1007/s00431-009-1104-z
- Hernandez-Escalante, V.M., Nava-Gonzalez, E.J., Voruganti, V.S., Kent, J.W., Haack, K., Laviada-Molina, H.A., Molina-Segui, F., Gallegos-Cabriales, E.C., Lopez-Alvarenga, J.C., Cole, S.A., Mezzles, M.J., Comuzzie, A.G., Bastarrachea, R.A., 2014. Replication of obesity and diabetes-related SNP associations in individuals from Yucatán, México. *Front. Genet.* 5. doi:10.3389/fgene.2014.00380

- Hession M, et al. (2008). Systematic review of randomized controlled trials of low-carbohydrate vs. low-fat/low-calorie diets in the management of obesity and its comorbidities. *Obes Rev*; 10(1): 36-50.
- Hidayati, Siti Nurul, et al (2006) Hubungan Asupan Zat Gizi dengan Indeks Massa Tubuh dengan Hiperlipidemia pada Murid SLTP yang obesitas di Yogyakarta, *Jurnal Sari Pediatri vol. 8 no. 1 pp 25-31*
- Hill, J.O., Kriketos, A.D., dan Peters, J.C. (2000). *Disturbances of Energy Balance dalam: Biochemical and Physiological Aspect of Human Nutrition*. Philadelphia: Saunders.
- Hingorjo, M.R., Qureshi, M.A., Mehdi, A., 2012. Neck circumference as a useful marker of obesity: a comparison with body mass index and waist circumference. *JPMA-J. Pak. Med. Assoc.* 62, 36.
- Hirsch, J., Salans, L.B., dan Aronne, L.J. (1995). *Obesity dalam Principles and Practice of Endocrinology and Metabolism Second Edition*. J.B. Philadelphia Lippincott Company
- Huertas-Vazquez, A., Plaisier, C., Weissglas-Volkov, D., Sinsheimer, J., Canizales-Quinteros, S., Cruz-Bautista, I., Nikkola, E., Herrera-Hernandez, M., Davila-Cervantes, A., Tusie-Luna, T., Taskinen, M.-R., Aguilar-Salinas, C., Pajukanta, P., 2007. TCF7L2 is associated with high serum triacylglycerol and differentially expressed in adipose tissue in families with familial combined hyperlipidaemia. *Diabetologia* 51, 62–69. doi:10.1007/s00125-007-0850-6
- Hunt JN, Stubbs DF. (1975). the volume and energy content of meals as determinants of gastric emptying. *J Physiol*; 245: 209– 225.
- Hunt JN. (1980). A possible relation between the regulation of gastric emptying and food intake. *Am J Physiol*; 239: G1–G4.
- Huriyati, E. (2006). Studi kohort prevalensi obesitas siswa-siswi sekolah lanjutan tingkat pertama Kota Yogyakarta. *Jurnal Gizi Klinik Indonesia, vol. 3, 1-5*.
- Huriyati, E., dan Julia, M. (2009). Tekanan darah siswa sekolah dasar obes dan tidak obes di Kota Yogyakarta. *Jurnal Gizi Klinik Indonesia, vol. 6, 60-63*.
- Huriyati, E., Harry Freitag Luglio, Prima Dhewi, A Fahmi Arif Tsani, Ahmad Hamim Sadewa, Mohammad Juffrie. Dyslipidemia, Insulin Resistance and Dietary Fat Intake in Obese and Normal Weight Adolescents: The Role of Uncoupling Protein 2 -866G/A Gene Polymorphism. Underreview of International Journal of Molecular and Epidemiology Genetic.
- Indriasari, D.(2008). *Hubungan antara resistensi insulin dengan tekanan darah pada remaja putri obes*. Skripsi: Universitas Gadjah Mada.

- Jensen, M. D. (2008). Role of body fat distribution and the metabolic complications of obesity. *The Journal of Clinical Endocrinology and Metabolism*, 93, S57–S63. doi:10.1210/jc.2008-1585
- Juhan-Vague, I., Alessi, M.C., Morange, P.E., 2000. Hypofibrinolysis and increased PAI-1 are linked to atherothrombosis via insulin resistance and obesity. *Ann. Med.* 32 Suppl 1, 78–84.
- Jun, H.S., Kim, I.K., Lee, H.J., Lee, H.J., Kang, J.H., Kim, J.R., Shin, H.D., Song, J., 2009. Effects of UCP2 and UCP3 Variants on the Manifestation of Overweight in Korean Children. *Obesity* 17, 355–362. doi:10.1038/oby.2008.531
- Kahn S.E., Hull R.L., Utzschneider K.M. (2006). Mechanisms linking obesity to insulin resistance and type 2 diabetes. *Nature*, vol 444. Doi:10.1038/nature05482.
- Kashyap S, Belfort R, Castaldelli A, Pratipanawatr T, Berria R, Pratipanawatr W, Bajaj M, Mandarino L, DeFronzo R, Cusi K. (2003). A sustained increase in plasma free fatty acids impairs insulin secretion in nondiabetic subjects genetically predisposed to develop type 2 diabetes. *Diabetes* 52:2461-2474
- Katz, S.L., Vaccani, J.-P., Clarke, J., Hoey, L., Colley, R.C., Barrowman, N.J., 2014. Creation of a reference dataset of neck sizes in children: standardizing a potential new tool for prediction of obesity-associated diseases? *BMC Pediatr.* 14, 159. doi:10.1186/1471-2431-14-159
- Katzmarzyk PT, et al. (2000). Familial resemblance for coronary heart disease risk: the HERITAGE Family Study. *Ethn Dis*; 10(2): 138-47.
- Kelishadi R, Hashemipour M, Sarraf-Zadegan N et al. (2001). Trend of atherosclerosis risk factors in children of Isfahan. *Asian Cardiovasc Thorac Ann*; 9:36-40.
- Keskin, M., Kurtoglu, S., Kendirci, M., Atabek, M. E., & Yazici, C. (2005). Homeostasis model assessment is more reliable than the fasting glucose/insulin ratio and quantitative insulin sensitivity check index for assessing insulin resistance among obese children and adolescents. *Pediatrics*, 115, e500–e503. doi:10.1542/peds.2004-1921
- Khoury, M., Manlihot, C., Dobbin, S., Gibson, D., Chahal, N., Wong, H., Davies, J., Stearne, K., Makerewich, O., Fisher, A., others, 2011. 22-The association of cardiovascular risk and high body mass index in adolescents is influenced by waist circumference. *Can. J. Diabetes* 35, 145.
- Klünder-Klünder, M., Mejía-Benitez, M.A., Flores-Huerta, S., Burguete-García, A.I., García-Mena, J., Cruz, M., 2011. rs12255372 Variant of TCF7L2 Gene Is Protective for Obesity in Mexican Children. *Arch. Med. Res.* 42, 495–501. doi:10.1016/j.arcmed.2011.05.006

- Kong, A.P., Choi, K.-C., Ko, G.T., Wong, G.W., Ozaki, R., So, W.-Y., Tong, P.C., Chan, J.C., 2008. Associations of overweight with insulin resistance,  $\beta$ -cell function and inflammatory markers in Chinese adolescents. *Pediatr. Diabetes* 9, 488–495. doi:10.1111/j.1399-5448.2008.00410.x
- Laakso M, Edelman SV, Brechtel G, Baron AD. (1990). Decreased effect of insulin to stimulate skeletal muscle blood flow in obese man. *J Clin Invest* 85:1844–1852.
- Landi Masquio, D.C., de Piano Ganen, A., da Silveira Campos, R.M., de Lima Sanches, P., Campos Corgosinho, F., Caranti, D., Tock, L., Túlio de Mello, M., Tufik, S., Dâmaso, A.R., 2015. Cut-off values of waist circumference to predict metabolic syndrome in obese adolescents. LOS VALORES CORTE CIRCUNFERENCIA CINTURA PARA PREDECIR EL Síndr. METABÓLICO EN Adolesc. *OBESOS* 31, 1540–1550. doi:10.3305/nh.2015.31.4.8442
- Laquatra, A. (2000). *Nutrition for Weight Management dalam Krause's Food, Nutrition, & Diet Therapy*. Philadelphia: Saunders.
- Large V, Hellström L, Reynisdóttir S, Lönnqvist F, Eriksson P, Lannfelt L & Arner P (1997). Human beta-2 adrenoceptor gene polymorphisms are highly frequent in obesity and associate with altered adipocyte beta-2 adrenoceptor function. *Clin Invest* 100, 3005–3013.
- Larsen, L.H., Ångquist, L., Vimalaswaran, K.S., Hager, J., Viguier, N., Loos, R.J., Handjieva-Darlenska, T., Jebb, S.A., Kunešova, M., Larsen, T.M., Martinez, J.A., Papadaki, A., Pfeiffer, A.F., Baak, M.A. van, Sørensen, T.I., Holst, C., Langin, D., Astrup, A., Saris, W.H., 2012. Analyses of single nucleotide polymorphisms in selected nutrient-sensitive genes in weight-regain prevention: the DIOGENES study. *Am. J. Clin. Nutr.* 95, 1254–1260. doi:10.3945/ajcn.111.016543
- Larson-Meyer, D.E., Newcomer, B.R., Ravussin, E., Volaufova, J., Bennett, B., Chalew, S., Cefalu, W.T., Sothorn, M., 2011. Intrahepatic and intramyocellular lipids are determinants of insulin resistance in prepubertal children. *Diabetologia* 54, 869–875. doi:10.1007/s00125-010-2022-3
- Laukkanen O, et al. (2004). Polymorphisms of the SUR1 (ABCC8) and Kir6.2 (KCNJ11) genes predict the conversion from impaired glucose tolerance to type 2 diabetes. *J Clin Endocrinol Metab*; 89(12): 6286-90.
- Laukkanen, O., Pihlajamäki, J., Lindström, J., Eriksson, J., Valle, T.T., Hämäläinen, H., Ilanne-Parikka, P., Keinänen-Kiukaanniemi, S., Tuomilehto, J., Uusitupa, M., Laakso, M., 2004. Polymorphisms of the SUR1 (ABCC8) and Kir6.2 (KCNJ11) Genes Predict the Conversion from Impaired Glucose Tolerance to Type 2 Diabetes. The Finnish Diabetes Prevention Study. *J. Clin. Endocrinol. Metab.* 89, 6286–6290. doi:10.1210/jc.2004-1204

- Lean ME, Han TS, Morrison CE. (1995). Waist circumference as a measure for indicating need for weight management. *BMJ.* ; 311:158-161
- Lebovitz HE. (2001). No Title. Insulin Resistance : Definition and Consequence, *Exp Clin E (109 Suppl)*, S135–48.
- Li, L., Shi, Y., Wang, X., Shi, W., Jiang, C., 2005. Single Nucleotide Polymorphisms in KATP Channels Muscular Impact on Type 2. *Diabetes* 54, 1592–1597. doi:10.2337/diabetes.54.5.1592
- Lin, W.-T., Huang, H.-L., Huang, M.-C., Chan, T.-F., Ciou, S.-Y., Lee, C.-Y., Chiu, Y.-W., Duh, T.-H., Lin, P.-L., Wang, T.-N., Liu, T.-Y., Lee, C.-H., 2013. Effects on uric acid, body mass index and blood pressure in adolescents of consuming beverages sweetened with high-fructose corn syrup. *Int. J. Obes.* 37, 532–539. doi:10.1038/ijo.2012.121
- Liu L, et al. (2012). An association between -866G/A polymorphism in the promoter of UCP2 and obesity: A meta-analysis. *Gene*; 514(1): 41-7.
- Lo´pez I, Marti A, Milagro F, Zulet MA, Moreno-Aliaga MJ & Mart´inez JA (2003) DNA microarray analysis of gene differentially expressed in diet-induced obese rats. *Obes Res* 11, 188–194.
- Loos RJ, et al. (2007). TCF7L2 polymorphisms modulate proinsulin levels and beta-cell function in a British Europid population. *Diabetes*; 56(7): 1943-7.
- Love-Osborne, K., Butler, N., Gao, D., Zeitler, P., 2006. Elevated fasting triglycerides predict impaired glucose tolerance in adolescents at risk for type 2 diabetes. *Pediatr. Diabetes* 7, 205–210. doi:10.1111/j.1399-5448.2006.00179.x
- Low S, et al. (2009). Review on epidemic of obesity. *Ann Acad Med Singapore*; 38(1): 57-9.
- Lyssenko, V., Lupi, R., Marchetti, P., Del Guerra, S., Orho-Melander, M., Almgren, P, Groop, L. (2005). Mechanisms by which common variants in the TCF7L2 gene increase risk of type 2 diabetes. *The Journal of clinical investigation (Vol. 117, pp. 2155–2163)*.doi: 10.1172/JCI30706
- Macho-Azcárate T, Marti A, Mart´inez JA & Ibañez J (2002) The Gln27Glu polymorphism outcome in obese women submitted to a maximal effort trial. *Int J Obes Relat Metab Disord* 26, 1434–1441.
- Macho-Azcárate T, et al. (2002). A maximal effort trial in obese women carrying the beta2-adrenoceptor Gln27Glu polymorphism. *J Physiol Biochem*; 58(2): 103-8.
- Macho-Azcárate T, et al. (2003). Basal fat oxidation and after a peak oxygen consumption test in obese women with a beta2 adrenoceptor gene polymorphism. *J Nutr Biochem*; 14(5): 275-9.

- Maedler K, Spinass GA, Lehmann R, Sergeev P, Weber M, Fontana A, Kaiser N, Donath MY (2001) Glucose induces  $\beta$ -cell apoptosis via upregulations of the Fas-receptor in human islets. *Diabetes* 50: 1683–1690
- Margareto J, Marti A, & Martínez JA (2002) Changes in UCP mRNA expression levels in brown adipose tissue and skeletal muscle after feeding a high-energy diet and relationships with leptin, glucose and PPARG2. *J Nutr Biochem* 12, 130– 137.
- Marti A, & Martínez JA (2006) Genetics of obesity: gene • diet interaction. *Int J Vitam Nutr Res* 76, 1–10.
- Marti A, Corbalan MC, Forga L, Martinez JA, Hinney A & Hebebrand J. (2003). A novel nonsense mutation in the melanocortin-4 receptor associated with obesity in a Spanish population. *Int J Obes Relat Metab Disord* 27, 385-388
- Marti A, Larrarte E, Novo FJ, García M, Warden CH & Martínez JA (2001) UCP2 muscle gene transfer modifies mitochondrial membrane potential. *Int J Obes Relat Metab Disord* 25, 68–75.
- Marti, A., Martínez-González, M. A., & Martínez, J. A. (2008). Interaction between genes and lifestyle factors on obesity. *The Proceedings of the Nutrition Society*.doi: 10.1017/S002966510800596X
- Marzi, C., Huth, C., Kolz, M., Grallert, H., Meisinger, C., Wichmann, H.-E., Rathmann, W., Herder, C., Illig, T., 2007. Variants of the Transcription Factor 7-Like 2 Gene ( *TCF7L2* ) are Strongly Associated with Type 2 Diabetes but not with the Metabolic Syndrome in the MONICA/KORA Surveys. *Horm. Metab. Res.* 39, 46–52. doi:10.1055/s-2007-957345
- Mason TM, Goh T, Tchipashvili V, Sandhu H, Gupta N, Lewis GF, Giacca A (1999) Prolonged elevation of plasma free fatty acids desensitizes the insulin secretory response to glucose in vivo in rats. *Diabetes* 48:524 – 530
- Matsuda J, Hosoda K, Itoha H, Sona C, Doi K, Tanaka T, Fukunaga Y, Inoue G' Nishimura H, Yoshimasa Y, Yamori Y, Nakaoa K. (1997). Cloning of rat uncoupling protein-3 and uncoupling protein-2 cDNAs: their gene expression in rats fed high-fat diet. *FEBS Letters* 418 200-204.
- Mattei, J., Qi, Q., Hu, F.B., Sacks, F.M., Qi, L., 2012. TCF7L2 genetic variants modulate the effect of dietary fat intake on changes in body composition during a weight-loss intervention. *Am. J. Clin. Nutr.* 96, 1129–1136. doi:10.3945/ajcn.112.038125
- Mazicioglu, M.M., Kurtoglu, S., Ozturk, A., Hatipoglu, N., Cicek, B., Ustunbas, H.B., 2010. Percentiles and mean values for neck circumference in Turkish children aged 6-18 years Mazicioglu et al. Neck circumference. *Acta Paediatr.* 99, 1847–1853. doi:10.1111/j.1651-2227.2010.01949.x

- Medawati, A, Hadi, H, Pramantara I D P, (2005). Hubungan antara asupan Energi, asupan lemak, dan obesitas pada remaja siswa, siswi SLTP di Kota Yogyakarta dan Kabupaten Bantul, *Jurnal Gizi Klinik Indonesia*, Vol. 1, no 3, 119-129.
- Miura, N., Ikezaki, A., Iwama, S., Matsuoka, H., Ito, K., Sugihara, S., 2006. Genetic factors and clinical significance of acanthosis nigricans in obese Japanese children and adolescents. *Acta Paediatr.* 95, 170–175. doi:10.1080/08035250500294015
- Moll, P. P., Burns, T. L., & Lauer, R. M. (1991). The genetic and environmental sources of body mass index variability: the Muscatine Ponderosity Family Study. *American Journal of Human Genetics*, 49, 1243–1255.
- Mo-suwan L, Junjana C, Puetpaiboon A. (1993). Increasing obesity in school children in a transitional society and the effect of the weight control program. *Southeast Asian J Trop Med Public Health*; 24:590-4
- Mozo J, Emre Y, Bouillaud F, Ricquier D & Criscuolo F (2005) Thermoregulation: what role for UCPs in mammals and birds? *Biosci Rep* 25, 227–249.
- Muhammad, H.F.L., Huriyati, E., Susilowati, R., dan Julia, M. 2009. Magnesium intake and insulin resistance in obese adolescent girls. *Paediatrica Indonesiana*, 49, 200-203.
- Munoz, J., Lok, K.H., Gower, B.A., Fernandez, J.R., Hunter, G.R., Lara-Castro, C., Luca, M.D., Garvey, W.T., 2006. Polymorphism in the Transcription Factor 7-Like 2 (TCF7L2) Gene Is Associated With Reduced Insulin Secretion in Nondiabetic Women. *Diabetes* 55, 3630–3634. doi:10.2337/db06-0574
- Nambiar, S., Truby, H., Davies, P.S., Baxter, K., 2013. Use of the waist-height ratio to predict metabolic syndrome in obese children and adolescents. *J. Paediatr. Child Health* 49, E281–E287. doi:10.1111/jpc.12147
- Nanfa, D., Sobngwi, E., Atogho-Tiedeu, B., Noubiap, J.J., Donfack, O.S., Mofo, E.P., Guewo-Fokeng, M., Metsadjio, A.N., Ngwa, E.N., Fosso, P.P., Djahmeni, E., Djokam-Dadjeu, R., Evehe, M.-S., Aminkeng, F., Mbacham, W.F., Mbanya, J.C., 2015. Association between the TCF7L2 rs12255372 (G/T) gene polymorphism and type 2 diabetes mellitus in a Cameroonian population: a pilot study. *Clin. Transl. Med.* 4, 17. doi:10.1186/s40169-015-0058-1
- Nielsen, E.-M. D., Hansen, L., Carstensen, B., Echwald, S. M., Drivsholm, T., Glümer, C., Pedersen, O. (2009). The E23K variant of Kir6.2 associates with impaired post-OGTT serum insulin response and increased risk of type 2 diabetes. *Diabetes*, 52, 573–577. doi:10.2337/diabetes.52.2.573
- Nielsen, E.-M.D., Hansen, L., Carstensen, B., Echwald, S.M., Drivsholm, T., Glümer, C., Thorsteinsson, B., Borch-Johnsen, K., Hansen, T., Pedersen, O., 2003. The E23K variant of Kir6.2 associates with impaired post-OGTT

serum insulin response and increased risk of type 2 diabetes. *Diabetes* 52, 573–577.

- Nielsen, L.B., Ploug, K.B., Swift, P., Ørskov, C., Jansen-Olesen, I., Chiarelli, F., Holst, J.J., Hougaard, P., Pörksen, S., Holl, R., Beaufort, C. de, Gammeltoft, S., Rorsman, P., Mortensen, H.B., Hansen, L., Group, on behalf of the H.S., 2007. Co-localisation of the Kir6.2/SUR1 channel complex with glucagon-like peptide-1 and glucose-dependent insulinotropic polypeptide expression in human ileal cells and implications for glycaemic control in new onset type 1 diabetes. *Eur. J. Endocrinol.* 156, 663–671. doi:10.1530/EJE-06-0756
- Nurwanti, E. (2013). *Paparan Iklan Junk Food Dan Pola Konsumsi Junk Food Sebagai Faktor Resiko Terjadinya Obesitas Pada Anak Sekolah Dasar Kota Dan Desa Di Daerah Istimewa Yogyakarta*. Thesis. FK UGM.
- Ochoa, M. C., Azcona, C., Biebermann, H., Brumm, H., Razquin, C., Wermter, A. K. Pelach, R. (2007). A novel mutation Thr162Arg of the melanocortin 4 receptor gene in a Spanish children and adolescent population. *Clinical Endocrinology*, 66, 652–658. doi:10.1111/j.1365-2265.2007.02788.x
- Ochoa, M. C., Marti A, & Martinez JA (2004) Obesity studies in candidate genes. *Med Clin (Barc)* 122, 542–551.
- Ochoa, M.C., Santos, J.L., Azcona, C., Moreno-Aliaga, M.J., Martínez-González, M.A., Martínez, J.A., Marti, A., 2007. Association between obesity and insulin resistance with UCP2–UCP3 gene variants in Spanish children and adolescents. *Mol. Genet. Metab.* 92, 351–358. doi:10.1016/j.ymgme.2007.07.011
- Oguzkan-Balci, S., Col-Araz, N., Nacak, M., Araz, M., Sabanci, H., Balat, A., Pehlivan, S., 2012. Mitochondrial uncoupling protein 2 (UCP2) gene polymorphisms are associated with childhood obesity and related metabolic disorders. *J. Pediatr. Endocrinol. Metab.* 0, 1–7. doi:10.1515/jpem-2012-0267
- Oktavianthi S, Trimarsanto H, Febinia CA, Suastika K, Saraswati MR, Dwipayana P, Arindrarto W, Sudoyo H, Malik, SG. (2012). Uncoupling protein 2 gene polymorphisms are associated with obesity. *Cardiovascular Diabetology* 11:41.
- Olson, T.M., Terzic, A., 2010. Human KATP channelopathies: diseases of metabolic homeostasis. *Pflüg. Arch. - Eur. J. Physiol.* 460, 295–306. doi:10.1007/s00424-009-0771-y
- Oppert JM, Vohl MC, Chagnon M, Dionne FT, Cassad-Doulcier AM, Ricquier D, Perusse L, Bouchard C. (1994). DNA Polymorphism in the Uncoupling Protein (UCP) Gene and Human Body Fat. *Int J Obes Relat Metab Disord* 18:526-531.

- Özer, S., Yılmaz, R., Kazancı, N.Ö., Sönmezgöz, E., Karaaslan, E., Altuntaş, B., Kuyucu, Y.E., 2015. Higher HDL levels are a preventive factor for metabolic syndrome in obese Turkish children. *LOS NIVELES HDL MÁS ALTOS SON UN FACTOR PARA EL Síndr. METABÓLICO LOS NIÑOS TURCOS QUE SUFREN Obes.* 31, 307–312. doi:10.3305/nh.2015.31.1.7691
- Padez C, Fernandes T, Mourao I, Moreira P, Rosado V. (2004). Prevalence of overweight and obesity in 7–9-year-old Portuguese children: trends in body mass index from 1970–2002. *Am J Hum Biol* 16, 670-678.
- Pick A, Clark J, Kubstrup C, Levisetti M, Pugh W, Boner-Weir S, Polonsky KS. (1998). Role of apoptosis in failure of  $\beta$ -cell mass compensation for insulin resistance and  $\beta$ -cell defects in the male Zucker diabetic fatty rat. *Diabetes* 47:358-364
- Pine DS, Goldstein RB, Wolk S, Weissman MM. (2001). The association between childhood depression and adulthood body mass index. *Pediatrics*; 107: 1049–1056.
- Plourde, G., 2002. Impact of obesity on glucose and lipid profiles in adolescents at different age groups in relation to adulthood. *BMC Fam. Pract.* 3, 18–14.
- Proietto J, Baur LA. (2004) Management of Obesity. *Med J Aust* 180:474-480.
- Proverawati, Atikah. (2010). *Obesitas dan gangguan perilaku makan pada remaja*. Yogyakarta: Nuha Medika.
- Pua YH, Ong PH. (2005). Anthropometric indices as screening tools for cardiovascular risk factors in Singaporean women. *Asia Pac J Clin Nutr* 14: 74-9
- Rahilly, S. (2001). Uncoupling protein 2: Adiposity angel and diabetes devil? *Nature Med.*, 7, 770–2.
- Rathnayake, K.M., Roopasingam, T., Wickramasighe, V.P., 2014. Nutritional and behavioral determinants of adolescent obesity: a case–control study in Sri Lanka. *BMC Public Health* 14, 1291. doi:10.1186/1471-2458-14-1291
- Rennie KL, Jebb SA. (2005). Prevalence of obesity in Great Britain. *Obes Rev* 6:11–12.
- Ricquier D, Bouillaud F. (2000). The uncoupling protein homologues: UCP1, UCP2, UCP3, StUCP and AtUCP. *Biochem J.* 345(2):161-79.
- Riskesdas, (2007), *Laporan Riskesdas*. Badan penelitian dan pengembangan Kementerian Kesehatan RI.
- Riskesdas, (2010), *Laporan Riskesdas*. Badan penelitian dan pengembangan Kementerian Kesehatan RI.

- Riskesmas, (2013), *Laporan Riskesdas*. Badan penelitian dan pengembangan Kementerian Kesehatan RI.
- Robertson, R. P., Harmon, J., Tran, P. O., Tanaka, Y., & Takahashi, H. (2003). Glucose toxicity in beta-cells: type 2 diabetes, good radicals gone bad, and the glutathione connection. *Diabetes*, 52, 581–587. doi:10.2337/diabetes.52.3.581
- Rolls ET. (2005). *Emotion Explained*. Oxford University Press: Oxford.
- Rolls ET. (2007). Understanding the mechanisms of food intake and obesity. *Obesity reviews* 8 (Suppl. 1), 67–72
- Rosen, E.D. & Spiegelman, B.M. (2000). 'Molecular Regulation of Adipogenesis'. *Annu. Rev. Cell Dev. Biol.*, 16: 145–171.
- Rosen, E.D., Hsu, C.H., Wang, X., Sakai, S., Freeman, M.W., Gonzalez, F.J. et al., (2002). 'C/EBP Induces Adipogenesis Through PPAR: A Unified Pathway'. *Genes Dev.*, 16: 22–26.
- Samekto, P T. (2008). *Hubungan aktivitas fisik dengan resistensi insulin pada remaja putri obes*. Skripsi: Universitas Gadjah Mada.
- Santangelo SL, et al. (2005). What is known about autism: genes, brain, and behavior. *Am J Pharmacogenomics*; 5(2): 71-92.
- Sayogo, S. (2006). *Gizi remaja putri*. Jakarta: Fakultas Kedokteran Universitas Indonesia.
- Schmitz, K.H., Jacobs, D.R.Jr., Hong, C.P., Steinberger, J., Moran, A., dan Sinaiko, A.R. (2002). Association of physical activity with insulin sensitivity in children. *International Journal of Obesity*, 26:1310-1316.
- Schwimmer JB, Deutsch R, Rauch JB, Behling C, Newbury R, Lavine JE. (2003). Obesity, insulin resistance, and other clinicopathological correlates of pediatric nonalcoholic fatty liver disease. *J Pediatr*; 143: 500–505.
- Schwimmer JS, Burwinkle TM, Varni JW. (2003) Health-related quality of life of severely obese children and adolescents. *JAMA*; 289: 1813–1819
- Scott, L.J., Bonnycastle, L.L., Willer, C.J., Sprau, A.G., Jackson, A.U., Narisu, N., Duren, W.L., Chines, P.S., Stringham, H.M., Erdos, M.R., Valle, T.T., Tuomilehto, J., Bergman, R.N., Mohlke, K.L., Collins, F.S., Boehnke, M., 2006. Association of Transcription Factor 7-Like 2 (TCF7L2) Variants With Type 2 Diabetes in a Finnish Sample. *Diabetes* 55, 2649–2653. doi:10.2337/db06-0341
- Selby, J.V., Newman, B., Quesenberry, C.P., Jr, Fabsitz, R.R., King, M.C. and Meaney, F.J. (1989). Evidence of genetic influence on central body fat in middle-aged twins. *Hum. Biol.*, 61, 179 – 194.

- Shashikiran U, Sudha V, Jayaprakash B. (2004). What is obesity? *Med J Malaysia* 59:130-3.
- Shen, H., Qi, L., Tai, E.S., Chew, S.K., Tan, C.E., Ordovas, J.M., 2006. Uncoupling Protein 2 Promoter Polymorphism -866G/A, Central Adiposity, and Metabolic Syndrome in Asians. *Obesity* 14, 656-661. doi:10.1038/oby.2006.74
- Shyng, S. L., Ferrigni, T., Shepard, J. B., Nestorowicz, a, Glaser, B., Permutt, M. a, & Nichols, C. G. (1998). Functional analyses of novel mutations in the sulfonylurea receptor 1 associated with persistent hyperinsulinemic hypoglycemia of infancy. *Diabetes*, 47(7), 1145-51. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9648840>
- Silva, D., Matos, A., Magalhães, T., Martins, V., Ricardo, L., Almeida, H., 2012. Prevalência de hipertensão arterial em adolescentes portuguesas da cidade de Lisboa. *Rev. Port. Cardiol.* 31, 789-794. doi:10.1016/j.repc.2012.02.022
- Sinaiko, A.R., Donahue, R.P., Jacobs, D.R., Jr., dan Prineas, R.J. (1999). Relation of Weight and Rate of Increase in Weight during Childhood and Adolescence to Body Size, Blood Pressure, Fasting Insulin, and Lipids in Young Adults: The Minneapolis Children's Blood Pressure Study. *Circulation*, 99, 1471-1476.
- Sorof J, Daniels S. (2002) Obesity hypertension in children: a problem of epidemic proportions. *Hypertension*; 40: 441-4417.
- Steinberg HO, Brechtel G, Johnson A, Fineberg N, Baron AD (1994) Insulin-mediated skeletal muscle vasodilation is nitric oxide dependent. A novel action of insulin to increase nitric oxide release. *J Clin Invest* 94:1172-1179.
- Steinberg HO, Paradisi G, Hook G, Crowder K, Cronin J, Baron AD (2000) Free fatty acid elevation impairs insulin-mediated vasodilation and nitric oxide production. *Diabetes* 49:1231-1238.
- Steinberger, J., Moran, A., Hong, C.P., Jacobs, D.R., Jr., dan Sinaiko, A.R. (2001). Adiposity in childhood predicts obesity and insulin resistance in young adulthood. *J Pediatr*, 138, 469-473.
- Sulistyoningrum, E. (2009). *Hubungan sindroma metabolik pada remaja putri obes dengan sindroma metabolik pada orang tua*. Thesis: Universitas Gadjah Mada.
- Sullivan PF, et al. (2003). Genetic and environmental contributions to obesity and binge eating. *Int J Eat Disord*; 33(3): 293-8.
- Sumiarsih, T. (2006). Peran biologis Uncoupling Protein 2 (UCP2) pada obesitas dan diabetes mellitus tipe 2 (DMT2). *Jurnal Kedokteran YARSI*; 14(3): 243-249.

- The International Association for the Study of Obesity and the International Obesity Task Force. (2000). *The Asia-Pacific Perspective: redefining obesity and its treatment*. Australia: IASO and IOTF.
- Tischler. *Neuroendocrinology: Obesity. Lecture 2004; 52:1-12.*
- Tschritter, O., Stumvoll, M., Machicao, F., Holzwarth, M., Weisser, M., Maerker, E., Fritsche, A. (2002). The Prevalent Glu23Lys Polymorphism in the Potassium Inward Rectifier 6.2 (KIR6.2) Gene Is Associated With Impaired Glucagon Suppression in Response to Hyperglycemia. *Diabetes*, 51(9), 2854–2860. <http://doi.org/10.2337/diabetes.51.9.2854>
- Turula, M., Kaprio, J., Rissanen, A. and Koskenvuo, M. (1990). Body weight in the Finnish Twin Cohort. *Diabetes Res. Clin. Pract.* 10 (Suppl. 1), S33–S36.
- Valet, P., Tavernier, G., Castan-Laurell, I., Saulnier-Blache, J.S. & Langin, D. (2002). 'Understanding Adipose Tissue Development from Transgenic Animal Models'. *J. Lipid Res.*, 43: 835–860.
- von Elm, E., Altman DG, Egger M., Pocock SJ., Gotsche PC., Vandenbroucke JP. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies. *Plos Med* 4(10): e296. [10.1371/journal.pmed.0040296](https://doi.org/10.1371/journal.pmed.0040296)
- Wajchenberg, B.L., (2007). Cell Failure in Diabetes and Preservation by Clinical Treatment. *Endocrine Reviews*. 28(2), pp.187–218.
- Wang C, et al. (2007). Adiponectin sensitizes insulin signaling by reducing p70 S6 kinase-mediated serine phosphorylation of IRS-1. *J Biol Chem*; 282(11): 7991-6.
- Wang, Y., Monteiro, C., Popkin, B.M. (2002). Trends of obesity and underweight in older children and adolescents in the United States, Brazil, China, and Russia. *Am J Clin Nutr.* 75, 971-977.
- WHO Expert Consultation. (2004). Appropriate-body mass index for Asian populations and its implications for policy and intervention strategies. *Lancet* 363:157-63.
- Wijga AH, et al. (2010). Comorbidities of obesity in school children: a cross-sectional study in the PIAMA birth cohort. *BMC Public Health*; 10: 184.
- Woodward, Mark. (1999). *Epidemiology: Study Design and Data Analysis*. United States of America: Chapman & Hall/CRC
- World Health Organization. (2000). *Obesity: preventing and managing the global epidemic. (WHO Technical Report Series no. 894)*. Geneva, Switzerland: World Health Organization.

- World Health Organization. (2008). *The global burden of disease: 2004 update*. Geneva, Switzerland: World Health Organization.
- Yeasmin, R., Choudhury, S., Parveen, S., Nahar, N., Begum, K., Akhter, F., 2015. Assessment of overweight, obesity and metabolic variables among students of a private medical college in Dhaka city. *Bangladesh J. Med. Sci.* 14, 53–58. doi:10.3329/bjms.v14i1.21559
- Yi, Y., Dongmei, L., Phares, D.A., Weiss, E.P., Brandauer, J., Hagberg, J.M., 2008. Association between *KCNJ11* E23K genotype and cardiovascular and glucose metabolism phenotypes in older men and women: *KCNJ11* E23K genotype and cardiovascular and glucose metabolism phenotypes. *Exp. Physiol.* 93, 95–103. doi:10.1113/expphysiol.2007.038893
- Yoshinaga M, Yuasa Y, Hatano H, Kono Y, Nomure Y, Oku S et al. (1995) Effect of total adipose weight and systemic hypertension on left ventricular mass in children. *Am J. Cardiol.* 76, 785-787
- Zhang, M., Wang, M., Zhao, Z.-T., 2014. Uncoupling protein 2 gene polymorphisms in association with overweight and obesity susceptibility: A meta-analysis. *Meta Gene* 2, 143–159. doi:10.1016/j.mgene.2013.10.009