



DAFTAR PUSTAKA

- [1] B. H. Urey, K. V Chellappan, E. Erden, and P. Surman, "State of the Art in Stereoscopic and Autostereoscopic Displays," vol. 99, no. 4, 2011.
- [2] M. H. P. H. V. B. W. A. I. J. F. Juola, "Effectiveness of Stereoscopic Displays in Medicine : A Review," vol. 01, 2012.
- [3] A. Zocco, S. Livatino, and L. T. D. Paolis, "Stereoscopic-3D vision to improve situational awareness in military operations," Switzerland: Springer International Publishing, 2014, pp. 351–362.
- [4] B. Mendiburu, *3D Movie Making Stereoscopic Digital Cinema from Script to Screen. .*
- [5] J. Park, H. Park, and M. Kim, "Stereoscopic 3D Model Viewer with Improved Depth Perception for Battlefield Visualization," pp. 377–384.
- [6] F. Newzoo and R. Report, "Preview of the Southeast Asian Games Market Opportunities in the world ' s fastest growing region," pp. 0–12, 2015.
- [7] M. Lambooi and W. I. I, "Visual Discomfort and Visual Fatigue of Stereoscopic Displays : A Review," vol. 53, no. 3, pp. 1–14, 2009.
- [8] K. Ukai and P. A. Howarth, "Visual fatigue caused by viewing stereoscopic motion images : Background , theories , and observations," vol. 29, pp. 106–116, 2008.
- [9] M. Lambooi, M. Fortuin, W. A. Ijsselsteijn, B. J. W. Evans, and I. Heynderickx, "Susceptibility to Visual Discomfort of 3-D Displays by Visual Performance Measures," vol. 21, no. 12, pp. 1913–1923, 2011.
- [10] G. Smith, "Nintendo's New 3D Console 'Makes Gamers Sick and Gives Them Headaches,'" *Mail Online*, 2011. [Online]. Available: <http://www.dailymail.co.uk/sciencetech/article-1371435/Nintendo-3DS-makes-gamers-sick-gives-headaches.html>.
- [11] S. Wibirama and K. Hamamoto, "Investigation of Visually Induced Motion Sickness in Dynamic 3D Contents based on Subjective Judgment , Heart Rate Variability , and Depth Gaze Behavior," *36th Annu. Int. Conf. IEEE Eng. Med. Biol. Soc.*, pp. 4803–4806, 2014.
- [12] V. De Silva, G. S. Member, A. Fernando, S. Member, S. Worrall, and H. K. Arachchi, "Sensitivity Analysis of the Human Visual System for Depth Cues in Stereoscopic 3-D Displays," vol. 13, no. 3, pp. 498–506, 2011.
- [13] J. J. LaViola, "A discussion of cybersickness in virtual environments,"



ACM SIGCHI Bull., vol. 32, no. 1, pp. 47–56, 2000.

- [14] A. Emery, “Playing Video Games Makes Me Sick,” *The Guardian*, 2011. [Online]. Available: <http://www.theguardian.com/lifeandstyle/2011/dec/19/video-games-makes-me-sick>.
- [15] C. T. Guo, C. W. Tsoi, Y. L. Wong, and K. C. Yu, “Visually induced motion sickness during computer game playing,” *Contemp. Ergon. Hum. Factors 2013 Proc. Int. Conf. Ergon. Hum. Factors 2013, Cambridge, UK, 15-18 April 2013*, pp. 51–58, 2013.
- [16] A. G. Solimini, A. Mannocci, D. Di Thiene, and G. La Torre, “A survey of visually induced symptoms and associated factors in spectators of three dimensional stereoscopic movies,” *BMC Public Health*, vol. 12, no. 1, p. 779, 2012.
- [17] S. Bruck and P. A. Watters, “Estimating Cybersickness of Simulated Motion Using the Simulator Sickness Questionnaire (SSQ): A Controlled Study,” pp. 486–488, 2009.
- [18] Aripin, “Pemodelan Karakter Animasi 3D,” *Techno.Com*, vol. 9, no. 2, pp. 53–64, 2010.
- [19] NVIDIA Corporation, “Game with Nvisia 3D Vision.” [Online]. Available: <http://www.nvidia.com/object/3d-vision-games.html>. [Accessed: 11-Jan-2016].
- [20] M. Poyade, A. Reyes-lecuona, and R. Viciania-abad, “Influence of Binocular Disparity in Depth Perception Mechanisms in Virtual Environments,” pp. 13–22, 2009.
- [21] 3D Consortium (3DC) Safety/Guidelines Section, “3DC Safety Guidelines for Dissemination of Human-friendly 3D,” 2011.
- [22] S. Wibirama, “Study on 3D Gaze Measurement And Its Application to Analyze Visually Induced Motion Sickness in Stereoscopic Environment,” Tokai University, 2014.
- [23] M. B. Flanagan, J. G. May, and T. G. Dobie, “The role ofvection, eye movements and postural instability in the etiology of motion sickness.,” *J. Vestib. Res.*, vol. 14, no. 4, pp. 335–46, 2004.
- [24] P. A. Howarth, “Potential hazards of viewing 3-D stereoscopic television, cinema and computer games: a review,” *Ophthalmic Physiol. Opt.*, vol. 31, no. 2, pp. 111–122, 2011.



- [25] A. Iijima, S. Komagata, T. Kiryu, T. Bando, and I. Hasegawa, "Vergence eye movements signifying 3D depth perception from 2D movies," *Displays*, vol. 33, no. 2, pp. 91–97, 2012.
- [26] R. S. Kennedy, N. E. Lane, S. Berbaum, and M. G. Lilienthal, "The International Journal of Aviation Psychology Simulator Sickness Questionnaire : An Enhanced Method for Quantifying Simulator Sickness," no. February 2013, pp. 203–220, 2009.
- [27] R. S. Kennedy, D. Ph, and K. S. Berbaum, "Disorientation and Postural Ataxia Following Flight Simulation," no. December 1995, pp. 13–17, 1997.
- [28] R. S. Kennedy, J. Drexler, and R. C. Kennedy, "Research in visually induced motion sickness," *Appl. Ergon.*, vol. 41, no. 4, pp. 494–503, 2010.
- [29] C. Jinjakam and K. Hamamoto, "Study on Parallax Affect on Simulator Sickness in One-screen and Three-screen Immersive Virtual Environment," *Proc. Sch. Inf. Telecommun. Eng. Tokai Univ.*, vol. 4, no. 4, pp. 34–39, 2011.
- [30] K. Benzeroual and R. S. Allison, "Cyber (Motion) Sickness In Active Stereoscopic 3D Gaming," pp. 1–7, 2013.
- [31] W. Fei, "A study of stereoscopic display technology and visual fatigue caused by viewing stereoscopic images," 2011.
- [32] S. Wibirama and K. Hamamoto, "Paper Development of Gaze Tracking System for Stereoscopic Virtual Environment by," vol. 5, no. 1, pp. 1–8, 2012.
- [33] S. T. Moore, T. Haslwanter, I. I. A. N. S. Curthoys, and S. T. Smiths, "A Geometric Basis for Measurement of Three-dimensional Eye Position Using Image Processing," vol. 36, no. 3, 1996.
- [34] S. Zugal and J. Pinggera, "Low – Cost Eye – Trackers: Useful for Information Systems Research ?," 2014.
- [35] J. W. Creswell, *Educational Research: Planning, Conducting, and Evaluating Quantitative and Qualitative Research*, Fourth Edi. Boston: Pearson Education, Inc., 2012.
- [36] Lund Research Ltd, "Two-way repeated measures ANOVA using SPSS Statistics," 2013. [Online]. Available: <https://statistics.laerd.com/spss-tutorials/two-way-repeated-measures-anova-using-spss-statistics.php>. [Accessed: 11-Jan-2016].
- [37] N. Setiawan, "Analisis Ragam / Analysis of variance (Anova) dua arah



- dengan interaksi,” *Statistik Ceria*, 2014. [Online]. Available: <http://statistikceria.blogspot.co.id/2014/01/analisis-ragam-analysis-of-variance-anova-dua-arah-dengan-interaksi.html>. [Accessed: 11-Jan-2016].
- [38] IBM Corp, “IBM SPSS Statistics for Windows [Computer software].” IBM Corp, Armonk, NY, 2013.
- [39] H. Coolican, *Research Methods and Statistics in Psychology*, 5th Editio. New York: Routledge, 2013.
- [40] A. G. Solimini, “Are There Side Effects to Watching 3D Movies? A Prospective Crossover Observational Study on Visually Induced Motion Sickness,” *PLoS One*, vol. 8, no. 2, p. e56160, 2013.
- [41] R. V. Lenth, “Java Applets for Power and Sample Size [Computer software].” 2011.
- [42] C. R. W. Vanvoorhis and B. L. Morgan, “Understanding Power and Rules of Thumb for Determining Sample Sizes,” vol. 3, no. 2, 2007.
- [43] S. Palmisano, F. Bonato, and A. Bubka, “Vertical Display Oscillation Effects on Forward Vection and Simulator Sickness,” vol. 78, no. 10, pp. 951–956, 2007.
- [44] S. Hu and R. M. Stern, “Optokinetic nystagmus correlates with severity of vection-induced motion sickness and gastric tachyarrhythmia,” *Aviat. Space. Environ. Med.*, vol. 69, no. 12, pp. 1162–1165, 1998.