



DAFTAR PUSTAKA

- Appelros, P. (2007). Prediction of length of stay for stroke patients. Acta Neurologica Scandinavica, 116, 15-19. Doi: 10.1111/j.1600-0404.2006.00756.x
- Arboix, A., Massons, J., Gracia-Eroles, L., Oliveres, M., & Comes, E. (2012). Clinical predictors of prolonged hospital stay after acute stroke: relevance of medical complications. International Journal of Clinical Medicine, 3, 502-507. Doi: 10.4236/ijcm.2012.36090
- Badan Penelitian dan Pengembangan Kesehatan. (2013). Riset Kesehatan Dasar 2013. Kementerian Kesehatan RI.
- Briggs, D.E., Felberg, R.A., Malkoff, M.D., Bratina, P., & Grotta, J.C. Should mild or moderate stroke patients be admitted to an intensive care unit. Stroke, 32, 871-876. Doi: 10.1161/01.STR.32.4.871
- Cooper, C. (2015, Mei 12). Sharp rise in strokes among people of working age leaves thousands devastated by 45 isability. The Independent. Diakses pada Juni 15, 2015, dari <http://www.independent.co.uk/life-style/health-and-families/health-news/sharp-rise-in-strokes-among-people-of-working-age-leaves-thousands-devastated-by-disability-10242663.html>
- Giraldo, E. (2013a). Intracerebral Hemorrhage. Merckmanuals.com. Retrieved 16 June 2015, from <https://www.merckmanuals.com/professional/neurologic-disorders/stroke-cva/intracerebral-hemorrhage>
- Giraldo, E. (2013b). Subarachnoid Hemorrhage (SAH). Merckmanuals.com. Retrieved 16 June 2015, from <https://www.merckmanuals.com/professional/neurologic-disorders/stroke-cva/subarachnoid-hemorrhage-sah>
- Halleivi, H., Albright, K.C., Aronowski, J., Barreto, A.D., Martins-Schild, S., Khaja, A.M., et al. (2008). Intraventricular hemorrhage: anatomic relationships and clinical implications. Neurology, 70(11), 848-852. Doi: 10.1212/01.wnl.0000304930.47751.75
- Herminawati, A., Suryani, M., & Sayono. (2013). Perbedaan lama rawat inap antara stroke hemoragik dan stroke non hemoragik di RSUD Tugurejo Semarang. Skripsi. STIKES Telogorejo Semarang.



- Kim, S.M., Hwang, S.W., Oh, E.W., & Kang, J.K. (2013). Determinants of the length of stay in stroke patients. Osong Public Health and Research Perspectives, 4(6), 342-346. Doi: 10.1016/j.phrp.2013.10.008
- Kumar, V., Abbas, A., & Aster, J. (2013). Robbins basic pathology (9<sup>th</sup> ed.). Philadelphia: Elsevier Saunders.
- Lim, J. & Cheon, S. (2015). Analysis of variation in length of stay (LOS) after ischemic and hemorrhagic stroke using the Charlson Comorbidity Index (CCI). Journal of Physical Therapy Science, 27(3), 799-803. Doi: 10.1589/jpts.27.799
- Liu, Y., Phillips, M., & Codde, J. (2001). Factors influencing patients' length of stay. Australian Health Review, 24(2), 63-70. Doi: 10.1071/AH010063
- Martono, H., & Kuswardani, RA. (2010). Strok dan penatalaksanaannya oleh internis. In A. Sudoyo, B. Setiyohadi, I. Alwi, M. Simadibrata K., & S. Setiati (Eds.), Buku ajar ilmu penyakit dalam (5<sup>th</sup> ed., pp. 892 - 903). Jakarta: InternaPublishing.
- Meschia, J., Bushnell, C., Boden-Albala, B., Braun, L., Bravata, D., & Chaturvedi, S. et al. (2014). Guidelines for the Primary Prevention of Stroke: A Statement for Healthcare Professionals From the American Heart Association/American Stroke Association. Stroke, 45(12), 3754-3832. Doi:10.1161/str.0000000000000046
- Morgenstern, L., Hemphill, J., Anderson, C., Becker, K., Broderick, J., & Connolly, E. et al. (2010). Guidelines for the management of spontaneous intracerebral hemorrhage: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. Stroke, 41(9), 2108-2129. Doi:10.1161/str.0b013e3181ec611b
- Nakao, S., Takata, S., Uemaru, H., Kashiwara, M., Osawa, T., Komatsu, K., et al. (2010). Relationship between Barthel Index scores during the acute phase of rehabilitation and subsequent ADL in stroke patients. The Journal of Medical Investigation, 57, 81-88.
- National Collaborating Centre for Chronic Conditions. (2008). Stroke: national clinical guideline for diagnosis and initial management of acute stroke



- and transient ischemic attack (TIA). London: Royal College of Physicians.
- Pan, X., Simon, T.A., Hamilton, M., & Kuznik, A. (2015). Comparison of costs and discharge outcomes for patients hospitalized for ischemic or hemorrhagic stroke with or without atrial fibrillation in the United States. Journal of Thrombosis and Thrombolysis, 39, 508-515. Doi: 10.1007/s11239-014-1144-8
- Ropper, A., Samuels, M., & Klein, J. (2014). Adam and Victor's principles of neurology (10<sup>th</sup> ed.). McGraw-Hill Education.
- Schlegel, D., Kolb, S.J., Luciano, J.M., Tovar, J.M., Cucchiara, B.L., Liebeskind, D.S., et al. (2003). Utility of the NIH Stroke Scale as predictor of hospital disposition. Stroke, 34, 134-137. Doi: 10.1161/01.STR.0000048217.44714.02
- Smith, W., English, J., & Claiborne Johnston, S. (2012). Cerebrovascular disease. In D. Longo, A. Fauci, D. Kasper, S. Hauser, J. Jameson & J. Loscalzo (Eds.), Harrison's principles of internal medicine (18<sup>th</sup> ed., pp. 3270 - 3299). New York: McGraw Hill Medical.
- Steiner, T., Al-Shahi Salman, R., Beer, R., Christensen, H., Cordonnier, C., & Csiba, L. et al. (2014). European Stroke Organisation (ESO) guidelines for the management of spontaneous intracerebral hemorrhage. International Journal Of Stroke, 9(7), 840-855. Doi:10.1111/ijs.12309
- Sulter, G., Steen, C., & De Keyser, J. (1999). Use of the Barthel index and modified rankin scale in acute stroke trials. Stroke, 30, 1539-1541. Doi: 10.1161/01.STR.30.8.1538
- Zhang, J., Wang, Y., Wang, G., Sun, H., Sun, T., & Shi, J. et al. (2011). Clinical factors in patients with ischemic versus hemorrhagic stroke in East China. World Journal Of Emergency Medicine, 2(1), 18 - 23. Doi:10.5847/wjem.j.1920-8642.2011.01.003