

REFERENCES

- Abbasi, T., Tauseef, S.M., Abbasi, S.A. (2012). *Biogas Energy*. Springer Business Media, New York.
- Ademark, P., Varga, A., Medve, J., Harjunpaa, V., Drakenberg, T., Tjerneld, F., and Stalbrand, H. (1998). Softwood hemicellulose-degrading enzymes from *Aspergillus niger*: Purification and properties of a beta-mannanase. *Journal of Biotechnology* 63(3): 199-210.
- Andre, L., Durante, M., Pauss, A., Lespinard, O., Ribeiro, T., Lamy, E. (2015). Quantifying physical structure changes and non-uniform water flow in cattle manure during dry anaerobic digestion process at lab scale: Implication for biogas production. *Bioresource Technology* 192: 660–669.
- APHA. (2005). *Standard Methods for the Examination of Water and Wastewater*. 21st ed. Ed. American Public Health Association, Washington DC.
- Altschul, S.F., Gish, W., Miller, W., Myers E.W., Lipman, D.J (1990). Basic local alignment search tool. *J Mol Biol* 215: 403–10.
- Batstone, D.J., Keller, J., Angelidaki, I.S.V., Pavlostathis, S.G., Rozzi, A., Sanders, W.T.M., et al. (2002). *Anaerobic Digestion Model No. 1. A task group on mathematical modeling of anaerobic digestion process*. IW A Scientific and Technical Report No. 13: IW A Publishing, London.
- Boone, D.R. (1982). *Appl. Environ. Microbiol.* 43, 57-64
- Brummeler, E, Horbach, H.C.J.M., and Koster, I.W. (1991). Dry Anaerobic Batch Digestion of the Organic Fraction of Municipal Solid Waste. *J . Chem. Tech. Biotechnol.* 50: 191-209.
- Buffiere, P., Loisel, D., Ndelgenes, B., Delgenes, J.P. (2006). Towards new indicators for the prediction of solid waste anaerobic digestion properties. *Water Sci Technol*; 53:233-41.
- Chanakya, H.N., Srikumar, K.G., Anand, V., Modak, J., and Jagadish, K.S. (1999). Fermentation properties of agro-residues, leaf biomass and urban market garbage in a solid phase biogas fermenter. *Biomass and Bioenergy* 16: 417-429.
- Cheng, J. (2010). *Biomass to renewable energy processes*. CRC Press, New York.

- Chynoweth, D.P., Turick, C.E., Owens, J.M., Jerger, D.E., Peck, M.W. (1993). Biochemical methane potential of biomass and waste feedstocks. *Biomass Bioenergy* 5: 95-111.
- Colazo, A., Sánchez, A., Font, X., Colón, J. (2015). Environmental impact of rejected materials generated in organic fraction of municipal solid waste anaerobic digestion plants: Comparison of wet and dry process layout. *Waste Management* 43: 84–97.
- Côté, C., Massé, D.I., Quessy, S. (2006). Reduction of indicator and pathogenic microorganisms by psychrophilic anaerobic digestion in swine slurries. *Bioresour Tech* 97, 686–91.
- De Baere, L., and Mattheeuws, B. (2010). Anaerobic Digestion of MSW in Europe. *Biocycle. February*: 24-26.
- Delmer, D.P., and Amor, Y. (1995). Cellulose Biosynthesis. *Plant Cell* 7: 987-1000.
- Deublein, D., and Angelika, S. (2008). *Biogas from waste and renewable resources*. WILEY-VCH Verlag GmbH & Co. KgaA, Weinheim.
- Dhaked, R.K., Singh, P., Singh, L. (2010). Biomethanation under psychrophilic conditions. *Waste Manag* 30: 2490–6.
- Emmel, A., Mathias, A.L., Wypych, F., and Ramos, L.P. (2003). Fractionation of Eucalyptus grandis chips by dilute acid-catalysed steam explosion. *Bioresource Technology* 86: 105-115.
- Fauzi, Y. (2005). *Kelapa Sawit, Budi Daya Pemanfaatan Hasil dan Limbah, Analisa Usaha dan Pemasaran*. Penebar Swadaya, Jakarta.
- Fernández-Rodríguez, J., Pérez, M., Romero, L.I. (2014). Dry thermophilic anaerobic digestion of the organic fraction of municipal solid wastes: Solid retention time optimization. *Chemical Engineering Journal* 251: 435–440.
- Fox, M.H., Noike, T., and Ohki, T. (2003). Alkaline subcritical-water treatment and alkaline heat treatment for the increase in biodegradability of newsprint waste. *Water Science and Technology* 48: 77-84.
- Frigon, J.C., Mehta, P., and Guiot, S.R. (2011). *Impact of mechanical, chemical and enzymatic pre-treatments on the methane yield from the anaerobic digestion of switchgrass*. Biomass and Bioenergy In Press, Corrected Proof.

- Gerardi, M.H. (2003). *The microbiology of anaerobic digesters*. Wiley, John & Sons, New York.
- Gomec, C.Y., Letsiou, I., Ozturk, I., Eroglu, Y., Wilderer, P.A. (2008). Identification of Archaeal population in the granular sludge of an UASB reactor treating sewage at low temperatures. *J. Environ. Sci. Health Pt. A* 43: 1504–1510.
- Guendouz, J., Buffire, P., Cacho, J., Carrre, M., and Delgenes, J.P. (2010). Dry anaerobic digestion in batch mode: Design and operation of a laboratory-scale, completely mixed reactor. *Waste Management* 30: 1768-1771.
- Hansen, K.H., Angelidaki, I., Ahring, B.K. (1998). Anaerobic digestion of swine manure: inhibition by ammonia. *Water Res* 3:, 5–12.
- Hashimoto, A.G. (1986). Pretreatment of wheat straw for fermentation to methane. *Biotechnol. Bioeng.* 28: 1857-1866.
- Himmel, M.E., Adney, W.S., Rivard, C.J., Grohmann, K. (1989). *Detection of Extra-Cellular Hydrolytic Enzymes in Anaerobic Digestion of MSW*. Solar Energy Research Institute. SERI/SP-231-3520. Pp. 9-23.
- Huber, H., Thomm, M., Konig, H., Thies, G., Stetter, K.O. (1982). *Methanococcus hermolithotrophicus*, a novel thermophilic lithotropic methanogen. *Arch Microbiol* 132: 47-50.
- Jewell, W.J., Cummings, R.J., and Richards, B.K. (1993). Methane fermentation of energy crops: Maximum conversion kinetics and in situ biogas purification. *Biomass and Bioenergy* 5: 261-278.
- Kaparaju, P., Buendia, I., Ellegaard, L., Angelidakia, I. (2008). Effects of mixing on methane production during thermophilic anaerobic digestion of manure: Lab-scale and pilot-scale studies. *Bioresource Technology* 99: 4919-4928.
- Kerdsuwan, S., Laohalidanond, K. (2011). *Renewable energy from palm oil empty fruit bunch*. In: Nayeripour M, editor. *Renewable energy: trends and applications*. Shanghai: InTech. pp. 50-123.
- Kim, J., Park, C., Kim, T.H., Lee, M., Kim, S., Kim, S.W., et al. (2003). Effects of various pretreatments for enhanced anaerobic digestion with waste activated sludge. *J Biosci Bioeng* 95: 271-5.
- Kothari, R., Kumar, V., Tyagi, W. (2011). *Assessment of waste treatment and energy recovery from dairy industrial waste by anaerobic digestion*. *Environ Manag Sustain Dev*;2:1-6 (IIOAB special issue).

- Kotsyurbenko, O.R., Nozhevnikova, A.N., Zavarzin, G.A. (1993). Methanogenic degradation of organic matter by anaerobic bacteria at low temperature. *Chemosphere* 27: 1745–61.
- Lavarack, B.P., Griffin, G.J., and Rodman, D. (2002). The acid hydrolysis of sugarcane bagasse hemicellulose to produce xylose, arabinose, glucose and other products. *Biomass & Bioenergy* 3: 367-380.
- Lehtomäki, A. (2006). *Biogas production from energy crops and crop residues*. University of Jyväskylä.
- Li, C., Mörtelmaier, C., Winter, J., Gallert, C. (2014). Effect of moisture of municipal biowaste on start-up and efficiency of mesophilic and thermophilic dry anaerobic digestion. *Bioresource Technology* 168: 23–32.
- Li, Y., Park, S.Y, and Zhu, J. (2011). Solid-state anaerobic digestion for methane production from organic waste. *Renewable and Sustainable Energy Reviews* 15: 821–826.
- Liew, L.N., Jian, S., Yebo, L. (2012). Methane production from solid-state anaerobic digestion of lignocellulosic biomass. *Biomass and Bioenergy* 46:125-132.
- Liu, C., Yuan, X., Zeng, G., Li, W., Li, J. (2008). Prediction of methane yield at optimum pH for anaerobic digestion of organic fraction of municipal solid waste. *Bioresource Technol*; 99: 882-8.
- Lübken, M., Koch, K., Gehring, T., Wichern, M., and Horn, H. (2010). Biogas from grass silage - Measurements and modeling with ADM1. *Bioresource Technology* 101: 8158-8165.
- Madigan, M.T., Martinko, J.M., Dunlap, P.V., Clark, D.P. (2009). *Brock biology of microorganisms*. 12th ed. Benjamin Cummings.
- Mansor, M., Ariffin, W.T.W. (2012). *More to oil palm than just palm oil*. [Internet]. New Straits Times; 2012 Mar 24 [cited 2015 Nov 17]. Available from: http://www.btimes.com.my/Current_News/BTIMES/articles/fleda2/Article/.
- Mass'e, D.I., Masse, L. (2000). The effect of temperature on slaughterhouse wastewater treatment in anaerobic sequencing batch reactors. *Bioresour Tech* 76: 91–8.

- Massé, D.I., Masse, L., Croteau, f. (2003). The effect of temperature fluctuations on psychrophilic anaerobic sequencing batch reactors treating swine manure. *Bioresource Technology*. 89: 57-62.
- Massé, D.I., Saady, N.M.C. (2015). Psychrophilic dry anaerobic digestion of dairy cow feces: Long-term operation. *Waste Management* 36: 86–92.
- Mod, R.R., Ory, R.L., Morris, N.M., and Normand, F.L. (1981). Chemical-Properties and Interactions of Rice Hemicellulose with Trace Minerals Invitro. *Journal of Agricultural and Food Chemistry* 29: 449-454.
- Mohammad, N., Alam, M.Z., Kabbashi, N.A., Ahsan, A. (2011). Effective composting of oil palm industrial waste by filamentous fungi: a review. *Resour Conservat Recycl* 58: 69-78.
- Mohammed, M.A.A., Salmiaton, A., Wan, A.W.A.K.G., Mohamad, A.M. (2012). Gasification of oil palm empty fruit bunches: a characterization and kinetic study. *Bioresour technol* 110: 628-36.
- Mora-Naranjo, N., Meima, J.A., Haarstrick, A., Hempel, D.C. (2004). Modelling and experimental investigation of environmental influences on the acetate and methane formation in solid waste. *Waste Manag.* 24: 763–773.
- Morohoshi, N. (1991). *Chemical characterization of wood and its components*. In Wood and cellulosic chemistry, 331-392. Hon, D. N. S., and Shiraishi, N., eds. Marcel Dekker, Inc, New York.
- Mosier, N., Wyman, C., Dale, B., Elander, R., Lee, Y.Y., Holtzapple, M., and Ladisch, M. (2005). Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresource Technology* 96: 673-686.
- National Renewable Energy Laboratory (NREL). 1992. *US Data Summary of Municipal Waste Management Alternatives*. X. Colorado.
- O'Dwyer, M.H. (1934). The hemicelluloses of the wood of English oak: the composition and properties of hemicellulose, isolated from samples of wood dried under various conditions. *Biochemistry Journal* 28: 2116-2124.
- Omar, R., Idris, A., Yunus, R., Khalid, K., Aida, I.M.I. (2011). Characterization of empty fruit bunch for microwave assisted pyrolysis. *Fuel*; 90:1536-44.
- Persson, T., Matusiak, M., Zacchi, G., and Jonsson, A.S. 2006. Extraction of hemicelluloses from process water from the production of masonite. *Desalination* 199: 411-412.

- Pommier, S., Chenu, D., Quintard, M., Lefebvre, X. (2007). A logistic model for the prediction of the influence of water on the solid waste methanization in landfills. *Biotechnol. Bioeng.* 97: 473–482.
- Rapport, J., Zhang, R., Jenkins, B.M., Williams, R.B. (2008). *Current anaerobic digestion technologies used for treatment of municipal organic solid waste*. California Environmental Protection Agency. [Internet] [cited 2014 Dec 14]. Available from: <http://www.calrecycle.ca.gov/Publications/Documents/1275/2008011.pdf>.
- Romano, R.T, Ruihong, Z., Sarah, T., Jeffery, A.M. (2009). The effect of enzyme addition on anaerobic digestion of Jose Tall Wheat Grass. *Bioresource Technology* 100: 4564-4571.
- Rubin, E.M. (2008). Genomics of cellulosic biofuels. *Nature*; 454: 841-5.
- Saady, N.M.C., Massé, D.I. (2015). Impact of organic loading rate on the performance of psychrophilic dry anaerobic digestion of dairy manure and wheat straw: Long-term operation. *Bioresource Technology* 182: 50–57.
- Schäfer, W., Lehto, M., and Teye, F. *Dry anaerobic digestion of organic residues on-farm —a feasibility study*. [Internet] [cited 2014 Dec 14]. Available from: <http://www.mtt.fi/met/pdf/met77.pdf>.
- Sharma, S.K., Mishra, I.M., Sharma, M.P., and Saini, J.S. (1988). Effect of Particle-Size on Biogas Generation from Biomass Residues. *Biomass* 17: 251-263.
- Shewani, A., Horgue, P., Pommier, S., Debenest, G., Lefebvre, X., Gandon, E., Paul, E. (2015). Assessment of percolation through a solid leach bed in dry batch anaerobic digestion processes. *Bioresource Technology* 178: 209–216.
- Simankova, M.V., Kotsyurbenko, O.R., Lueders, T., Nozhevnikova, A.N., Wagner, B., Conrad, R., Friedrich, M.W. (2003). Isolation and characterization of new strains of methanogens from cold terrestrial habitats. *Syst Appl Microbiol* 26: 312-318.
- Sluiter, A., Hames, B., Ruiz, R., Scarlata, C., Sluiter, J., Templeton, D., Croker, D. 2011. Laboratory Analytical Procedure (LAP): Determination of Structural Carbohydrate and Lignin in Biomass. Golden, CO, USA: National Renewable Energy Laboratory.
- Sponza, D.T., Ağdağ, O.N., (2004). Impact of leachate recirculation and recirculation volume on stabilization of municipal solid wastes in simulated anaerobic bioreactors. *Process Biochem* 39: 2157–2165.

- Sun, Y., and Cheng, J. (2002). Hydrolysis of lignocellulosic materials for ethanol production: a review. *Bioresource Technology* 83: 1-11.
- Taherzadeh, M.J., and Karimi, K. (2008). Pretreatment of Lignocellulosic Wastes to Improve Ethanol and Biogas Production: A Review. *International Journal of Molecular Sciences* 9: 1621-1651.
- Teh, C.B.S., Goh, K.J., Kamarudin, K.N. (2010). Physical changes to oil palm empty fruit bunches (EFB) and EFB mat (Ecomat) during their decomposition in the field. *Pertanika J Trop Agric Sci.* 33: 39-44.
- Themelis, N.J., Verma, S. (2004). *The better option: anaerobic digestion of organic waste in MSW*. Waste Manag World.
- Tong, X., Smith, L.H., and McCarty, P.L. (1990). Methane Fermentation of Selected Lignocellulosic Materials. *Biomass* 21: 239-255.
- Vandevivere, P., Baere, L., and Verstraete, W. (2002). *Types of anaerobic digesters for solid wastes*. In: Mata-Alvarez J. (Editor): Biomethanization of the Organic Fraction of Municipal Solid Wastes, London.
- Vavilin, V.A., Lokshina, L.Y., Rytov, S.V., Kotsyurbenko, O.R., Nozhevnikova, A.N., Parshina, S.N. (1997). Modelling methanogenesis during anaerobic conversion of complex organic matter at low temperatures. *Water Sci. Technol.* 36: 531–538.
- Wellinger, A. (1999). *Process design of agricultural digesters*. Nova Energie GmbH Elggerstrasse, Report.
- Wujcik, W.J., and Jewell, W.J. (1980). Dry Anaerobic Fermentation. Biotechnology and Bioengineering. Wiley & Sons, Inc, New York.
- Yang, S.T., Okos, M.R. (1987). Kinetic study and mathematical modeling of methanogenesis of acetate using pure culture of methanogens. *Biotechnol Bioeng* 30: 661-7.
- Yusoff, S. (2006). Renewable energy from palm oil: innovation on effective utilization of waste. *J Clean Prod* 14(1):87-93.
- Zahedi, S., Sales, D., Romero, L.I., Solera, R. (2013). Optimisation of the two-phase dry-thermophilic anaerobic digestion process of sulphate-containing municipal solid waste: Population dynamics. *Bioresource Technology* 148: 443–452.
- Zhang, Y.H.P., & Lynd, L.R. (2004). Toward an aggregated understanding of enzymatic hydrolysis of cellulose: noncomplexed cellulase systems. *Biotechnol Bioeng* 88: 797–824.

- Zhao, H.W., Viraraghavan, T. (2004). Analysis of the performance of an anaerobic digestion system at the Rigna wastewater treatment plant. *Bioresour Tech* 95: 301–7.
- Zheng, Y., Pan, Z., and Zhang, R. (2009). Overview of biomass pretreatment for cellulosic ethanol production. *Int J Agric & Biol Eng* 2: 51-68.
- Zhu, G., and Jha, A.K. (2013). Psychrophilic dry anaerobic digestion of cow dung for methane production: Effect of inoculum. *ScienceAsia* 39: 500-510.
- Zhu, J., Cai, W., Li, Y. (2010). Enhanced solid-state anaerobic digestion of corn stover by alkaline pretreatment. *Bioresour Technol* 101: 7523-7528.