

DAFTAR PUSTAKA

- Belliere, J., Casemayou, A., Ducasse, L., Zakaroff-Girard, A., Martins, F., Iacovoni, J.S., et al. Specific macrophage subtypes influence the progression of rhabdomyolysis-induced kidney injury. *J Am Soc Nephrol* 2015; 26(6): 1363-1377.
- Billiet, L., Doaty, S., Katz, J.D. and Velasquez, M.T. Review of hyperuricemia as new marker for metabolic syndrome. *ISRN Rheumatol* 2014; 852954: 1-7.
- Bobulescu, I.A. and Moe, O.W. Renal transport of uric acid: evolving concepts and uncertainties. *Adv Chronic Kidney Dis* 2012; 19(6): 358–371.
- Brunton, L.L., Chabner, B.A. and Knollman, B.C. (editor). Goodman & Gilman's the pharmacological basis of therapeutics, 12th ed. New York : McGraw-Hill, 2011.
- Carew, R.M., Wang, B. and Kantharidis, P. The role of EMT in renal fibrosis. *Cell Tissue Res* 2012; 347(1): 103–116.
- Chaudhary, K., Malhotra, K., Sowers, J. and Aroor, A. Uric acid – key ingredient in the recipe for cardiorenal metabolic syndrome. *Cardiorenal Med* 2013;3:208–220.
- Chia, R., Achilli, F., Festing, M.F.W., and Fisher, E.M.C. The origins and uses of mouse outbred stocks. *Nat Genet* 2005; 37(11):1181-1186.
- Cochrane, A.L., Kett, M.M., Samuel, C.S., Campanale, N.V., Anderson, W.P., Hume, D.A., et al. Renal structural and functional repair in a mouse model of reversal of ureteral obstruction. *J Am Soc Nephrol* 2005; 16: 3623–3630.
- Correa-Costa, M., Braga, T.T., Semedo, P., Hayashida, C.Y., Bechara, L.R.G., Elias, R.M., et al. Pivotal role of toll-like receptors 2 and 4, its adaptor molecule MyD88, and inflammasome complex in experimental tubule-interstitial nephritis. *PLoS ONE* 2011; 6(12): e29004.
- Darmawan, J., Valkenburg, H.A., Muirden, K.D. and Wigley, R.D. The epidemiology of gout and hyperuricemia in a rural population of Java. *J Rheumatol* 1992; 19(10): 1595-9.
- David dan Arkeman, H. Evaluation of the oral toxicity of formaldehyde in rats. *Univ Med* 2008; 27(3): 106-112.
- Derbre, F., Ferrando, B., Gomez-Cabrera, M.C., Sanchis-Gomar, F., Martinez-Bello, V.E., Olaso-Gonzalez, G., et al. Inhibition of xanthine oxidase by allopurinol prevents skeletal muscle atrophy: role of p38 MAPKinase and E3 ubiquitin ligases. *PLoS ONE* 2012; 7(10): e46668.
- Diwan, V., Mistry, A., Gobe, G. and Brown, L. Adenine-induced chronic kidney and cardiovascular damage in rats. *J Pharmacol Toxicol Method* 2013; 68(2): 197–207.
- Filiopoulos, V., Hadjiyannakos, D. and Vlassopoulos, D. New insights into uric acid effects on the progression and prognosis of chronic kidney disease. *Ren Fail* 2012; 34(4): 510– 520.
- Gustafsson, D. and Unwin R. The pathophysiology of hyperuricaemia and its possible relationship to cardiovascular disease, morbidity and mortality. *BMC Nephrol* 2013; 14: 164.

- Hakoda, M. Recent trends in hyperuricemia and gout in Japan. *Japan Med Assoc J* 2012; 55(4): 319–323.
- Han, Y., Ma, F.Y., Tesch, G.H., Manthey, C.L. and Nikolic-Paterson, D.J. c-fms Blockade reverses glomerular macrophage infiltration and halts development of crescentic anti-GBM glomerulonephritis in the rat. *Lab Invest* 2011; 91: 978–991.
- Han, Y., Ma, F.Y., Tesch, G.H., Manthey, C.L. and Nikolic-Paterson, D. J. Role of macrophages in the fibrotic phase of rat crescentic glomerulonephritis. *Am J Physiol Renal Physiol* 2013; 304: F1043–F1053.
- Harmita dan Radji, M. Buku ajar analisis hayati. Edisi ke-3. Jakarta : EGC, 2008.
- Helal, I., Fick-Brosnahan, G.M., Reed-Gitomer, B., and Schrier, R.W. Glomerular hyperfiltration: definitions, mechanisms and clinical implications. *Nat Rev Nephrol* 2012; 8(5):293-300.
- Henderson, N.C., Mackinnon, A.C., Farnworth, S.L., Kipari, T., Haslett, C., Iredale, J.P, et al. Galectin 3 expression and secretion links macrophages to the promotion of renal fibrosis. *Am J Pathol* 2008; 172: 288–298.
- Ikezumi, Y., Atkins, R.C. and Nikolic-Paterson, D.J. Interferon- γ augments acute macrophage-mediated renal injury via a glucocorticoid-sensitive mechanism. *J Am Soc Nephrol* 2003; 14: 888–898.
- Ikezumi, Y., Suzuki, T., Karasawa, T., Hasegawa, H., Kawachi, H., Nicolice-Paterson, D.J. and Uchiyama, M. Contrasting effects of steroids and mizoribine on macrophage activation and glomerular lesions in rat thy-1 mesangial proliferative glomerulonephritis. *Am J Nephrol* 2010; 31: 273–282.
- Ikezumi, Y., Suzuki, T., Karasawa, T., Hasegawa, H., Yamada, T., Imai, N., et al. Identification of alternatively activated macrophages in new-onset paediatric and adult immunoglobulin A nephropathy: potential role in mesangial matrix expansion. *Histopathol* 2011; 58: 198–210.
- Janeway, C.A. and Medzhitov, R. Innate immune recognition. *Annu Rev Immunol* 2002; 20: 197-216.
- Jin, M., Yang, F., Yang, I., Yin, Y., Luo, J.J., Wang, H. and Yang, X.F. Uric acid, hyperuricemia and vascular diseases. *Front Biosci* 2012; 17: 656–669.
- Johnson, R.J., Nakagawa, T., Jalal, D., S´anchez-Lozada, L.G., Kang, D. H. and Ritz, E. Uric acid and chronic kidney disease: which is chasing which? *Nephrol Dial Transplant* 2013; 28(9): 2221–2228.
- Kanellis, J., Watanabe, S., Li, J.H., Kang, D.H., Li, P., Nakagawa, T., et al. Uric acid stimulates monocyte chemoattractant protein-1 production in vascular smooth muscle cells via mitogen-activated protein kinase and cyclooxygenase-2. *Hypertension* 2003; 41(6): 1287–1293.
- Kang, D.H., Nakagawa, T., Feng, L., Watanabe, S., Han, L., Mazzali, M., et al. A role for uric acid in the progression of renal disease. *J Am Soc Nephrol* 2002; 13(12): 2888–2897.
- Kang, D.H., Park, S.K., Lee, I.K. and Johnson, R.J. Uric acid-induced C-reactive protein expression: implication on cell proliferation and nitric oxide production of human vascular cells. *J Am Soc Nephrol* 2005; 16(12): 3553–3562.

- Katzung, B.G., Masters, S.B., Trevor, A.J. (editor). Basic & clinical pharmacology, 11th ed. New York : McGraw-Hill, 2009.
- Kim, I.Y., Lee, D.W., Lee, S.B. and Kwak, I.S. The role of uric acid in kidney fibrosis: experimental evidences for the causal relationship. *BioMed Res Int* 2014; 638732: 1-9.
- Klopfleisch, R. Multiparametric and semiquantitative scoring systems for the evaluation of mouse model histopathology - a systematic review. *BMC Vet Res* 2013; 9:123.
- Kurniari, P.K., Kambayana, G., dan Putra, T.R. Hubungan hiperurisemia dan fraction uric acid clearance di desa tenganan pegringsingan karangasem bali. *J Peny Dalam* 2011; 12(2): 77-80.
- Leary, S., Underwood, W., Anthony, R., Cartner, S., Corey, D., Grandin, T., et al. AVMA Guidelines for the Euthanasia of Animals: 2013 Edition. Available at: https://www.avma.org/issues/animal_welfare/euthanasia.pdf. Accessed 11 November 2015.
- Lech, M., Grobmayr, R., Ryu, M., Lorenz, G., Hartter, I., Mulay, S.R., et al. Macrophage phenotype controls long-term aki outcomes—kidney regeneration versus atrophy. *J Am Soc Nephrol* 2014; 25: 292–304.
- Lee, S., Huen, S., Nishio, H., Nishio, S., Lee, H.K., Choi, B.S., et al. Distinct macrophage phenotypes contribute to kidney injury and repair. *J Am Soc Nephrol* 2011; 22: 317-326.
- Lin, S.L. and Duffield, J.S. Macrophages in kidney injury and repair. *Acta Nephrologica* 2012; 26(2): 45-57.
- Liu, B., Wang, T., Zhao, H.N., Yue, W.W., Yu, H.P., Liu, C.X., et al. The prevalence of hyperuricemia in China: a meta-analysis. *BMC Public Health* 2011; 11:832.
- Lu, J., Cao, Q., Zheng, D., Sun, Y., Wang, C., Yu, X., et al. Discrete functions of M2a and M2c macrophage subsets determine their relative efficacy in treating chronic kidney disease. *Kidney Int* 2013; 84: 745–755.
- Martinez, F.O. and Gordon, S. The M1 and M2 paradigm of macrophage activation: time for reassessment. *F1000 Prime Rep* 2014; 6:13.
- Mazzali, M., Hughes, J., Kim, Y.G., Jefferson, J.A., Kang, D.H., Gordon, K.L., et al. Elevated uric acid increases blood pressure in the rat by a novel crystal independent mechanism. *Hypertension* 2001; 38(5): 1101–1106.
- Mazzali, M., Kanellis, J., Han, L., Feng, L., Xia, Y.Y., Chen, Q., et al. Hyperuricemia induces a primary renal arteriopathy in rats by a blood pressure-independent mechanism. *Am J Physiol Renal Physiol* 2002; 282(6): F991–F997.
- Meng, X.M., Nikolic-Paterson, D.J. and Lan, H.Y. Inflammatory processes in renal fibrosis. *Nat Rev Nephrol* 2014; 10(9): 493-503.
- Mills, C.D. Anatomy of a discovery: M1 and M2 macrophages. *Front Immunol* 2015; 6: 212.
- Murray, P.J. and Wynn, T.A. Protective and pathogenic functions of macrophage subsets. *Nat Rev Immunol* 2011; 11: 723-737.
- Nomura, J., Busso, N., Ives, A., Tsujimoto, S., Tamura, M., So, A., et al. Febuxostat, an inhibitor of xanthine oxidase, suppresses lipopolysaccharide-

- induced MCP-1 production via MAPK phosphatase-1-mediated inactivation of JNK. *PLoS ONE* 2013; 8(9): e75527.
- Pan, B., Liu, G., Jiang, Z. and Zheng, D. Regulation of renal fibrosis by macrophage polarization. *Cell Physiol Biochem* 2015; 35: 1062-1069.
- Patcher, P., Nivorozhkin, A. and Szabo, C. Therapeutic effects of xanthine oxidase inhibitors: renaissance half a century after the discovery of allopurinol. *Pharmacol Rev* 2006; 58: 87–114.
- Peglow, S., Toledo, A.H., Anaya-Prado, R., Lopez-Neblina, F. and Toledo-Pereya, L.H. Allopurinol and xanthine oxidase inhibition in liver ischemia reperfusion. *J Hepatobiliary Pancreat Sci* 2011 Mar;18(2):137-46.
- Robert, C., Lu, X., Law, A., Freeman, T.C. and Hume, D.A. Macrophages.com: an on-line community resource for innate immunity research. *Immunobiology* 2011; 216(11): 1203-11.
- Ryu, E.S., Kim, M.J., Shin, H.S., Jang, Y.H., Choi, H.S., Jo, I., et al. Uric acid-induced phenotypic transition of renal tubular cells as a novel mechanism of chronic kidney disease. *Am J Physiol Renal Physiol* 2013; 304(5): F471–F480.
- Sánchez-Lozada, L. G., Tapia, E., Santamaria, J., Avila-Casado, C., Soto, V., Nepomuceno, T., et al. Mild hyperuricemia induces vasoconstriction and maintains glomerular hypertension in normal and remnant kidney rats. *Kidney Int* 2005; 67(1): 237–247.
- Sánchez-Lozada, L.G., Lanaspá, M.A., Cristóbal-García, M., García-Arroyo, F., Soto, V., Cruz-Robles, D., et al. Uric acid-induced endothelial dysfunction is associated with mitochondrial alterations and decreased intracellular ATP concentrations. *Nephron Exp Nephrol* 2012; 121(3-4): e71–e78.
- Sánchez-Lozada, L.G., Soto, V., Tapia, E., Avila-Casado, C., Sautin, Y.Y., Nakagawa, T., et al. Role of oxidative stress in the renal abnormalities induced by experimental hyperuricemia. *Am J Physiol Renal Physiol* 2008; 295(4): F1134–F1141.
- Sánchez-Lozada, L.G., Tapia, E., Avila-Casado, C., Soto, V., Franco, M., Santamaria, J., et al. Mild hyperuricemia induces glomerular hypertension in normal rats. *Am J Physiol Renal Physiol* 2002; 283(5): F1105–F1110.
- Sánchez-Lozada, L.G., Tapia, E., López-Molina, R., Nepomuceno, T., Soto, V., Avila-Casado, C., et al. Effects of acute and chronic L-arginine treatment in experimental hyperuricemia. *Am J Physiol Renal Physiol* 2007; 292(4): 1238–1244.
- Sautin, Y.Y. and Johnson, R.J. Uric acid: the oxidant-antioxidant paradox. *Nucleosides Nucleotides Nucleic Acids* 2008; 27 (6-7): 608–619.
- So, A. and Thorens, B. Uric acid transport and disease. *J Clin Invest* 2010; 120(6): 1791–1799.
- Stefater, J.A., Ren, S., Lang, R.A. and Duffield, J.S. Metchnikoff's policemen: macrophages in development, homeostasis and regeneration. *Trends Mol Med* 2011; 17: 743-752.
- Vinuesa, E., Hotter, G., Jung, M., Herrero-Fresneda, I., Torras, J., Sola, A. Macrophage involvement in the kidney repair phase after ischaemia/reperfusion injury. *J Pathol* 2008; 214: 104–113.

- Watanabe, S., Kang, D.H., Feng, L., Nakagawa, T., Kanellis, J., Lan, H., et al. Uric acid, hominoid evolution, and the pathogenesis of salt-sensitivity. *Hypertension* 2002; 40(3): 355–360.
- Wu, X., Wakamiya, M., Vaishnav, S., Geske, R., Montgomery, C., Jones, P., et al. Hyperuricemia and urate nephropathy in urate oxidase-deficient mice. *Proc Natl Acad Sci U S A* 1994; 91(2): 742–746.
- Zhang, M., He, Y., Sung, X., Li, Q., Wang, W., Zhao, A., and Di, W. A high M1/M2 ratio of tumor-associated macrophages is associated with extended survival in ovarian cancer patients. *J Ovarian Res* 2014; 7:19.
- Zhou, Y., Fang, L., Jiang, L., Wen, P., Cao, H., He, W., et al. Uric acid induces renal inflammation via activating tubular NF-kB signaling pathway. *PLoS ONE* 2012; 7(6): e39738.
- Zhu, Y., Pandya, B.J. and Choi, H.K. Prevalence of gout and hyperuricemia in the us general population. *Arthritis Rheum* 2011; 63(10): 3136-3141.