

DAFTAR PUSTAKA

- Abegunde, O. K., Mu, T. -H., Chen, J. -W., and Deng, F. -M. (2013). Physicochemical characterization of sweet potato starches popularly used in Chinese starch industry. *Food Hydrocolloids*, 33(2), 169–177.
- Anderson, J. W., Baird, P., Davis Jr, R. H., Ferreri, S., Knudtson, M., Koraym, A. Waters, V., and Williams, C. L. (2009). Health benefits of dietary fiber: a Lead Article. *Nutrition Review*, 67(4), 188–205.
- Anonim. (2001). AACC report: The definition of dietary fiber. *Cereal Foods World*, 46(3), 112-126.
- Anonim. (2007a). Dietary Fibre: An Information Statement. *Institute of Food Science and Technology, London* (April), 1–10.
- Anonim. (2007b). Information Statement: Dietary Fibre. *Institute of Food Science and Technology*. 1–10.
- Anonim. (2008). Prebiotics and probiotics. *World Gastroenterology Organisation Practice Guide*.
- Asp, N.G., Johansson, C.G., Hallmer, H., and Siljestrom, M. (1983). Rapid Enzymatic Assay of Insoluble and Soluble Dietary Fiber. *Journal Agricultural Food Chemistry* 31, 476-482.
- Avanza, M. V., Chaves, M. G., Acevedo, B. A., and Añón, M. C. (2012). Functional properties and microstructure of cowpea cultivated in north-east Argentina. *LWT - Food Science and Technology*, 49(1), 123–130.
- Azorín-Ortuño, M., Urbán, C., Cerón, J. J., Tecles, F., Allende, A., Tomás-Barberán, F. A., and Espín, J. C. (2009). Effect of low inulin doses with different polymerisation degree on lipid metabolism, mineral absorption, and intestinal microbiota in rats with fat-supplemented diet. *Food Chemistry*, 113(4), 1058–1065.
- Bedani, R., Rossi, E. A., and Saad, S. M. I. (2013). Impact of inulin and okara on *Lactobacillus acidophilus* La-5 and *Bifidobacterium animalis* Bb-12 viability in a fermented soy product and probiotic survival under in vitro simulated gastrointestinal conditions. *Food Microbiology*, 34(2), 382–389.
- BeMiller, J., and Whistler, R. (2009). Starch Chemistry and Technology (3rd ed.). *Academic Press*.
- Bertolini, A. C. (2010). Starches: characterization, properties, and application. *CRC Press*.
- Bird, A. R., Vuaran, M., Crittenden, R., Hayakawa, T., Playne, M. J., Brown, I. JL., and Topping, D. L. (2009). Comparative effects of high-amylose starch and a fructooligosaccharide on fecal bifidobacteria numbers and short-chain fatty acids in pig fed *Bifidobacterium animalis*. *Dig Dis Sci*, 54, 947-954.

- Bosscher, D., Van Loo, J., and Franck, A. (2006). Inulin and oligofructose as functional ingredients to improve bone mineralization. *International Dairy Journal*, 16(9), 1092–1097.
- Brennan, C. S., Kuri, V., and Tudorica, C. M. (2004). Inulin-enriched pasta: effects on textural properties and starch degradation. *Food Chemistry*, 86(2), 189–193.
- Brumovsky, J. O., and Thompson, D. B. (2001). Production of boiling-stable granular resistant starch by partial acid hydrolysis and hydrothermal treatments of high-amylose maize starch. *Cereal Chemistry*, 78(6), 680–689.
- Campos, F. M., Figueiredo, A. R., Hogg, T. A., and Couto, J. A. (2009). Effect of phenolic acids on glucose and organic acid metabolism by lactic acid bacteria from wine. *Food Microbiology*, 26, 409–414.
- Cara, C., Ruiz, E., Carvalheiro, F., Moura, P., Ballesteros, I., Castro, E., and Gírio, F. (2012). Production, purification and characterisation of oligosaccharides from olive tree pruning autohydrolysis. *Industrial Crops and Products*, 40, 225–231.
- Carvalheiro, F., Esteves, M. P., Parajó, J. C., Pereira, H., and Gírio, F. M. (2004). Production of oligosaccharides by autohydrolysis of brewery's spent grain. *Bioresource Technology*, 91, 93–100.
- Carvalho, A. F. A., Neto, P. d. O., da Silva, D. F., and Pastore, G. M. (2013). Xylo-oligosaccharides from lignocellulosic materials: Chemical structure, health benefits and production by chemical and enzymatic hydrolysis. *Food Research International*, 51(1), 75–85.
- Charalampopoulos, D., and Rastall, R. A. (2012). Prebiotics in foods. *Current Opinion in Biotechnology*, 23(2), 187–91.
- Chung, H. -J., Liu, Q., and Hoover, R. (2009). Impact of annealing and heat-moisture treatment on rapidly digestible, slowly digestible and resistant starch levels in native and gelatinized corn, pea and lentil starches. *Carbohydrate Polymers*, 75(3), 436–447.
- Colussi, R., Pinto, V. Z., Halal, S. L. M E., Vanier, N. L., Villanova, F. A., e Silva, R. M., Zavareze, E. d. R., and Dias, A. R. G. (2014). Structural, morphological, and physicochemical properties of acetylated high-, medium-, and low-amylose rice starches. *Carbohydrate Polymers*, 103, 405–413.
- Danjo, K., Nakaji, S., Fukuda, A., Shimoyama, T., Sakamoto, J., and Sugawara, K. (2003). The resistant starch level of heat moisture-treated high amylose cornstarch is much lower when measured in the human terminal ileum than when estimated *in vitro*. *The Journal of Nutrition*, 133(7), 2218–2221.
- Datar, R., Huang, J., Maness, P., Mohagheghi, A., Czernik, S., and Chornet, E. (2007). Hydrogen production from the fermentation of corn stover biomass pretreated with a steam-explosion process. *International Journal of Hydrogen Energy*, 32(8), 932–939.

- Davis, L. M. G., Martínez, I., Walter, J., and Hutkins, R. (2010). A dose dependent impact of prebiotic galactooligosaccharides on the intestinal microbiota of healthy adults. *International Journal of Food Microbiology*, 144(2), 285–292.
- Davoodabadi, A., Dallal, M. M. S., Foroushani, A. R., Douraghi, M., Yazdi, M. K. S. and Harati, F. A. (2015). Antibacterial activity of *Lactobacillus* spp. isolated from the feces of healthy infants against enterohatogenic bacteria. *Anaerobe*, 34, 53-58.
- Deepa, B., Abraham, E., Cherian, B. M., Bismarck, A., Blaker, J. J., Pothan, L. A., Leao, A. L., de Souza, S. F., and Kottaisamy, M. (2011). Bioresource Technology Structure , morphology and thermal characteristics of banana nano fibers obtained by steam explosion. *Bioresource Technology*, 102(2), 1988-1997.
- Dobre, T., Stroescu, M., Stoica, A., Draghici, E., and Antohe, N. (2008). Inulin Extraction and Encapsulation. *Chemical Bulletin "POLITEHNICA" Univ. (Timișoara)*, 53(67), 215–217.
- Elleuch, M., Bedigian, D., Roiseux, O., Besbes, S., Blecker, C., and Attia, H. (2011). Dietary fibre and fibre-rich by-products of food processing: Characterisation, technological functionality and commercial applications: A review. *Food Chemistry*, 124(2), 411–421.
- Favier, C. F., de Vos, W. M., and Akkermans, A. D. L. (2003). Development of bacterial and bifidobacterial communities in feces of newborn babies. *Anaerobe*, 9(5), 219–29.
- Franck, A., and Leenher, L. D. (2003). Inulin. *Biopolymers Vol. 6*, 439-473.
- Fu, Z-q., Wang, L-j., Li, D., and Adhikari, B. (2012). Effects of partial gelatinization on structure and thermal properties of corn starch after spray drying. *Carbohydrate Polymers*, 88, 1319-1325.
- Fuentes-Zaragoza, E., Riquelme-Navarrete, M. J., Sánchez-Zapata, E., and Pérez-Álvarez, J. A. (2010). Resistant starch as functional ingredient: A review. *Food Research International*, 43(4), 931–942.
- Futch, J. (2009). Altering Sweet Potato Starch Functionality by Amino Acids and pH Treatments: A Thesis. Louisiana State University and Agricultural and Mechanical College.
- Gibson G. R., and Roberfroid, M. B. (1995). Dietary modulation of the human colonie microbiota: Introducing the concept of prebiotics *J. Nutr.* 125, 1401-1412.
- Gibson, G. R. (2004). Fibre and effects on probiotics (the prebiotic concept). *Clinical Nutrition Supplements*, 1(2), 25–31.
- Gibson, G. R., Scott, K. P., Rastall, R. A., Tuohy, K. M., Hotchkiss, A., Dubert-Ferrandon, A., Gareau, M., Murphy, E. F., Saulnier, D., Loh, G., Macfarlane, S., Delzenne, N., Ringel, Y., Kozianowski, G., Dickmann, R., Lenoir-Wijnkoop, I., Walker, C., and Buddington, R. (2010). Dietary

- prebiotics: current status and new definition. *Food Science and Technology Bulletin: Functional Foods*, 7(1), 1-19.
- Gordillo, C. A. S., Valencia, G. A., Zapata, R. A. V, and Henao, A. C. A. (2014). Physicochemical Characterization of Arrowroot Starch (*Maranta arundinacea* Linn) and Glycerol/Arrowroot Starch Membranes. *International Journal of Food Engineering*, 10(4), 727–735.
- Grizard, D., and Barthomeuf, C. (1999). Non-digestible oligosaccharides used as prebiotic agents: mode of production and beneficial effects on animal and human health. *Reproduction, Nutrition, Development*, 39, 563–588.
- Grootaert, C., Delcour, J. A., Courtin, C. M., Broekaert, W. F., Verstraete, W., and de Wiele, T. V. (2007). Microbial metabolism and prebiotic potency of arabinoxylan oligosaccharides in the human intestine. *Trends in Food Science and Technology*, 18, 64–71.
- Gunaratne, A. And Hoover, R. (2002). Effect of heat-moisture treatment on the sturcture and physicochemical properties ot tuber and root starches. *Carbohydrate polymers*, 49, 425-437.
- Guo, J., Liu, L., Lian, X., Li, L., and Wu, H. (2014). The properties of different cultivars of Jinhai sweet potato starches in China. *International Journal of Biological Macromolecules*, 67, 1–6.
- Harmayani, E., Kumalasari, I. D., and Marsono, Y. (2011). Effect of arrowroot (*Maranta arundinacea* L.) diet on the selected bacterial population and chemical properties of caecal digesta of Sprague Dawley rats, *International Research Journal of Microbiology*, 2(8), 278–284.
- Huebner, J., Wehling, R. L., and Hutkins, R. W. (2007). Functional activity of commercial prebiotics. *International Dairy Journal*, 17(7), 770–775.
- Jacquet, N., Quiévy, N., Vanderghem, C., Janas, S., Blecker, C., Wathélet, B., Devaux, J., and Paquot, M. (2011). Influence of steam explosion on the thermal stability of cellulose fibres. *Polymer Degradation and Stability*, 96, 1582–1588.
- Jiang, Q., Gao, W., Shi, Y., Li, X., Wang, H., Huang, L., and Xiao, P. (2013). Physicochemical properties and *in vitro* digestion of starches from different *Dioscorea* plants. *Food Hydrocolloids*, 32, 432-439.
- Jyothi, A. N., Sheriff, J. T., and Sajiev, M. S., Physical and functional properties of arrowroot starch extrudates. *Journal of Food Science*, 74(2), 97-104.
- Kelly, G. (2008). Inulin-Type Prebiotics : A Review (Part 1). *Alternative Medicine Review*, 13(4), 315–329.
- Kondepudi, K. K., Ambalam, P., Nilsson, I., Wadström, T., and Ljungh, Å. (2012). Prebiotic-non-digestible oligosaccharides preference of probiotic bifidobacteria and antimicrobial activity against *Clostridium difficile*. *Anaerobe*, 18(5), 489–497.
- Kumar, S. B., and Prabhasankar, P. (2014). Low glycemic index ingredients and modified starches in wheat based food processing : A review. *Trends in*

Food Science and Technology, 35(1), 32–41.

- Kusharto, C. M., Surono, I. S., and Adi, A. C. (2009). Efek sinbiotik prebiotik asal pangan lokal dengan *Enterococcus faecium* IS27526 terhadap BAL dan berat badan pada tikus percobaan. *Prosiding Seminar Hasil-Hasil Penelitian IPB*.
- Lee, C. J., Kim, Y., Choi, S. J., and Moon, T. W. (2012). Slowly digestible starch from heat-moisture treated waxy potato starch: Preparation, structural characteristics, and glucose response in mice. *Food Chemistry*, 133(4), 1222–1229.
- Leite, J. T. C., Martinelli, P., Murr, F. E. X., and Park, K. J. (2004). Study of The Inulin Concentration by Physical Methods. *Proceeding 14th International Drying Symposium* (pp. 868–875). Sao Paulo-Brazil.
- Leszczyński, W. (2004). Resistant starch – classification, structure, production. *Polish Journal Food Nutrition Science*, 13(54), 37–50.
- Li, S. L., and Gao, Q. Y. (2010). Effect of Heat-Moisture Treatment on the Formation and Properties of Resistant Starches From Mung Bean (*Phaseolus radiatus*). *World Academy of Science, Engineering and Technology*, 4(12), 703–710.
- Li, W., Wang, K., Sun, Y., Ye, H., Hu, B., and Zeng, X. (2015). Influence of structure of galactooligosaccharides and fructooligosaccharide on the fermentation *in vitro* by human intestinal microbiota. *Functional Foods*, 13, 158–168.
- López-Molina, D., Navarro-Martínez, M. D., Melgarejo, F. R., Hiner, A. N. P., Chazarra, S., and Rodríguez-López, J. N. (2005). Molecular properties and prebiotic effect of inulin obtained from artichoke (*Cynara scolymus* L.). *Phytochemistry*, 66(12), 1476–84.
- Ma, Z., Boye, J. I., Simpson, B., K., Prasher, S. O., Monpetit, D., and Malcolmson, L., Thermal processing effects on the functional properties and microstructure of lentil, chickpea, and pea flours. *Food Research International*, 44, 2534–2544
- Manning, T. S., and Gibson, G. R. (2004). Prebiotics. *Best Practice and Research Clinical Gastroenterology*, 18(2), 287–298.
- Marsono, Y. (2002a). Indeks glikemik umbi-umbian (Glycemic Index of tuber foods). *Agritech*, 22(1).
- Marsono, Y. (2002b). Pengaruh pengolahan terhadap pati resisten pisang kepok (*Musa paradisiaca* fa. *typica*) dan pisang tanduk (*Musa paradisiaca* fa. *Corniculata*). *Agritech*, 22(2).
- Martínez, M., Gullón, B., Yáñez, R., Alonso, J. L., and Parajó, J. C. (2010). Kinetic assessment on the autohydrolysis of pectin-rich by-products. *Chemical Engineering Journal*, 162, 480–486.
- Meyer, D., Bayarri, S., Tárrega, a., and Costell, E. (2011). Inulin as texture modifier in dairy products. *Food Hydrocolloids*, 25(8), 1881–1890.

- Montane, D., Farriol, X., Salvado, J., Jollez, P., and Chornet, E. (1998). Application of steam explosion to the fractionation and rapid vapor-phase alkaline pulping of wheat straw. *Biomassa and Bioenergy*, 14(3), 261-276.
- Moore, K. E. (2011). Bioogical analysis of prebiotics in various processed food matrices: A Thesis. *University of Nebraska*.
- Moro, G., Minoli, I., Mosca, M., Fanaro, S., Jelinek, J., Stahl, B., and Boehm, G. (2002). Dosage-related bifidogenic effects of galacto- and fructooligosaccharides in formula-fed term infants. *Journal of Pediatric Gastroenterology and Nutrition*, 34(March), 291–295.
- Mosfegh, A. L., Friday, J. E., Goldman, J. P., and Ahuja, J. K. C. (1999). Presence of inulin and oligofructose in the diets of Americans. *Journal of Nutritions*, 129, 1407S-1411S.
- Murphy, M. M., Douglass, J. S., and Birkett, A. (2008). Resistant starch intakes in the United States. *Journal of the American Dietetic Association*, 108, 67–78.
- Nabarlatz, D., Ebringerová, A., and Montané, D. (2007). Autohydrolysis of agricultural by-products for the production of xylo-oligosaccharides. *Carbohydrate Polymers*, 69, 20–28.
- Noda, Y., Asada, C., Sasaki, C., Hashimoto, S., and Nakamura, Y. (2013). Extraction method for increasing antioxidant activity of raw garlic using steam explosion. *Biochemical Engineering Journal*, 73, 1–4.
- Nurliyani, Julia, M., and Harmayani, E. (2013). Effect of arrowroot (*Maranta arundinacea*) cookies intervention on fecal secretory immunoglobulin A and physical properties of children under five years. *Interntional Research Journal of Microbiology*, 4(1), 21-28.
- Ojetti, V., Gigante, G., Ainora, M. E., Fiore, F., Barbaro, F., and Gasbarrini, A. (2009). Microflora imbalance and gastrointestinal diseases. *Digestive and Liver Disease*, 3, 35-39.
- Parada, J., and Aguilera, J. M. (2012). Effect of native crystalline structure of isolated potato starch on gelatinization behavior and consequently on glycemic response. *Food Research International*, 45(1), 238–243.
- Phelps, C. F. (1965). The physical properties of inulin solutions. *The Biochemical Journal*, 95, 41–47.
- Pompei, A., Cordisco, L., Raimondi, S., Amaretti, A., Pagnoni, U. M., Matteuzzi, D., and Rossi, M. (2008). *In vitro* comparison of the prebiotic effects of two inulin-type fructans. *Anaerobe*, 14(5), 280–6.
- Pramodao, K. S., and Riar, C. S. (2014). Comparative study of effect of modification with onic gums and dry heating on the physicochemical characteristic of sweet potato and taro starches. *Food Hydrocolloids*, 35, 613-619
- Puncha-arnon, S., and Uttapap, D. (2013). Rice starch vs. rice flour: Differences in their properties when modified by heat-moisture treatment.

Carbohydrate Polymers, 91(1), 85–91.

- Rachmadani, A. D. dan Estiasih, T. (2015). Beras analog berbasis umbi garut (*Maranta arundinacea*) dan alginat sebagai pangan berkhasiat obat (*Medicinal Foods*) yang diujikan pada tikus hiperglikemik. <http://jpa.ub.ac.id/index.php/jpa/article/download/357/368> diakses Desember 2015
- Rehinan, Z., Rashid, M., and Shah, W.H. (2004). Insoluble dietary fibre components of food legumes are affected by soaking and cooking processes. *Food Chemistry*, 85, 245-249.
- Roberfroid, M. (2002). Functional food concept and its application to prebiotics. *Digestive and Liver Disease Supplements*, 34(2), 105–110.
- Rodríguez-Cabezas, M. E., Camuesco, D., Arribas, B., Garrido-Mesa, N., Comalada, M., Bailón, E., Cuerto-Sola, M., Utrilla, P., Guerra-Hernández, E., Pérez-Roca, C., Gálvez, J., and Zarzuelo, A. (2010). The combination of fructooligosaccharides and resistant starch shows prebiotic additive effects in rats. *Clinical Nutrition*, 29(6), 832–839.
- Ronkart, S. N., Blecker, C. S., Fourmanoir, H., Fougnes, C., Deroanne, C., Van Herck, J. C., and Paquot, M. (2007). Isolation and identification of inulooligosaccharides resulting from inulin hydrolysis. *Analytica Chimica Acta*, 604, 81–87.
- Russell, P. L., Berry, C. S., and Greenwell, P. (1989). Characterisation of resistant starch from wheat and maize. *Journal of Cereal Science*, 9(1), 1–15.
- Saad, N., Delattre, C., Urdaci, M., Schmitter, J. M., and Bressollier, P. (2013). An overview of the last advances in probiotic and prebiotic field. *LWT - Food Science and Technology*, 50(1), 1–16.
- Sajilata, M. G., Singhal, R. S., and Kulkarni, P. R. (2006). Resistant starch— A Review,. *Comprehensive reviews in food science and food safety*, 5, 1-17.
- Sanz, T., Martínez-Cervera, S., Salvador, A., and Fiszman, S. M. (2010). Resistant starch content and glucose release of different resistant starch commercial ingredients: effect of cooking conditions. *European Food Research and Technology*, 231(5), 655–662.
- Scott, K. P., Gratz, S. W., Sheridan, P. O., Flint, H. J., and Duncan, S. H. (2013). The influence of diet on the gut microbiota. *Pharmacological Research*, 69(1), 52–60.
- Sims, I., Ryan, J. L. J., and Kim, S. H. (2014). *In vitro* fermentation of prebiotic oligosaccharides by *Bifidobacterium Lctis* HNO19 and *Lactobacillus* spp. *Anaerobe*, 25, 11-17
- Slavin, J. L. (2005). Dietary fiber and body weight. *Nutrition*, 21, 411–418.
- Slavin, J. (2013). Fiber and prebiotics: mechanism and health benefits. *Nutrients*, 5, 1417-1435.
- Ślizewska, K., Kapuśniak, J., Barczyńska, R., and Jochym, K. (2012). Resistant

dextrins as prebiotic in Comprehensive Studies on Glycobiology and Glycotechnology. 261–288.

- Teng, C., Yan, Q., Jiang, Z., Fan, G., and Shi, B. (2010). Production of xylooligosaccharides from the steam explosion liquor of corncobs coupled with enzymatic hydrolysis using a thermostable xylanase. *Bioresource Technology*, 101(19), 7679–7682.
- Tuohy, K. M., Rouzaud, G. C. M., Brück, W. M., and Gibson, G. R. (2005). Modulation of the human gut microflora towards improved health using prebiotics--assessment of efficacy. *Current Pharmaceutical Design*, 11, 75–90.
- Villegas, B., Tárrega, A., Carbonell, I., and Costell, E. (2010a). Optimising acceptability of new prebiotic low-fat milk beverages. *Food Quality and Preference*, 21(2), 234–242.
- Wichienchot, S., Thammarutwasik, P., Jongjareonrak, A., Chansuwan, W., Preeya Hmadhlu, Hongpattarakere, T., Itharat, A., and Ooraikul, B. (2011). Extraction and analysis of prebiotics from selected plants from southern Thailand. *Songklanakarin Journal of Science and Technology*, 33(5), 517–523.
- Xiao, X., Bian, J., Peng, X. P., Xu, H., Xiao, B., and Sun, R. C. (2013). Bioresource Technology Autohydrolysis of bamboo (*Dendrocalamus giganteus* Munro) culm for the production of xylo-oligosaccharides. *Bioresource Technology*, 138, 63–70.
- Xiaoli, X., Liyi, Y., Shuang, H., Wei, L., Yi, S., Hao, M., Jusong, Z., and Xiaoxiong, Z. (2008). Determination of oligosaccharide contents in 19 cultivars of chickpea (*Cicer arietinum* L) seeds by high performance liquid chromatography. *Food Chemistry*, 111(1), 215–219.
- Yadav, B. S., Guleria, P., and Yadav, R. B. (2013). Hydrothermal modification of indian water chesnut starch: Influence of heat-moisture treatment and annealing on the physicochemical, gelatinization and pasting characteristic. *LWT-Food Science and Technology*, 53: 211–217.
- Yang, L., Zhou, Y., Wu, Y., Meng, X., Jiang, Y., Zhang, H., and Wang, H. (2016). Preparation and physicochemical properties of three types of modified glutinous rice starches. *Carbohydrate Polymers*, 137, 305–313.
- Yang, Z., Hu, J., and Zhao, M. (2011). Isolation and quantitative determination of inulin-type oligosaccharides in roots of *Morinda officinalis*. *Carbohydrate Polymers*, 83(4), 1997–2004.
- Zhang, Y., Zhao, W., Yang, R., Ahmed, M. A., Hua, X., Zhang, W., and Zhang, Y. (2013). Preparation and functional properties of protein from heat-denatured soybean meal assisted by steam flash-explosion with dilute acid soaking. *Journal of Food Engineering*, 119(1), 56–64.
- Zhang, Z., Jiang, B., Liao, X., Yi, J., Hu, X., and Zhang, Y. (2012). Inactivation of *Bacillus subtilis* spores by combining high-pressure thermal sterilization and ethanol. *International Journal of Food Microbiology*, 160(2), 99–104.



Zobel, H. F., Young, S. N., and Rocca, L. A. (1988). Starch gelatinization: An X-ray diffraction study. *Cereal Chemistry*, 65(6), 443–446.

Zuleta, A., and Sambucetti, M. E. (2001). Inulin determination for food labeling. *J. Agric. Food. Chem*, 49, 4570-4572.