

DAFTAR PUSTAKA

- Aasheim, E.T.M., Hofsø, D., Hjelmessaeth, J., Sandbu, R., 2008. Peripheral neuropathy and severe malnutrition following duodenal switch. *Obes Surg.*, 18(12):1640-1643.
- Adler, A.I., Boyko, E.J., Ahroni, J.H., Stensel, V., Forsberg, R.C., 1997. Risk factors for diabetic peripheral sensory neuropathy. Results of the Seattle Prospective Diabetic Foot Study. *Diabetes Care*, 20: 1162-1167.
- Akbaş, E., Dericci, E., Söylemez, F., Kanik, A., Polat, F., 2004. An investigation of effects of toluene and cigarette smoking on some blood parameters and lymphocyte life span. *Cell Biol Toxicol.*, 20(1):33-40.
- Akisu, M., 1996. Severe acute thinner intoxication. *Turk Y Pediatr*; 38(2):223-5
- Ameno, K., Kiriu, T., Fuke, C., Ameno, S., Shinohara, T., Ijiri, I., 1992. Regional brain distribution of toluene in rats and in a human autopsy. *Archives of toxicology*, 66: 153–156.
- American Chemistry Council, 2006. *Voluntary Children's Chemical Evaluation Program (VCCEP) Tier 1 Pilot submission Toluene (CAS No. 108-88-3)*. Washington, DC: American Chemistry Council.
- American Conference of Governmental Industrial Hygienists, 2000. *Threshold limit values for chemical substances and physical agents and biological exposure indices 2000*. Cincinnati, OH: American Conference of Governmental Industrial Hygienists.
- Angerer, J., 1985. Occupational chronic exposure to organic solvents, XII: o-cresol excretion after toluene exposure. *Iny Arch Occup Environ Health*; 56:323-328.
- Appenzeller, O., Parks, R.D., McGee, J., 1968. Peripheral neuropathy in chronic disease of the respiratory tract. *Am J Med.*, 11: 873-880
- Azhary, H., Farooq, M.U., Bhanushali, M., 2010. Peripheral neuropathy: differential diagnosis and management. *American Family Physician*; 81(7).
- Badan Standardisasi Nasional, 2005. *SNI 19-0232-2005: Nilai ambang batas (NAB) zat kimia di udara tempat kerja*. Jakarta: Badan Standardisasi Nasional.
- Badanich KA, Adler KJ, Kirstein CL., 2006. Adolescents differ from adults in cocaine conditioned place preference and cocaine-induced dopamine in the nucleus accumbens septi. *Eur J Pharmacol.*, 550:95–106.
- Balai Lingkungan Yogyakarta Propinsi DI Yogyakarta, 2011. *Kualitas udara ambien kota Yogyakarta tahun 2011*. Diunduh pada tanggal 3 November 2014 dari laman <http://blh.jogjaprovo.go.id/2012/04/kualitas-udara-ambien-kota-yogyakartatahun-2011/>
- Bauer, B., 1993. *Miljø og arbejdsmiljø i den grafiske branche*. Bind 1. Ydre miljø. Miljøprojekt No. 212. Miljøstyrelsen.
- Bennett, M.I., Attal, N., Backonja, M.M., Baron, R., Bouhassira, D., Freynhagen, R., Scholz, J., To'ille, T.R., Wittchen, H.-U., Jensen, T.S., 2007. Using screening tools to identify neuropathic pain. *Pain*; 127:199-203.

- Berger, A.R., Schaumburg, H.H., 2005. *Human toxic neuropathy caused by industrial agents*. Dalam: Dyck, P.J., Thomas, P.K., (Editors). *Peripheral neuropathy (4th Ed)*. Philadelphia, PA: Elsevier Inc.
- Bowen, S.E., Charlesworth, J.D., Tokarz, M.E., Wright, M.J. Jr., Wiley, J.L., 2007. Decreased sensitivity in adolescent vs adult rats to the locomotor activating effects of toluene. *Neurotoxicol Teratol*, 29(6):599-606.
- Brugnone F, Gobbi M, Ayyad K, Giuliari C, Cerpelloni M, Perbellini L., 1995. Blood toluene as a biological index of environmental toluene exposure in the "normal" population and in occupationally exposed workers immediately after exposure and 16 hours later. *Int Arch Occup Environ Health*, 66(6):421-5.
- Cacchione, P.Z., 2010. Sensory Changes. *Consult GeriRN.org*
- Callaghan, B., Feldman, E., 2013. The metabolic syndrome and neuropathy: therapeutic challenges and opportunities. *Ann Neurol.*, 74: 397-403.
- Canadian Center for Occupational Health and Safety, 2008. *Chemical profiles toluene*. Diunduh pada tanggal 18 Juli 2014 dari laman http://www.ccohs.ca/oshanswers/chemicals/chem_profiles/toluene
- Carlsson, A., 1982. Exposure to toluene: Uptake, distribution and elimination in man. *Scandinavian journal of work & environmental health*, 8: 43-55.
- Carstensen, M., 1997. *Biochemicals for the printing industry*. Minneapolis, MN: The Institute for Local Self-Reliance.
- Caselli, M., de Gennaro, G., Marzocca, A., Trizio, L., Tutino, M., 2010. Assessment of the impact of the vehicular traffic on BTEX concentration in ring roads in urban areas of Bari (Italy). *Chemosphere*, 81(3):306-311.
- Cheng, Y.J., Gregg, E.W., Kahn, H.S., Willians DE, Rekeneire, N.D., Narayan, K.M.V., 2006. Peripheral insensate neuropathy- a tall problem in US adults? *American Journal of Epidemiology*, 8:873-80
- Collins SL, Izenwasser S., 2002. Cocaine differentially alters behavior and neurochemistry in periadolescent versus adult rats. *Brain Res Dev Brain Res.*, 138:27-34. [PubMed]
- Chouanière, D., Pascal Wild, P., Fontana, J.-M., Héry, M., Fournier, M., Baudin, V., Subra, I., Rousselle, D., Toamain, J.-P., Saurin, S., Ardiot, M.-R., 2002. Neurobehavioral disturbances arising from occupational toluene exposure. *American Journal of Industrial Medicine*; 41(2):77-88.
- de Joode, B.W., Tielemans, E., Vermeulen, R., Wegh, H., Kromhout, H., 2005. Dermal exposure assessment to benzene and toluene using charcoal cloth pads. *Journal of Exposure Analysis and Environmental Epidemiology*; 15:47-50.
- Dick, F., Sample, S., Chen, R., Seaton, A., 2000. Neurological deficits in solvent-exposed painters: a syndrome including impaired colour vision, cognitive defects, tremor and loss of vibration sensation. *Q J Med.*, 93:655-661.
- Draper, T.H.J., Bamiou, D-E., 2009. Auditory neuropathy in a patient exposed to xylene: case report. *The Journal of Laryngology & Otology*, 123:462-465.
- Ellis, A., Benner, D.L.H., 2013. Neuroinflammation and the generation of neuropathic pain. *British Journal of Anaesthesia*, 111(1):26-37.

- Fauci, B., Kasper, L.H., Jameson, L., 2008. *Harrison's Principles of Internal Medicine 17th Chapter 379: Peripheral neuropathy*. New York: McGraw Hill; pp. 2651–66.
- Frantz, K.J., O'Dell, L.E., Parsons, L.H., 2007. Behavioral and neuro-chemical responses to cocaine in periadolescent and adult rats. *Neuropsychopharmacology*, 32:625–637.
- Gandevia, S.C., 2001. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev.*;81(4):1725-89.
- Ghanei, E., Homayooni, M., Nasrollahi, A., 2007. Rhabdomyolysis and thinner intoxication. *Medical Journal of the Islamic Republic of Iran*; 21(3):162-166
- Gilron, I., Watson, C. P. N., Cahill, C. M., Moulin, D. E., 2006. Neuropathic pain: a practical guide for the clinician. *Canadian Medical Association Journal*, 175:265–275. doi: 10.1503/cmaj.060146
- Glockle, A.H, Cuno, E., Hensel, T., Schultz, H.J., Sandstede, R.P., 1996. *Feldstudie Toluol in Tiefdruckerien*. Wiesbaden: Berufsgenossenschaft Druck und Papierverarbeitung.
- Gospe, S.M., Calaban, M.J., 1988. Central nervous system distribution of inhaled toluene. *Fundamental and applied toxicology*, 11: 540–545.
- Graham, D.I., 1997. Greenfield's neuropathology (6th Edition). *Arnold*; 779-780.
- Graham Hole University of Sussex, 2013. *Graham Hole Research Skills Kruskal-Wallis handout, version 1.0., for autumn 2013*. Brighton: University of Sussex, UK. Diunduh pada tanggal 20 Desember 2014 dari laman <http://www.sussex.ac.uk/Users/grahamh/RM1web/teaching08-RS.html>
- Grzybowski, A., Holder, G.E., 2010. Tobacco optic neuropathy (TON): the historical and present concept of the disease. *Acta Ophthalmologica*, 89(5):495-499.
- Gupta, S., Kline, L., Lincoff, N., Palmer, C., Cure, J., & Balos, L., 2010. *Toluene optic neuropathy: MRI and pathological features*. Poster Presentations at North American Neuro-Ophthalmology Society on Tuesday, March 9, 2010
- Gupta, S.R., Palmer, C.A., Curé, J.K., Balos, L.L., Lincoff, N.S., & Kline, L.B., 2011. Toluene optic neurotoxicity: magnetic resonance imaging and pathologic features. *Hum Pathol*; 42(2): 295-8. doi: 10.1016/j.humpath.2010.08.005.
- Hansen, O.C., Eggert, T., 2003. *Survey of chemical compounds in consumer products: survey, emission, and evaluation of volatile organic chemicals in printed matter*. Göteborg, Swedia: Danish Technological Institute.
- Hapsari, E.D., 2011. *Hubungan antara lama menderitanya dengan kadar glukosa plasma pada penderita diabetes mellitus tipe 2 suku jawa* [Skripsi]. Yogyakarta: Universitas Gadjah Mada.
- Hicks, A.L., Kent-Braun, J., Ditor, D.S., 2001. Sex differences in human skeletal muscle fatigue. *Exerc Sport Sci Rev.*, 29: 109–112, 2001.
- Hinkle, D.E., Wiersma, W., Jurs, S.G., 2003. *Applied Statistics for the Behavioral Sciences (5th Ed)*. Boston: Houghton Mifflin.

- Huang, C-C., 2008. Polyneuropathy Induced by n-Hexane Intoxication in Taiwan. *Acta Neurol Taiwan*, 17 (1):3-10
- Hughes, R.A.C., 2002. Peripheral neuropathy, *BMJ*; 324(7335):466-469.
- Ikeda, M., Ohtsuji, H., 1969. Significance of urinary hippuric acid determination as an index of toluene exposure. *Br J Ind Med*, 26:244-246.
- International Association for the Study of Pain, 2011. *IASP pain terminology*. Washington, DC: International Association for the Study of Pain. Retrieved from www.iasp-pain.org/AM/Template.cfm?Section¼Pain_Definitions
- International Programme on Chemical Safety, 1985. *Environmental health criteria 52 toluene*. Geneve: International Programme on Chemical Safety, WHO.
- International Programme on Chemical Safety, 2000. *Air quality guidelines toluene*. Geneve: WHO.
- Jung, K-H., Jeon, Cu, K., Kim, Y.A., Jeon, B.S., 2007. Rapidly Progressive Toxic Leukoencephalomyelopathy with Myelodysplastic Syndrome: a Clinicopathological Correlation. *J Clin Neurol.*, 3(1):45-49.
- Kang, D-M., & Kim, I., 2014. Compensation for occupational neurological and mental disorders. *J Korean Med Sci*, 29 (Suppl):S59-S65.
- Kathleen, A., 2006. Peripheral Neuropathy: Pathogenic Mechanisms and Alternative Therapies. *Alternative Medicine Review*, 11, 294-329.
- Katulanda, P., Ranasinghe, P., Jayawardena, R., Constantine, G.R., Sheriff, M.H., 2012. The prevalence, patterns and predictors of diabetic peripheral neuropathy in a developing country. *Diabetol Metab Syndr*, 4: 21.
- Kesler A, Pianka P., 2003. Toxic optic neuropathy. *Curr Neurol Neurosci Rep.*, 3:410-414.
- Kiyokawa, M., Mizota, A., Takasoh, M., Adachi-Usami, E., 1999. Pattern visual evoked cortical potentials in patients with toxic optic neuropathy caused by toluene abuse. *Japan J Ophthalmol*, 43(5):438-442.
- Kutlu, G., Gomceli, Y.B., Sonmez, T., Inan, L.E., 2009. Peripheral neuropathy and visual evoked potential changes in workers exposed to n-hexane. *J of Clin Neuroscience*, 16(10):1296-1299.
- Koffman, B.M., Greenfield, L.J., Ali, I.I., Pirzada, N.A., 2006. Neurologic complications after surgery for obesity. *Muscle and Nerve*, 33(2):166-176.
- Kohn, M.C., Melnick, R.L., 1993. Species differences in the production and clearance of 1,3-butadiene metabolites: a mechanistic model indicates predominantly physiological, not biochemical, control. *Carcinogenesis*, 14(4): 619-628.
- Konuk, 2012. Effects of α -lipoic acid on DNA damage, protein oxidation, lipid peroxidation, and some biochemical parameters in sub chronic thinner addicted rats. *Turk J Biology*, 36:702-710.
- Kote, G.S.S., Bhat, A.N., Thajuddeen, K., Ismail, M.H., Gupta, A., 2013. Peripheral insensate neuropathy-is height a risk factor? *J Clin Diagn Res.*, 7(2):296-301; doi: [10.7860/JCDR/2013/5140.2751](https://doi.org/10.7860/JCDR/2013/5140.2751)

- Kramer, L.D., Locke, G.E., 1987. Wernicke's encephalopathy. Complication of gastric plication. *Journal of Clinical Gastroenterology*, 9(5):549–552.
- La Dou, J., 2004. *Lange medical books: current occupational & environmental medicine*. NY: Mc-Graw Hill, Inc.
- Lataye, R., Maguin, K., Campo, P., 2007. Increase in cochlear microphonic potential after toluene administration. *Hearing Research*; 230:34-42.
- Lazar, R.B., & Ho, S.U., 1983. Progressive optic neuropathy and sensorineural hearing loss due to chronic glue sniffing. *J Neurol Neurosurg Psychiatry*, 46(11): 1060.
- Lee, Y.-L., Pai, M.-C., Chen, J., -H., Guo, Y.L., 2003. Central neurological abnormalities and multiple chemical sensitivity caused by chronic toluene exposure. *Occupational Medicine*, 53:479–482
- Lees-Haley, P.R., Williams, C.W., 1997. Neurotoxicity of chronic low-dose exposure to organic solvents: a skeptical review. *J Clin Psychol*; 53:699–712.
- Lennon, S.M., Koblar, S., Hughes, R.A.C., Goeller, J., Riser, A.C., 1993. Reasons for persistent disability in Guillain-Barré syndrome. *Clin Rehab*, 7(1):1–8.
- Lin, Y.S., Smith, T.J., Kelsey K.T., Wypij, D., 2001. Human physiologic factors in respiratory uptake of 1,3-butadiene. *Environ Health Perspect*, 109(9): 921–926.
- Lin, Y.S., Egeghy, P.P., Rappaport, S.M., 2008. Relationships between levels of volatile organic compounds in air and blood from the general population. *Journal of Exposure Science and Environmental Epidemiology*, 18:421–429; doi:10.1038/sj.jes.7500635
- Liu, Y., Rao, D., Fechter, D., 1997. Correspondence between middle frequency auditory loss in vivo and outer hair cell shortening in vitro. *Hearing Research*; 112:134-140
- Lubman, D.I., Yücel, M., Lawrence, A.J., 2008. Inhalant abuse among adolescents: neurobiological considerations. *British Journal of Pharmacology*, 154:316–326.
- Magy, L., Vallat, J.-M., 2008. Nerve biopsy without muscle sampling: is it enough for diagnosing vasculitis? *J Neurol Neurosurg Psychiatry*; 79:1307.
- Manikantan, P., Balachandrar, V., Sasikala, K., Mohanadevi, S., 2010. Cytogenetic methods for assessing human exposure to toluene in Coimbatore, South India. *Asian Pac J Cancer Prev*, 11(6):1687-1693.
- Martinez-Alfaro, M., Alcarazz-Contreras, Y., Cárabez-Trejo, A., Leo-Amador, G.E., 2011. Oxidative stress effects of thinner inhalation. *Indian J Occup Environ Med*, 15(3):87-92.
- Martyn, C., Hughes R.A.C., 1998. *Peripheral neuropathies*. London: BMJ.
- Mäki-Paakkanen, J., Husgafvel-Pursiainen, K., Kalliomäki, P.L., Tuominen, J., Sorsa, M., 1980. Toluene-exposed workers and chromosome aberrations. *J Toxicol Environ Health*; 6:775-781.
- Merck, 2012. *Perhimpunan dokter spesialis saraf (PERDOSSI) dan Merck peduli kesehatan saraf (Siaran Pers)*. Diunduh pada tanggal 25 Oktober 2014 dari:

http://www.merck.co.id/country/id/id/images/Press%20Rilis%20N5000%208Jun12-Ind_tcm663_94645.pdf?Version=

- Merkies, I.S, Schmitz, P.I, Samijn, J.P, van der Meche, F.G., van Doorn, P.A., 1999. European Inflammatory Neuropathy Cause and Treatment (INCAT) Group. Fatigue in immune-mediated polyneuropathies. *Neurology*; 53(8):1648–54
- Michael, Ash, I., 2011. *Handbook of fillers, extenders, and diluents* (2nd Ed.). NY: Synapse Information Resources Inc.
- Mochammad, I., 2012. *Korelasi antara kadar timbal dalam darah dengan neuropati perifer pada petugas stasiun pengisian bahan bakar untuk umum* [Tesis] Yogyakarta: Universitas Gadjah Mada.
- Mold, J.W., Vesely, S.K., Keyl, B.A., Schenk, J.B., Roberts, M., 2004. The Prevalence, Predictors, and Consequences of Peripheral Sensory Neuropathy in Older Patients. *J Am Board Fam Pract.*, 17:309–18.
- Mold, J.W., Vesely, S.K., Keyl, B.A., 2010. The prevalence, predictors, and consequences of peripheral sensory neuropathy in older patients. *J Am Board Fam Pract.*, 17(5):309-318.
- Morley-Fletcher, S., Bianchi, M., Gerra, G., Laviola, G., 2002. Acute and carryover effects in mice of MDMA ('ecstasy') administration during periadolescence. *Eur J Pharmacol.*, 448:31–38
- Moszyński, P., Rutowski, J., Słowiński, S., 1996. The effect of cigarettes smoking on the blood counts of T and NK cells in subjects with occupational exposure to organic solvents. *Cent Eur J Public Health*, 4 (3):164-168.
- Motz, G.T., Eppert, B.L., Borchers, M.T., 2010. Chronic Cigarette Smoke Exposure Generates Pathogenic T Cells Capable of Driving COPD-like Disease in Rag2^{-/-} Mice. *Am J Respir Crit Care Med.*, 181(11):1223-1233.
- MØLHAVE, L. & PEDERSEN, O.F. Measurements of alveolar concentrations of toluene. *International archives of occupational and environmental health*, 54: 65–71 (1984).
- Mørck, H.I., Winkel, P., Gyntelberg, F., 1988. Health effects of toluene exposure. *Dan. Med. Bull.*; 35:196-200.
- Mukaka, M., 2012. A guide to appropriate use of Correlation coefficient in medical research. *Malawi Medical Journal: The Journal of Medical Association of Malawi*, 24(3):69-71.
- Murena, F., 2007. Air quality nearby road traffic tunnel portals: BTEX monitoring. *Journal of Environmental Science*, 19:578-583.
- National Institute for Occupational Safety and Health, 1978. *Occupational health guideline for toluene*. Atlanta, GA: CDC.
- National Institute for Occupational Safety and Health, 2003. *NIOSH 150: hydrocarbons aromatic, NIOSH Manual of Analytical Methods (Issue 2)*. Atlanta, GA: Centers for Disease Control & Prevention (CDC).
- Nguyen, T.P., Chaudry, V., 2011. Multifocal motor neuropathy. *Neurol India*; 59(5): 700-706.
- Nikolajsen, L., 2012. Postamputation pain: studies and mechanism. *Dan Med J.*, 59(10):B4527.

- Nikolajsen, L., & Jensen, T.S., 2001. Phantom limb pain. *Br J Anaesth*, 87:107-116.
- Nise, G., Attewell, R., Skrefving, S., Ørbæk, P., 1989. Elimination of toluene from venous blood and adipose tissue after occupational exposure. *British Journal of Industrial Medicine*, 46:407-411.
- Novella, S.P., Inzucchi, S.E., Goldstein, J.M., 2001. The frequency of undiagnosed diabetes and impaired glucose tolerance in patients with idiopathic sensory neuropathy. *Muscle Nerve*, 24: 1229–31
- Occupational Safety and Health Administration, 2012. *OSHA infosheet: Toluene safety in the workplace*. Washington, DC: US. Department of Labor.
- Odebode, T.O., Odebode, S.O., 2005. Protein Energy Malnutrition and the Nervous System: the Impact of Socioeconomic Condition, Weaning Practice, Infection and Food Intake, an Experience in Nigeria. *Pakistan Journal of Nutrition*, 4(5):304-309.
- Palencia, G., Hernández-Pedro, N., Saavedra-Perez, D., Peña-Curiel, O., Ortiz-Plata, A., Ordoñez, G., Flores-Estrada, D., Sotelo, J. and Arrieta, O., 2014. Retinoic acid reduces solvent-induced neuropathy and promotes neural regeneration in mice. *J. Neurosci. Res.*, 92:1062–1070. doi: 10.1002/jnr.23376
- Parra, M.A., Elustondo, D., Bermejo, R., Santamaría, J.M., 2009. Ambient air levels of volatile organic compounds (VOC) and nitrogen dioxide (NO₂) in a medium size city in Northern Spain. *Sci Total Environ*, 407:999-1009
- Perkins, B.A, Orszag, A., Ngo, M., Ng, E., New, P., 2010. Prediction of incident diabetic neuropathy using the monofilament examination: a 4-year prospective study. *Diabetes Care*, 33: 1549-1554.
- Pfeiffer, G., Wicklein, E.M., Ratusinski, T., Schmitt, L., Kunze, K., 2001. Disability and quality of life in Charcot-Marie-Tooth disease type 1. *Journal of Neurology, Neurosurgery and Psychiatry*;70(4):548–50.
- Puri, V., Chaudry, N., Tatke, M., 2007. N-hexane neuropathy in scree printers. *Electromyogr Clin Neurophysiol.*; 47(3):145-52.
- Poncelet, A.N., 1998. An algorithm for the evaluation of peripheral neuropathy. *Am Fam Physician*; 57(4):755-764.
- Pradeepa, R., Remana, J., Vignesh, J., 2008. Prevalence and risk factors for diabetic neuropathy in an urban south Indian population: the Chennai urban rural epidemiology study (CURES-55). *Diabetic Medicine*, 25:407-412.
- Pyor, G.T., 1991. A toluene-induced motor syndrome in rats resembling that seen in some human solvent abusers. *Neurotoxicology and teratology*; 13(4):387-400.
- Rahayu, D.P., 2013. *Deteksi toksisitas pajanan toluene terhadap penurunan jumlah sel sertoli tikus wistar jantan* [Tesis]. Depok: Fakultas Kedokteran Universitas Indonesia.
- Rebert, C.S., Schwartz, R.W., Svendsgaard, D.J., Pryor, G.T., Boyes, W.K., 1995. Combined effects of paired solvents on the rat's auditory system. *Toxicology*; 105:345-354.

- Reza, J., Salazar, G., Trejo, A., 1997. Evaluation of composition and evaporation behavior of commercial thinner samples expended in Mexico City. *Rev Int Contam Ambient.*, 13:87–95.
- Riegel, A.C., Zapata, A., Shippenberg, T.S., French, E.D., 2007. The abused inhalant toluene increases dopamine release in the nucleus accumbens by directly stimulating ventral tegmental area neurons. *Neuropsychopharmacology*, 32:1558–1569.
- Ropper, A. H., Samuels, M. A., 2000. Disease of peripheral nerve. *Adams & Victor's Principles of Neurology (7th Edition)*. New York: The McGraw Hill.
- Rusdy, M.D.R., 2012. *Analisis gejala neurotoksik akibat pajanan xylene pada pekerja pembuatan cat PT. X tahun 2012* [Tesis]. Depok: Universitas Indonesia.
- Scelsi, R., Candura, S.M., 2012. Occupational toxic neuropathies: morphology in peripheral nerve biopsies. *G Ital Med Lav Ergon.*; 34(4):410-419.
- Shigeta, S., Misawa, T., Aikawa, H., 1980. Effects of concentration and duration of toluene exposure on Sidman avoidance in rats. *Neurobehav Toxicol.*, 2(2): 85-88.
- Smith, A.G., Singleton, J.R., 2013. Obesity and hyperlipidemia are risk factors for early diabetic neuropathy. *J Diabetes Complications*, 27: 436-442
- Smith, H.S., Meek, P.D., 2011. Pain responsiveness to opioids: central versus peripheral neuropathic pain. *J Opioid Manag.*; 7(5):391-400.
- Sophianita, 2003. *Hubungan antara kadar asam hipurat urin akibat pajanan toluen dengan efek kesehatan akut pada tenaga kerja percetakan "X"* [Karya ilmiah]. Jakarta: Universitas Indonesia.
- Sorensen, L., Molyneaux, L., Yue, D.K., 2002. Insensate versus painful diabetic neuropathy: the effects of height, gender, ethnicity, and glycaemic control. *Diabetes Res Clin Pract*, 57(1):45-51.
- Spear, L.P, 2000. Modeling adolescent development and alcohol use in animals. *Alcohol Res Health*, 24:115–23. [PubMed]
- Spear, L.P., 2000. The adolescent brain and age-related behavioral manifestations. *Neuroscience and Biobehavioral Reviews*, 24:417–63. [PubMed]
- Stubblefield, M.D., Burstein, H.J., Burton, A.W., Custodio, C.M., Deng, G.E., Ho, M., 2009. NCCN Task Force Report: Management of Neuropathy in Cancer, *Journal of the National Comprehensive Cancer Network*; 7(Suppl 5).
- Suzuki, R., Dickenson, A., 2005. Spinal and supraspinal contributions to central sensitization in peripheral neuropathy. *Neurosignals*; 2005;14(4):175-81.
- Svendsen, K., Rognes, K.S., 2000. Exposure to organic solvents in the offset printing industry in Norway. *Ann. Occup. Hyg.*; 44(2):119-124.
- Tanaka, K., Maeda, T., Kobayashi, T., 2003. A survey of urinary hippuric acid and subjective symptoms among occupational low toluene exposed workers. *Fukushima J. Med. Sci.*; 49:129-139.
- Tavee, J., 2014. *Overview of peripheral neuropathies (1st consult)*. Philadelphia, PA: Elsevier BV.

- Tokunaga, R., Takahata, S., Onoda, M., Ishi-i, T., Sato, K., Hayashi, M., Ikeda, M., 1974. Evaluation of the exposure to organic solvent mixture. Comparative studies on detection tube and gas-liquid chromatographic methods, personal, and stationary sampling and urinary metabolite determination. *Int Arch Arbeitsmed*; 33:257-267.
- Troy, J., Jensen, R., 2002. New perspectives on the management of diabetic peripheral neuropathic pain. *Diabetes and Vascular Disease Research*, 3(2):108-119.
- Tseng, C.H., 2006. Prevalence of lower-extremity amputation among patients with diabetes mellitus: is height a factor? *MAJ*, 174:319–23.
- Tyle, H., Larsen, H.S., Andersen, L.K., Brandt-Lassen, R., 2003. *European Union Risk Assessment Report: toluene*. Luxembourg: Office for Official Publications of the European Communities.
- US Environmental Protection Agency, 1985. *Method 524.1: volatile organic compounds in water by purge-and-trap gas chromatography/mass spectrometry*. Cincinnati, OH: US Environmental Protection Agency, Environmental Monitoring and Support Laboratory.
- von Burg, R., 1993. Toxicology update: toluene. *Jour f Appl Toxicology*; 13: 441–446
- von Euler, M., Pham, T.M., Hillefors, M., Bjelke, B., Henriksson, B., von Euler, G., 2000. Inhalation of low concentrations of toluene induces persistent effects on a learning retention task, beam-walk performance, and cerebrocortical size in the rat. *Exp Neurol*, 163: 1–8.
- WHO, 1995. *Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Consultation. WHO Technical Report Series Number 854*. Geneva: World Health Organization.
- WHO, 1997. *Xylenes, Environmental Health Criteria 190*. Geneva, Switzerland: WHO.
- WHO, 2006. *Neurological disorders: public health challenges - Chapter 3.6: Neurological disorders associated with malnutrition*. Geneva: World Health Organization.
- Warsito, A., 2007. *Analisis pemajanan toluena terhadap profil darah pada pekerja sektor industri penyulingan minyak bumi* [Tesis] Semarang: Universitas Diponegoro.
- Wee, J.Y.M., Hopman, W.M., 2008. Effects of smoke exposure and other lifestyle factors on pain response to electrical stimulation in women. *Pain Res Manage*, 13(3):231-235.
- White, C.M., van Doorn P.A., Garssen, M.P.J, Stockley R.C., 2009. Interventions for fatigue in peripheral neuropathy. *Cochrane Database of Systematic Reviews*; Issue 4. Art. No.: CD008146. doi: 10.1002/14651858.CD008146
- Wiaderna, D., Tomas, T., 2002. Assessment of long term effects of exposure to toluene based on the analysis of selected behavioral responses with particular reference to the ability to trigger behavioral hypersensitivity in rats. *Int J Occup Med Environ Health*, 15(3):239-245.
- Wiles, C.M., 1996. Neurological complications of severe illness and prolonged mechanical ventilation. *Thorax*; 51(Suppl 2):S40–44.

- Wiley, J.L., Evan, R.L., Grainger, D.B., Nicholson, K.L., 2008. Age-dependent differences in sensitivity and sensation to cannabinoids and “club drugs” in male adolescent and adults rats. *Addict Boil.*, 13(3-4):277-286.
- Wills, A.J., O`Connor, S., McMillan, A., 2008. Sub-acute demyelinating neuropathy associated with an NK/T cell lymphoma. *J Neurol Neurosurg Psychiatry*, 79:484-485.
- Windebank, A.J., 1999. Chemotherapeutic neuropathy. *Curr Opin Neurol.*; 12(5):565-571.
- Wjst, M., Reitmar, P., Dold, S., Wulff, A., Nicolai, T., Von Loeffelholz-Colberg, E.F., 1993. Road traffic and adverse effects on respiratory health in children. *BMJ*, 307:596-600.
- Woodward, J.J., Beckley, J., 2014. Effects of the abused inhalant toluene on the mesolimbic dopamine system. *J Drug Alcohol Res.*, 3:doi:10.4303/jdar/235838
- Yüksel, G., Karlikaya, G., Tutkavul, K., Yildirim, B., Örken, C., Tireli, H., 2007. Clinical and electrodiagnostic findings of n-hexane neuropathy. *J Neurol Sci [Turk]*; 24:219-225.
- Zavalic, M., Mandic, Z., Turk, R., 1998. Quantitative assessment of color vision impairment in workers exposed to toluene. *Am J Ind Med*; 33(3):297-304.
- Zhang, Y., Mu, Y., Liu, J., Mellouki, A., 2012. Levels, sources and health risks of carbonyls and BTEX in the ambient air of Beijing. *China. J Environ Sci (China)*, 24:124-30
- Ziegler, D., Rathmann, W., Dickhaus, T., Meisinger, C., Mielck, A., KORA Study Group, 2008. Prevalence of polyneuropathy in pre-diabetes and diabetes is associated with abdominal obesity and macroangiopathy: the MONICA/KORA Augsburg Surveys S2 and S3. *Diabetes Care*, 31: 464-469.
- Keputusan Presiden Republik Indonesia, Nomor 22, Tahun 1993, tentang Penyakit yang Timbul karena Hubungan Kerja.
- Keputusan Menteri Tenaga Kerja Republik Indonesia, Nomor Kep. 187/Men/1997, Tahun 1997, tentang Pengendalian Bahan Kimia Berbahaya di Tempat Kerja.