

## DAFTAR PUSTAKA

- Anderson, J.W., Smith, B.M., Washnockk, C.S. (1999). Cardiovascular and renal benefits of dry bean and soybean intake. *Am J Clin Nutr*, 70(3):464-74.
- Astuti. M. 1999. History of the Development of Tempe. Di dalam Agranoff J., pp.:2-13.
- Atmaca, G. (2004). Antioxidant effects of sulfur-containing amino acids. *Yonsei Med J*, 45:776-788.
- Azadbakht, L., Kimiagar, M., Mehrabi, Y., Esmailzadeh, A., Frank, B. Hu, Walter, C.W. (2007). Soy Consumption, Markers of Inflammation, and Endothelial Function: A cross-over study in postmenopausal women with the metabolic syndrome. *Diabet C*, 30(4):967-973.
- Barik, R., Jain, S., Qwatra, D., Joshi, A., Tripathi, G.S., Goyal, R. (2008). Antidiabetic Activity of Aqueous Root Extract of *Ichnocarpus frutescens* in Sterptozotocin-Nicotinamide Induced Type-II Diabetes in Rats. *Indian J Pharmacol*. 40(1):19-22.
- Bastard, J.P., Maachi, M., Lagathu, C., Kim, M.J., Caron, M., Vidal, H., Capeau, J., Feve, B. (2006). Recent advances in the relationship between obesity, inflammation, and insulin resistance. *Eur Cytokine Netw*, 17(1):4-12.
- Beavers, K.M., Serra, M.C., Beavers, D.P., Cooke, M.B., Willoughby, D.S. (2009). Soymilk supplementation does not alter plasma markers of inflammation and oxidative stress in postmenopausal women. *Nut Res*, 29:616-22
- Boden, G., Shulman, G.I. (2002). Free fatty acids in obesity and type 2 diabetes: defining their role in the development of insulin resistance and beta-cell dysfunction. *Eur J Clin Invest*, 32(3):14-23.
- Borradaile, N.M., de Dreu, L.E., Wilcox, L.J., Edwards, J.Y., Huff, M.W. (2002). Soya Phytoestrogens, Genistein and Daidzein Decrease Apolipoprotein B Secretion from HepG2 Cells through Multiple Mechanisms. *Biochem J*, 366(2):531-9.
- Braunwald, E. (2008). Biomarkers in Heart Failure. *N Engl J Med*. 358:2148-2159.
- \_\_\_\_\_. (1997). Shattuck lecture - cardiovascular medicine at the turn of the millennium: triumphs, concerns, and opportunities. *N Engl J Med*, 337:1360-1369.

- Brownlee, M.(2005). The Pathobiology of Diabetic Complications: A Unifying Mechanism. *Diab*, 54(6):1615-25.
- Cade, W.T.(2008). Diabetes-related Microvascular and Macrovascular Diseases in The Physical Therapy Setting. *Phys Ther*, 88(11):1322-35.
- Captoge, S., Di Angelantonio, E., Lowe, G., Pepys, M.B., Thompson, S.G., Collins, R., Danesh, J.(2010). C-reactive protein concentration and risk of coronary heart disease, stroke, and mortality: an individual participant meta-analysis. *Lancet*, 375:132-140.
- Cheng, K.C., Lin, J.T., Liu, W.H.(2011). Extracts from Fermented Black Soybean Milk Exhibit Antioxidant and Cytotoxic Activities. *Food Technol Biotechnol*, 49(1):111-17.
- Chin-Lin, H., dan Gow-Chin, Y.(2007). Effects of Flavonoids and Phenolic Acids on the Inhibition of Adipogenesis in 3T3-L1, *J Agri Food Chem*, 55 (21):8404-8410.
- Choudhary, N., Ahlawat, R.S.(2008). Interleukin-6 and C-Reactive Protein in Pathogenesis of Diabetic Nephropathy: New Evidence Linking Inflammation, Glycemic Control, and Microalbuminuria. *Iran J Kid Dis.*, 2(2):72-9.
- Dajanta, K., Chukeatirote, E., Apichartsrangkoon, A., Frazier, R.A.(2009). Enhanced Aglycone Production of Fermented Soybean Products by *Bacillus* Species. *Acta Biologica Szegediensis*, 53(2):93-98.
- DeFronzo, R.A., Simonson, D., Ferrannini, E.(1982). Hepatic and Peripheral Insulin Resistance: A Common Feature of Type 2 (Non-Insulin-Dependent) and Type 1 (Insulin-Dependent) Diabetes Mellitus. *Diabetologia*, 23(4):313-9.
- Dini (2010). *Kompas*. [Online] Available at: <http://megapolitan.kompas.com> [Accessed 03 April 2014].
- Dong, J.Y., Wang, P., He, K., Qin, L.Q.(2011). Effects of soy isoflavones on circulating C-reactive protein in postmenopausal women: meta-analysis of randomized controlled trials. *Meno*, 18(11):1256-62.
- Ejtahed, H.S., Mohtadi-Nia, J., Homayouni-Rad, A., Niafar, M., Asghari-Jafarabadi, M., Mofid, V.(2012). Probiotic Yogurt Improves Antioxidant Status in Type 2 Diabetic Patients. *Nut*, 28(5):539-43.

- Elmarakby, A.A. dan Sullivan, J.C.(2012). Relationship between Oxidative Stress and Inflammatory Cytokines in Diabetic Nephropathy. *Cardio Ther*, 30:49-59.
- EOL, 2014. EOL. [Online] Available at: <http://eol.org> [Accessed 6 April 2014].
- Fanti, P., Asmis, R., Stephenson, T.J., Sawaya, B.P., Frankee, A.A.(2006). Positive Effect of Dietary Soy in ESRD Parients with Systemic Inflammation- Correlation Between Blood Levels of The Soy Isoflavones and The Acute-Phase Reactans. *Nephrol Dial Transplant*, 21(8):2239-46.
- Farnworth, E.(2005). *Kefir- A Complex Probiotic*, s.l.: Food Science and Technology Bulletin: Functional Food.
- Filiberto, A.C., Mumford, S.L., Pollack, A.Z., Zhang, C., Yeung, E.H., Perkins, N.J., et al.(2013). Habitual Dietary Isoflavone Intake Is Associated with Decreased C-Reactive Protein Concentrations Among Healthy Premenopausal Women. *J Nutr*, 143(6):900-6.
- Fowler, M.(2008). Microvascular and Macrovascular Complications of Diabetes. *Clin Diab*, 26(2):77-82.
- Ghozi, M., Agni, A.N., Gunawan, Ulfah, N.M., Gunawan, W. (1991). Ilmu Penyakit Mata IV. Laboratorium Ilmu Penyakit Mata. Fakultas Kedokteran UGM, Yogyakarta.
- Giacco, F., Brownlee, M.(2010). Oxidative Stress and Diabetic Complications. *Circ Res*, 107(9):1058-70.
- Ginter, E. dan Simko,V.(2012). Global Prevalence and Future of Diabetes Mellitus. *Adv Exp Med Biol.*, Volume 771:35-41.
- Ginter, E. dan Vlado, S.(2012). Endocrine. In: *Diabetes: An Old Disease, A new Insight*. s.l.:Springer.
- Goldberg, R.(2009). Cytokine and Cytokine-like Inflammation Markers, Endothelial Dysfunction, and Imbalanced Coagulation in Development of Diabetes and Its Complications. *J Clin Endocrinol Metab*, 94(9):3171-82.
- Guacuc standard procedure retro-orbital sinus/plexus blood collection in mice and rats.(2006). Available in PDF.
- Gunawan H.(2014). *Tribun News*. [Online] Available at: <http://www.tribunnews.com> [Accessed 3 April 2014].
- Hadisaputro, S., Djokomoeljanto, R.R.J., Judiono, Soesatyo.(2012). The Effects of Oral Plain Kefir Supplementation on Proinflammatory Cytokine

- Properties of the Hyperglycemia Wistar Rats Induced by Streptozotocin. *Acta Med Indones.*, 44(2):100-4.
- Haenlein, G.F.W.(2004). Goat milk in human nutrition. *Small Rum Research*, 51:155-163.
- Halliwell, B.(2005). Free Radicals and Other Reactive Species in Disease. eLS.
- Held, P.(2010). *An Introduction to Reactive Oxygen Species - Measurement of ROS in Cells*, s.l.: s.n.
- Hernández-Ledesma, B., Recio, I., Amigo, L.(2008). Lactoglobulin as source of bioactive peptides. *Amino Acids*, 35:257-265.
- Higdon, J.(2009). *Linus Pauling Institute*. [Online] Available at: [lpi.oregonstate.edu](http://lpi.oregonstate.edu) [Accessed 06 April 2014].
- Hulley, Stephen, B., Cumming, Steven, R., Grady, D.(2007). Designing a Randomized Blinded Trial, Chapter 10. Designing Clinical Research, 3rd Edition. Philadelphia, USA: Lippincott Williams & Wilkins. pp.:147 - 161.
- Jakus, V., dan Rietbrock, N.(2004). Advanced Glycation End-Products and The Progress of Diabetic Vascular Complications. *Physiol Res*, 53(2):131-42.
- Kano, M., Takayanagi, T., Harada, K., Sawada, S., Ishikawa, F.(2006). Bioavailability of isoflavones after ingestion of soy beverages in healthy adults. *J Nut*, 136:2291-2296.
- Kaptoge, S., Di Angelantonio, E., Lowe G., Pepys, M.B., Thompson, S.G., Collins, R, et al.(2010). C-reactive Protein Concentration and Risk of Coronary Heart Disease, Stroke, and Mortality: An Individual Participant Meta-Analysis. *Lancet*, 375(9709):132-40.
- Karunakaran, U. dan Park, K-G.(2013). A Systematic Review of Oxidative Stress and Safety of Antioxidants in Diabetes: Focus on Islets and Their Defense. *Diab Met J*, 37(2):106-12.
- Kasenkas, H., Dinkci, N., Seckin, K., Kinik, O., Gonc, S.(2011). Antioxidant Properties of Kefir Produced from Different Cow and Soy Milk Mixtures. *J Agric Sci.*, 17:253-9.
- Kim, Y.G., Lohakare, J.D., Yun, J.H., Heo, J., Chae, B.J.(2007). Effect of feeding levels of microbial fermented soy protein on the growth performance, nutrient digestibility and intestinal morphology in weaned piglets. *Asian-Aust. J. Anim. Sci.*, 20:399-404.

- Koya, D.(2000). Amelioration of Accelerate Diabetic Mesangial Expansion by Treatment by PKC beta Inhibitor in Diabetic Mice, a Rodent Model for Type 2 Diabetes. *The FASEB J.* 14:439-47.
- Kuklinska, A.M., Mroczko, B., Musial, W.J., Sawicki, R., Kozieradzka, A., Waszkiewicz, E., Szmitkowski, M.(2009). High-sensitivity C-reactive protein and total antioxidant status in patients with essential arterial hypertension and dyslipidemia. *Adv Med Sci*; 54(2):225-32.
- Kullisar, T.(2003). Antioxidative probiotic fermented goat's milk decreases oxidative stress-mediated atherogenicity in human subjects. *B J Nut*, 90:449-456.
- Kwon, D.Y., Jang, J.S., Lee, J.E., Kim, Y.S., Shin, D.H., Park, S.(2006). The Isoflavonoid Aglycone-Rich Fractions of Chungkookjang, Fermented Unsalted Soybeans, Enhance Insulin Signaling and Peroxisome Proliferator-Activated Receptor-Gamma Activity In Vitro. *Bifactors*, 26(4):245-58.
- Kwon, K.B., Kim, E.K., Jeong, E.S., Lee, H.Y., Lee, Y.R., Park, J.W., Ryu, D.G., Park, B.H.(2006). Cortex cinnamomi extract prevents streptozotocin and cytokine-induced beta-cell damage by inhibiting NF-B. *World J Gastroenterol*, 12(27):4331-37.
- Lara-Villoslada, F., Debras, E., Nieto, A., Concha, A., Galves, J., Lopez-Huertas, E.(2004). Oligosaccharides Isolated from Goat Milk Reduce Intestinal Inflammation in A Rat Model of Dextran Sodium Sulfate-Induced Colitis. *Clin Nut*, 25(3):477-88.
- Ling, Z., Pipeleers, D.G.(1994). Preservation of glucose-responsive islet beta cells during serum-free culture. *End*, 134:2614-2621.
- Liu, J.R., Chen, M.J., Lin, C.W.(2005). Antimutagenic and Antioxidant Properties of Milk-Kefir and Soymilk-Kefir. *J Agric Food Chem.*, 53(7):2467-74.
- Liu, J.R., Wang, S.Y., Lin, Y.Y., Lin, C.W.(2002). Antitumor Activity of Milk Kefir and Soy Milk Kefir in Tumor-Bearing Mice.. *Nutr Cancer*, 44(2):183-7.
- Londhe, G.(2014). Effect of different levels of goat milk, soy milk and cow milk on chemical composition of rasogolla. *J Veterinar Sci Technolo*, 5(3):180.

- Lv, Y.Y., Jin, Y., Han, G.Z., Liu, Y.X., Wu, T., Liu, P.(2012). Ursolic Acid Supresses IL-6 Induced C-reactive Protein Expression in HepG2 and Protects HUVECs from Injury Induced by CRP. *Eur J Pharm Sci*, 45(1-2):190-94.
- Mahasneh, A.M., dan Abbas, M.M.(2010). Probiotics and traditional fermented foods: the eternal connection. *Jordan J Biol Sci.*, 3(4):133-140.
- Majello, B., Arcone, R., Tonniati, C., Ciliberto, G.(1990). Constitutive and IL-6-induced Nuclear Factors That Interact with Teh Human C-reactive Protein Promoter. *EMBO J*, 9(2):457-65.
- Marczuk-Krynicka, D., Hryniewiecki, T., Piatek, J., Paluszak, J.(2003). The effect of brief food withdrawal on the level of free radicals and other parameters of oxidative status in the liver. *Med Sci Monit*, 9(3):131-5.
- Marwah, M.P., Suranindyah, Y.Y., Murti, T.W.(2012). Produksi dan Komposisi Susu Kambing Peranakan Ettawa yang Diberi Suplemen Daun Katu (*Sauropus androgynus* (L.) Merr) pada Awal Masa Laktasi (Milk Production and Milk Composition of Ettawa Crossbred Goat, Fed Katu Leaves (*Sauropus androgynus* (L.) Merr) as. *Buletin Peternakan*, 34(2):94-102.
- Mayer, C., Gruber, H.J., LandL, E.M., Pailer, S., Scharnagl, H., Truschnig-Wilders, M., März, W.(2007). Rosuvastatin Reduces Interleukin-6-Induced Expression of C-Reactive Protein in Human Hepatocytes in a STAT3- and C/EBP-Dependent Fashion. *Int J Clin Phar Ther.*, 45(6):319-27.
- Meyer, M., Schreck, R., Baeuerle, P.A.(1993). H2O2 and Antioxidants Have Opposite Effects on Activation of NF-kappa B and AP-1 in Intact Cells: AP-1 as Secondary Antioxidant-responsive Factor. *EMBO J*, 12:2005-15.
- Nanri, A., Moore, M.A., Kono, S.(2007). Impact of C-reactive Protein on Disease Risk and Its Relation to Dietary Factors. *Asian Pac J Cancer Prev*, 8(2):167-77.
- Nelson, D.L. dan Cox, M. M. (2005). *Principles of Biochemistry* (4th ed.). New York: W. H. Freeman
- Nielsen, I.L., Williamson, G.(2007). Review of the factors affecting bioavailability of soy isoflavones in humans. *Nutr Cancer*;57:1-10.
- Nieuwenhuis, R., dan Nieuweik, J.(2005). Soya in the Daily Diet. In: *Cultivating of Soya and Other*

- Legumes. pp.:40-9. Aromisa Foundation, Wageningen, Netherlands.
- Nurliyani, Harmayani, E., dan Soesatyo, M.H.N.E.(2010). The effect of goat milk supplementation on Th1 and Th2 responses of Peyer's patch lymphocyte culture in dinitrochlorobenzene sensitized rat. *WASET*, 68:926-931
- Oakenfull, D.G.(2001). Soy protein, saponins and plasma cholesterol. *J. Nutr*, 131:2971.
- Otles, S., dan Cangindi, O.(2003). Kefir: A Probiotic Dairy-Composition, Nutritional and Therapeutic Aspect. *Pakistan J Nutr*. 2(2):54-59.
- Patel, R.P., McAndrew, J., Sellak, H., White, C.R., Jo, H., Freeman, BA., Darley-Usmar, V.M.(1999). Biological aspects of reactive nitrogen species. *Biophys Acta*, 1411:385-400.
- Powers, A.C.(2006). Diabetes Melitus. In : Jameson, L.J., ed. *Harisson's Endocrinology*. USA: The McGraw-Hill Companies, pp.:314-317.
- Pratama, SE., dan Probosari, E.(2012). Pengaruh Pemberian Kefir Susu Sapi terhadap Kadar Kolesterol LDL Tikus Jantan *Sparague Dawley* Hiperkolesterolemia. *J Nutr Colleague*, 1(1):358-64.
- Rattray, F.P., dan O'Connel, M.J.(2011). Kefir: Fermented Milks. pp.:1-10. Elsevier Saunders. Philadelphia.
- Reeves, .PG., Neilsen, F.H., Fahey, J.R.(1993). AIN-93 Purified Diets for Laboratory Rodents: Fianal Report of the American Institute of Nutrition Ad Hoc Writing Committe on the Formulation of the AIN-76A Rodent Diet. *J Nutr.*, 123(11):1939-51.
- Sartika, C.R.(2006). Penanda Inflamasi, Stress Oksidatif dan Disfungsi Endotel pada Sindroma Metabolik. *Forum Diagnosticum*. Prodia Diagnostics Educational Services. No. 2.
- Schmidt, A.M., Yan, S.D., Wautier, J.L., Stern,D.(1999). Activation of Receptor for Advanced Glycation End Products: A Mechanism for Chronic Vascular Dysfunction in Diabetic Vasculopathy and Atherosclerosis. *Circ Res*, 84:489-97.
- Schmitz, F.J., Ksebati, M.B., Gunasekera, S.P., Agarwal, S.(1988). *J Org Chem*, 53(25):5941-47.
- Schreck, R., Rieber, P., Baeuerle, P.A.(1991). Reactive Oxygen Intermediates as Apparently Widely Used Messengers in The Activation of the NF-kappa B Transcription Factor and HIV-1. *EMBO J*,10:2247-58.

- Seo, S.M., Baek, S.H., Jeon, H.K., Kang, S.M., Kim, D.S., Kim, W.S. (2013). Correlations Between The Level of High Sensitivity C-Reactive Protein and Cardiovascular Risk Factors in Korean Adults with Cardiovascular Disease or Diabetes Mellitus: A CALLISTO Study. *J Atheroscler Thromb.*, 20(7):616-22
- Shilpa, V., Hati, S., Yadav, D. (2011). Biofunctionality of probiotic soy yoghurt. *FNS*, 2(5):223-50.
- Song, J., Kwon, O., Chen, S., Daruwala, R., Eck, P., Park, J.B. (2002). Flavonoid inhibition of sodium-dependent Vitamin C transporter 1 (SVCT1) and glucose transporter isoform 2 (GLUT2), intestinal transporters for vitamin C and glucose. *J Biol Chem*, 277:15252-60.
- Songisepp E. Evaluation of technological and functional properties of the new probiotic *Lactobacillus fermentum* ME-3. *Dissertationes Medicinae Universitatis Tartuenss. Tartu*, 2005.
- Suryaatmadja, M. (2003). High sensitive C-reactive Protein : Parameter Baru Resiko Kardiovaskular (dalam) Pendidikan Berkesinambungan Patologi Klinik, pp.:93-100.
- Szkudelski, T. (2012). Streptozotocin-nicotinamide-induced diabetes in the rat. Characteristics of the experimental model. *Exp Biol Med*, 237(5):481-90.
- Targher, G., Bertolini, L., Zoppini, G., Zenari, L., Falezza, G. (2005). Increased Plasma Markers of Inflammation and Endothelial Dysfunction and Their Association with Microvascular Complications in Type 1 Diabetic Patients without Clinically Manifest Macroangiopathy. *Diabet Med*, 22(8):999-1004.
- Teixeira, S.R., Tappenden, K.A., Carson, L., Jones, R., Prabhudesai, M., Marshall, W.P., Erdman, J.W. Jr. (2004). Isolated Soy Protein Consumption Reduces Urinary Albumin Excretion and Improves the Serum Lipid Profile in Men with Type 2 Diabetes Mellitus and Nephropathy. *J Nutr*, 134(8):1874-80.
- Teruya, K., Yamashita, M., Tominaga, R., Nagira, T., Shim, S.Y., Katakura, Y. (2002). Fermented Milk, Kefir Enhances Glucose Uptake Into Insulin-Responsive Muscle Cells. *Cytotech*, 40(1-3):107-16.
- Thoreux, K., dan Schmucker, D.L. (2001). Kefir milk enhances intestinal immunity in young but not old rats. *Jour of Nut*, 131:807-12.

- Tremblay F, Marette A. (2001). Amino Acid and Insulin Signaling Via The mTOR/p70 S6 Kinase Pathway. A Negative Feedback Mechanism Leading to Insulin Resistance in Skeletal Muscle Cells. *J Biol Chem*, 276(41):38052-60.
- USDA, (2014). USDA. [Online] Available at: <http://ndb.nal.usda.gov> [Accessed 7 April 2014].
- Virtanen, T., Pihlanto, A., Akkanen, S., Korhonen, H. (2007). Development of antioxidant activity in milk whey during fermentation with lactic acid bacteria. *Jour of Appl Micro*, 102:106-115.
- Volanakis, J.E. (2001). Human C-Reactive Protein: Expression, Structure, and Function. *Mol Immunol*, 38(2-3):189-97.
- WHO, 2011. *Fact Sheet: Diabetes*. [Online] Available at: [www.who.int](http://www.who.int) [Accessed 26 Juni 2013].
- WHO, 2013. *Fact Sheet: Diabetes*. [Online] Available at: [www.who.int](http://www.who.int) [Accessed 26 Juni 2013].
- Yayu, Z., Maheswari, R.R.A., Susanty, H. (2011). *Karakteristik Kualitas Susu Segar Dan Yoghurt*. s.l., Seminar Nasional Teknologi Peternakan dan Veteriner.
- Zimmer, Z., Pattama, A. (2001). Socioeconomic status and health among older adults in Thailand: an examination using multiple indicators, *Soc Sci & Med*, 52(8):1297-1311.