



DAFTAR PUSTAKA

- American Thoracic Society, 2002, ATS Statement:
Guidelines for the Six-Minute Walk Test, *American Journal of Respiratory and Critical Care Medicine*, vol. 166, pp. 111-7.
- Callens E, Graba S, Gillet-Juvin K, Essalhi M, Bidaud-Chevalier B, Peiffer C, et al. 2009, Measures of Dynamic Hyperinflation After a 6-Minute Walk Test in Patients With COPD, *Chest*, vol. 136, pp. 1466-72.
- Chakrabarti B, Calverley PMA, Davies PDO, 2007, Tuberculosis and its incidence, special nature, and relationship with chronic obstructive pulmonary disease, *International Journal of COPD*, vol. 2, no. 3, pp. 263-72.
- Crisafulli E, Clinici EM, 2010, Measures of dyspnea in pulmonary rehabilitation, *Multidisciplinary Respiratory Medicine*, vol. 5, no. 3, pp. 202-10.
- Enright PL, 2003, The Six-Minute Walk Test, *Respiratory Care*, vol. 48, no. 8, pp. 783-5.
- Harada S, Harada Y, Kitara Y, Takamoto M, Ishibashi T, Sininoda A, 1990, Tuberculosis Sequelae: Clinical Aspects, *Kekkaku journal*, vol. 65, no. 12, pp. 831-8.
- Kim HY, Song KS, Goo JM, Lee JS, Lee KS, Lim TH, 2001, Thoracic Sequelae and Complication of Tuberculosis, *RadioGraphics*, vol. 21, no. 4, pp. 839-60.



Launois C, Barbe C, Bertin C, Nardi J, Perotin J, Dury S, et al. 2012, The modified Medical Research Council scale for the assessment of dyspnea in daily living in obesity: a pilot study, *BMC Pulmonary Medicine*, vol. 12, no. 61, pp. 1-7.

Longo DL, Fauci AS, Kasper DL, Hauser SL, Jameson JL, Loscalzo J, 2012, *Harrison's Principles of Internal Medicine*, 18 edn, McGraw-Hill, United States of America.

Machida K, Maekura R, 2005. State of Art: Sequelae of Tuberculosis. *Kekkaku*, vol. 80, no. 10, pp. 665-74.

Marin JM, Carrizo SJ, Gascon M, Sanchez A, Gallego B, Celli BR, 2001, Inspiratory Capacity, Dynamic Hyperinflation, Breathlessness, and Exercise Performance during the 6-Minute-Walk Test in Chronic Obstructive Pulmonary Disease, *American Journal of Respiratory and Critical Care Medicine*, vol. 163, pp. 1395-9.

Naso FC, Pereira JS, Schuh SJ, Unis G, 2011, Functional evaluation in patients with pulmonary tuberculosis sequelae, *Pneumologia Portuguese Journal of Pulmonology*, vol. 39, pp. 1-6.

Pasipanodya JG, Miller TL, Vecino M, Munguia G, Garmon R, Bae S, Drewyer G, Weis SE, 2007, Pulmonary Impairment After Tuberculosis, *CHEST*, vol. 131, no. 6, pp. 1817-24.

Pusat Data dan Informasi Kementerian Kesehatan Republik Indonesia, 2011, Prevalensi TB per 100.000 Penduduk Provinsi D.I. Yogyakarta.

Stenton C, 2008, The MRC breathlessness scale, *Occupatioal Medicine*, vol. 58, pp. 226-7.



**HUBUNGAN ANTARA DERAJAT SESAK NAFAS DENGAN KAPASITAS FUNGSIONAL PADA PASIEN
SEKUELLE TUBERKULOSIS
PARU**

EKO NUGROHO, dr. Heni Retnowulan, M. Kes., Sp.PD; dr. Deddy Nur Wachid A., M. Kes., Sp. PD-KR
Universitas Gadjah Mada, 2015 | Diunduh dari <http://etd.repository.ugm.ac.id/>

World Health Organization, 1994, *TB: A Global Emergency*, WHO, Geneva.

World Health Organization, 2009, *Epidemiology. Global Tuberculosis Control: Epidemiology, Strategy, Financing*, WHO, Geneva.

World Health Organization, 2012, *Global Tuberculosis Report*, WHO, Geneva.

World Health Organization, 2012, *Tuberculosis Control in the South-East Asia Region*, WHO Regional Office for South-East Asia, New Delhi.

Yoshida N, Yoshiyama T, Asai E, Komatsu Y, Sugiyama Y, Mineta Y, 2006, Exercise Training for the Improvement of Exercise Performance of Patients with Pulmonary Tuberculosis Sequelae, *Internal Medicine*, vol. 45, no. 1505, pp. 399-403.