

CHAPTER VI

REFERENCES

- Amin SB. Clinical assessment of bilirubin-induced neurotoxicity in premature infants. *Semin Perinatol.* 2004; 28(5):340-347.
- American Speech-Language-Hearing Association, AHA <http://jslhr.pubs.asha.org/>. Accessed October 23, 2014.
- Bess FH, Dodd-Murphy J, Parker RA. Children with minimal sensorineural hearing loss: prevalence, educational performance, and functional status. *Ear Hearing* 1998; 19:339-54
- Burden V, Stott CM, Forge J, Goodyer I. The Cambridge Language and Speech Project (CLASP). I. Detection of language difficulties at 36 to 39 months. *Dev Med Child Neurol.* 1996;38(7):613-631.
- Bhandari V, Narang A, Mann SB, Raghunathan M, Bhakoo ON. Brain stem electrical response audiometry in neonates with hyperbilirubinaemia. *Indian J Paediatr* 1993;60:409-13.
- Billings KR, Kenna MA. Causes of pediatric sensorineural hearing loss: yesterday and today. *Arch Otolaryngol Head Neck Surg* 1999;125:517-21
- British Medical Journal (BMJ). <http://bestpractice.bmj.com/bestpractice/monograph/672/treatment/details.html>. Accessed 24 October 2014.
- Davis AC. The prevalence of hearing impairment and reported hearing disability among adults in Great Britain. *Int J Epidemiol* 1989; 18:911-17.
- Davis A, Wood S. The epidemiology of childhood hearing impairment: factors relevant to planning of services. *Br J Audiol* 1992; 26: 77-90.
- Ebbesen F, Andersson C, Verder H, et al. Extreme hyperbilirubinaemia in term and near-term infants in Denmark. *Acta Paediatrica.* 2005;94:59-64.
- Graziani LJ, Mitchell DG, Kornhauser M, Pidcock FS, Merton DA, Stanley C, et al. Neurodevelopment of preterm infants: neonatal neurosonographic and

- serum bilirubin studies. *Pediatrics*. 1992; 89(2):229-234.
- Hack M, Wilson-Costello D, Friedman H, Taylor GH, Schluchter M, Fanaroff AA. Neurodevelopment and predictors of outcomes of children with birth weights of less than 1000 g 1992-1995. *Arch Pediatr Adolesc Med*. 2000; 154(7):725-731.
- Hearing Conservation Amendment. In: Occupational Safety and Health Administration: occupational noise exposure federal register, part III. Washington DC: Department of Labor, 1981:37773-78.
- Hoing R. Inner ear changes in fetal asphyxia. *Laryngorhinotologie* 1991;70:72-7.
- Huang MJ, Kua KE, Teng HC, Tang KS, Weng HW, Huang CS. Risk factors for severe hyperbilirubinemia in neonates. *Pediatr Res*. Nov 2004;56(5):682-9.
- Koyama S, Kaga K, Sakata H, et al. Pathological findings in the temporal bone of newborn infants with neonatal asphyxia. *Acta Otolaryngol* 2005;125:1028-32
- Kumar RK. Neonatal jaundice. An update for family physicians. *Aust Fam Physician*. 1999;28:679-682.
- Law J, Garrett Z, Nye C. Speech and language therapy interventions for children with primary speech and language delay or disorder. *Cochrane Database Syst Rev*. 2003; (3):CD004110.
- Law J, Boyle J, Harris F, Harkness A, Nye C. Screening for primary speech and language delay: a systematic review of the literature. *Int J Lang Commun Disord*. 1998;33(Suppl):21-23.
- Leonard DR, Shen T, Howe HL, Egler T, eds. Trends in the prevalence of birth defects in Illinois and Chicago 1989 to 1997. *Epidemiologic report series 99:4*. Springfield, IL: Illinois Department of Public Health, 1999.
- Maisels MJ. What's in a name? Physiologic and pathologic jaundice: the conundrum of defining normal bilirubin levels in the newborn. *Pediatrics*. 2006;118:805-807.

- Manning D, Todd P, Maxwell M, et al. Prospective surveillance study of severe hyperbilirubinaemia in the newborn in the UK and Ireland. *Arch Dis Child Fetal Neonatal Ed.* 2007;92:F342-F346.
- Mohr PE, Feldman JJ, Dunbar JL, et al. The societal costs of severe to profound hearing loss in the United States. *Int J Technol Assess Health Care* 2000; 16: 1120-35.
- Nakamura, H., Takada, S., Shimabuku, R., Matsuo, M., Matsuo, T., & Negishi, H. (1985). Auditory nerve and brainstem responses in newborn infants with hyperbilirubinemia. *Pediatrics*; 75(4): 703-708.
- NHS National Institute for Health and Clinical Excellence. Neonatal Jaundice: A Clinical Guideline. <http://guidance.nice.org.uk/CG98/Guidance/pdf/English>
- Nelson HD, et al. Screening for speech and language delay in preschool children. Evidence synthesis no 41. Rockville MD: Agency for healthcare research and quality. 2006. <http://www.ahrq.gov/downloads/pub/prevent/pdf/ser/speechsyn.pdf>.
- Oh W, Tyson JE, Fanaroff AA, Vohr BR, Perritt R, Stoll BJ, et al. Association between peak serum bilirubin and neurodevelopmental outcomes in extremely low birth weight infants. *Pediatrics*. 2003; 112(4):773-779.
- Orchik DJ, Wark DJ. Hearing hazard of toy cellular telephones and walkie talkies. *Clin Pediatr* 1995; 34:278-80.
- Peltola H. Haemophilus influenzae type B disease and vaccination in Europe: lessons learned. *Pediatr Infect Dis* 1998;17(suppl):5126-32.
- Randall D, Reynell J, Curwen M. A study of language development in a sample of 3 year old children. *Br J Disord Commun.* 1974;9(1):3-16.
- Sasroasmoro S., Ismael S., 1995. *Dasar-dasar Metode Penelitian Klinis*. Jakarta : Binarupa Aksara.
- Shetty P. Speech and language delay in children: A review and the role of a pediatric dentist. *J Indian Soc Pedod Prev Dent* 2012;30:103-8

- Smitherman H, Stark AR, Bhutani VK. Early recognition of neonatal hyperbilirubinemia and its emergent management. *Semin Fetal Neonatal Med.* 2006;11:214-224.
- Smith RJH, Van Camp G. Nonsyndromic hereditary hearing loss and deafness, DFNB1. *GeneReviews at GeneTests: Medical Genetics Information Resource* <http://www.geneclinics.org/>
- Stern LM, et al. The Adelaide preschool language unit: results of follow-up. *J Paediatr Child Health.* 1995;31(3):207-212.
- Stierman L. Birth defects in California: 1983-1990. The California Birth Defects Monitoring Program, California Department of Health Services, 1994.
- Subcommittee on hyperbilirubinaemia, American Academy of Pediatrics. Management of hyperbilirubinaemia in the newborn infant 35 or more weeks of gestation, *Clinical Practice Guideline. Pediatrics* 2004, 114(1): 297-316.
- Sutton GJ, Rowe SJ. Risk factors for childhood sensorineural hearing loss in the Oxford Region. *Br J Audiol* 1997; 31: 39-54.
- White KR. Early hearing detection and intervention programs: opportunities for genetic services. *Am J Med Genet A* 2004; 130:29-36.
- Wellman MB, Sommer DD, McKenna J. Sensorineural hearing loss in postmeningitic children. *OtolNeurotol* 2003; 24:907-12.
- Wilson DH, Walsh PG, Sanchez L, et al. The epidemiology of hearing impairment in an Australian adult population. *Int J Epidemiol* 1999;28: 247-52.
- Zarina N, and Valeria N. Prevalence of sensorineural hearing loss on Asian children. 1996; 30(5): 332-339.