

### Daftar Pustaka

- Adam, C.L., Williams, P. a, Dalby, M.J., Garden, K., et al., 2014. Different Types of Soluble Fermentable Dietary Fiber Decrease Food Intake, Body Weight Gain and Adiposity in Young Adult Male Rats. *J Nutr Metab*, 11(1), pp.36.
- American Diabetes Association, 2014. Standards of medical care in diabetes 2014. *Diabetes Care*. 37 Suppl 1, S14-S80.
- Aminov, R.I., Walker, A.W., Duncan, S.H., Harmsen, H.J.M., et al., 2006. Molecular Diversity, Cultivation, and Improved Detection by Fluorescent in situ Hybridization of a Dominant Group of Human Gut Bacteria Related to *Roseburia* spp. or *Eubacterium rectale*. *J Appl Environ Microbiol*, 72(9), pp.6371-6376.
- Barcenilla, A., Pryde, S.E., Martin, J.C., Duncan, H., et al., 2000. Phylogenetic Relationships of Butyrate-Producing Bacteria from the Human Gut Phylogenetic Relationships of Butyrate-Producing Bacteria from the Human Gut. *J Appl Environ Microbiol*, 66(4), pp.1654-1661.
- Behall, K.M., Scholfield, D.J., Hallfrisch, J.G. & Liljeberg-Elmstahl, H.G.M., 2006 Consumption of Both Resistant Starch and  $\beta$ -Glucan Improves Postprandial Plasma Glucose and Insulin in Women. *Diabetes Res Care*, 29(5), pp.976-981.
- Besten, G., van Eunen, K., Groen, A.K., Venema, K., et al., 2013. The Role of Short-Chain Fatty Acids in the Interplay between Diet, Gut Microbiota, and Host Energy Metabolism. *J Lipid Res*, 54(9), pp.2325-2340.
- Bodinham, C.L., Smith, L., Thomas, E.L., Bell, J.D., et al., 2014. Efficacy of Increased Resistant Starch Consumption in Human type 2 Diabetes. *Endocr Connect*, 3(2), pp.75-84.
- Campbell, J.E. & Drucker, D.J., 2013. Pharmacology, Physiology, and Mechanisms of Incretin Hormone Action. *Cell Metab*, 17(6), pp.819-837.

- Cani, P.D., Daubioul, C.A., Reusens, B., Remacle, C., et al., 2005. Involvement of Endogenous Glucagon-like Peptide-1 (7-36) Amide on Glycaemia-lowering Effect of Oligofructose in Streptozotocin-treated Rats. *J Endocrinol Diabetes*, 185(3), pp.457-465.
- Cani, P.D., Hoste, S., Guiot, Y. & Delzenne, N.M., 2007. Dietary non-Digestible Carbohydrates Promote L-cell Differentiation in the Proximal Colon of Rats. *Br J Nutr*, 98(1), pp.32-37.
- Drucker, D.J., 1998. Perspectives in Diabetes: Glucagon-Like Peptides. *Diabetes*, 47(21), pp.159-169.
- Duncan, S.H. & Flint, H.J., 2008. Proposal of a Neotype Strain (A1-86) for *Eubacterium rectale*. *Int J Syst Evol Microbiol*, 58(7), pp.1735-1736.
- Edfalk, S. & Edlund, H., 2008. Gpr40 Is Expressed in Enteroendocrine Cells and Mediates Free Fatty Acid Stimulation of Incretin Secretion. *Diabetes*, 57(9), pp.2280-2287.
- Everard, A. & Cani, P.D., 2014. Gut microbiota and GLP-1. *Rev Endocr Metab Disord*, 15(3), pp.189-196.
- Fowler, M.J., 2008. Microvascular and Macrovascular Complications of Diabetes. *Clin Diabetes*, 26(2), pp.77-82.
- Fuentes-Zaragoza, E., Riquelme-Navarrete, M.J., Sánchez-Zapata, E. & Pérez-Álvarez, J.A., 2010. Resistant Starch as Functional Ingredient: A review. *Food Res Int*, 43(4), pp.931-942.
- Fujii, H., Iwase, M., Ohkuma, T., Ogata-kaizu, S., et al., 2013. Impact of Dietary Fiber Intake on Glycemic control, Cardiovascular Risk factors and Chronic Kidney Disease in Japanese Patients with Type 2 Diabetes Mellitus: the Fukuoka Diabetes Registry. *Nutr J*, 12, pp.159-167.
- Hindy, G., Sonestedt, E., Ericson, U., Jing, X., 2012. Role of TCF7L2 Risk Variant and Dietary Fibre Intake on Incident Type 2 Diabetes. *Diabetologia*, 55, pp.2646-2654.

- Holst, J.J., 2007. The physiology of glucagon-like peptide 1. *Physiol Rev*, 87(4), pp.1409-1439.
- Iswani, S., Suhartono, M.T., 2011, *Short Chain Fatty Acid Profile During in vitro Fermentation Of EuBacterium Rectale On Suku Sweet Potato Type-III Resistant Starch Media*. Institut Pertanian Bogor.
- Kaku, K., 2010. Pathophysiology of Type 2 Diabetes and Its Treatment Policy. *Japan Med Assoc J*, 53(1), pp.41-46.
- Kimura, I., Inoue, D., Hirano, K., Tsujimoto, G., 2014. The SCFA Receptor GPR43 and Energy Metabolism. *Front Endocrinol*, 5(85), pp.1-3.
- Kuwahara, A., 2014. Contributions of Colonic Short-Chain Fatty Acid Receptors in Energy Homeostasis. *Front Endocrinol*, 5 (144), pp.1-5.
- Lattimer, J.M. & Haub, M.D., 2010. Effects of Dietary Fiber and Its Components on Metabolic Health. *Nutrients*, 2(12), pp.1266-1289.
- Li, S.K., 2013. *Role of Vesicle-Associated Membrane Protein 2 in Glucagon-like Peptide-1 Secretion by Peptide-1 Secretion*. University of Toronto.
- Lin, H.V., Frassetto, A., Kowalik, E.J., Nawrocki, A.R., et al., 2012. Butyrate and Propionate Protect Against Diet-induced Obesity and Regulate Gut Hormones via Free Fatty Acid Receptor 3-Independent Mechanisms. *PloS one*, 7(4), pp.1-9.
- Litwak, L., Goh, S.Y., Hussein, Z., Malek, R., et al., 2013. Prevalence of Diabetes Complications in People With Type 2 Diabetes Mellitus and Its Association with Baseline Characteristics in the Multinational Achieve study. *Diabetol Metab Syndr*, 5(1), pp.57-66.
- Louis, P. & Flint, H.J., 2009. Diversity, Metabolism and Microbial Ecology of Butyrate-Producing Bacteria from the Human Large Intestine. *FEMS Microbiol Ecol*, 294(1), pp.1-8.
- Lu, S., Yu, Y., Zhu, H., Liu, X.D., et al., 2009. Berberine Promotes Glucagon-like Peptide-1 (7-36)

Amide Secretion in Streptozotocin-induced Diabetic Rats. *J Endocrinol*, 200(2), pp.159-165.

Mansour, A., Hosseini, S., Larijani, B., Pajouhi, M., et al., 2013. Nutrients Related to GLP1 Secretory Responses. *Nutrition*, 29(6), pp.813-820.

Massimino, S.P., Mcburney, M.I., Field, C.J., Thomson, A.B.R., et al., 1998. Nutrient Requirements and Interactions Fermentable Dietary Fiber Increases GLP-1 Secretion and Improves Capacity in Healthy Dogs. *J Endocrinol*, 128, pp.1786-1793.

Maziarz, M.P., 2013. Role of Fructans and Resistant Starch in Diabetes Care. *Diabetes Spectr*, 26(1), pp.2635-39.

Mitra, A., Bhattacharya, D. & Roy, S., 2007. Role of Resistant Starches Particularly Rice Containing Resistant Starches in Type 2 Diabetes. *J Hum Ecol*, 21(1), pp.47-51.

Nøhr, M.K., Pedersen, M.H., Gille, A., Egerod, K.L., et al., 2013. GPR41/FFAR3 and GPR43/FFAR2 as Cosensors for Short-chain Fatty Acids in Enteroendocrine Cells vs FFAR3 in Enteric Neurons and FFAR2 in Enteric Leukocytes. *Endocrinology*, 154(10), pp.3552-3564.

Olokoba, A.B., Obateru, O.A. & Olokoba, L.B., 2012. Type 2 Diabetes Mellitus: a Review of Current Trends. *Oman Med J*, 27(4), pp.269-273.

Ozougwu, O., 2013. The Pathogenesis and Pathophysiology of Type 1 and Type 2 Diabetes Mellitus. *J Physiol Pathophysiol*. 4(4), pp.46-57.

Petersen, N., Reimann, F., Bartfeld, S., Farin, H.F., et al., 2014. Generation of L cells in Mouse and Human Small Intestine Organoids. *Diabetes*. 63(2), pp.410-420.

Post, R.E., Mainous, A.G., King, D.E. & Simpson, K.N., 2012. Dietary Fiber for the Treatment of Type 2 Diabetes Mellitus: a Meta-Analysis. *J Am Board Fam Med*, 25(1), pp.16-23.

Psichas, A., Sleeth, M.L., Murphy, K.G., Brooks, L., et al., 2014. The Short Chain Fatty Acid Propionate

Stimulates GLP-1 and PYY Secretion via Free Fatty Acid Receptor 2 in Rodents. *Int J Obes*, pp.1-6.

Puddu, A., Sanguineti, R., Montecucco, F., Viviani, G.L., 2014. Evidence for the Gut Microbiota Short-Chain Fatty Acids as Key Pathophysiological Molecules Improving Diabetes. *Mediators Inflamm*, 2014, pp.1-9.

Riccardi, G. & Rivellese, A.A., 1991. Effects of Dietary Fiber and Carbohydrate on Glucose and Lipoprotein Metabolism in Diabetic Patients. *Diabetes Care*, 14(12), pp.1115-1125.

Schwiertz, A., Lehmann, U., Jacobasch, G. & Blaut, M., 2002. Influence of Resistant Starch on the SCFA Production and Cell Counts of Butyrate-producing Eubacterium spp. in the Human Intestine. *J Appl Microbiol*, 93 pp.157-162.

Setiawan, T.A., 2013. *Pengaruh Pemberian Tepung Gembili (Dioscorea esculenta) Terhadap Glukosa Darah Tikus Wistar yang Diinduksi Nikotinamida dan Streptozotosin*. Universitas Gadjah Mada.

Shao, W., Wang, D., Chiang, Y., Ip, W., et al., 2013. The Wnt Signaling Pathway Effector TCF7L2 Controls Gut and Brain Proglucagon Gene Expression and Glucose Homeostasis. *Diabetes*, 62(3), pp.789-800.

Thévenod, F., 2008. Pathophysiology of Diabetes Mellitus Type 2: Roles of Obesity, Insulin Resistance and  $\beta$ -Cell Dysfunction. *Front Diabetes*, pp.191-196.

Tolhurst, G., Heffron, H., Lam, Y.S., Parker, H.E., et al., 2012. Short-chain Fatty Acids Stimulate Glucagon-like Peptide 1 Secretion via the G-protein-coupled Receptor FFAR2. *Diabetes*, 61(2), pp.364-371.

Tolhurst, G., Reimann, F., Gribble, F.M., 2009. Nutritional Regulation of Glucagon-like Peptide1 Secretion. *J Physiol*, 587(1), pp.27-32.

Trinh, D.K.Y., Zhang, K., Hossain, M., Brubaker, P.L., et al., 2003. Expression in the Rodent Gastrointestinal Epithelium. *Diabetes*, 52(22), pp.425-433.

- Tuomilehto, J., Schwarz, P., Lindström, J., 2011. Long-term Benefits from Lifestyle Interventions for Type 2 Diabetes Prevention. *Diabetes Care*, 34(2), pp.210-214.
- Weickert, M.O., Pfeiffer, A.F.H., 2008. Metabolic Effects of Dietary Fiber Consumption and Prevention of Diabetes 1. *J Nutr*, 138, pp.439-442.
- Weickert, M.O., Arafat, A.M., Blaut, M., Alpert, C., et al., 2011. Changes in Dominant Groups of the Gut Microbiota do not Explain Cereal-fiber Induced Improvement of Whole-body Insulin sensitivity. *Nutr Metab*, 8, pp.1-10.
- Yu, Y., Liu, L., Wang, X., Liu, X., et al., 2010. Modulation of Glucagon-like Peptide-1 Release by Berberine: in vivo and in vitro Studies. *Biochem Pharmacol*, 79, pp.1000-1006
- Zeng, H., Lazarova, D.L., Bordonaro, M., 2014. Mechanisms Linking Dietary Fiber, Gut Microbiota and Colon Cancer Prevention. *World J Gastrointest Oncol*, 6(2), pp.41-51.
- Zhou, J., Hegsted, M., Mccutcheon, K.L., Keenan, M.J., et al., 2006. Peptide YY and Proglucagon mRNA Expression Patterns and Regulation in the Gut. *Obesity*, 14(4), pp.683-689.
- Zhou, J., Martin, R.J., Tulley, R.T., Raggio, A.M., et al., 2008. Dietary Resistant Starch Upregulates Total GLP-1 and PYY in a Sustained Day-Long Manner Through Fermentation in Rodents. *Am J Physiol Endocrinol Metab*, 295(5), pp.1160-1166.