



DAFTAR PUSTAKA

- Cannon, C.P. & Braunwald, E., 2010. Unstable angina and non-ST-elevation myocardial infarction. In J. Loscalzo, ed 17th. *Harrison's Cardiovascular Medicine*. New York: McGraw Hill, pp. 387-394.
- Enkhmaa, B., Anuurad, E., Zhang, Z., Pearson, T.A., Berglund, L., 2011. Usefulness of apolipoprotein B/apolipoprotein A-I ratio to predict coronary artery disease independent of the metabolic syndrome in African-Americans. *American Journal of Cardiology*, vol. 106, no. 9, pp. 1264-1269.
- Fodor, G., 2011. Primary prevention of CVD: treating dyslipidaemia. *British Medical Journal*, vol. 83, no. 10, pp. 1207-1208.
- Ghazali, M.V., 2011. Studi cross sectional. In S. Sastroasmoro & S. Ismael, eds. *Dasar-dasar Metodologi Penelitian Klinis*. Jakarta: Sagung Seto, pp. 131-132.
- Go, A.S., Mozaffarian, D., Roger, V.L., et al., 2013. Heart disease and stroke statistics--2013 update: a report from the American Heart Association. *Circulation*, vol. 127, no. 1, pp. e6-e245.
- Hamm, C.W., Heeschen, C., Falk, E., Fox, K.A.A., 2009. Acute coronary syndrome: pathophysiology, diagnosis, and risk stratification. In A. J. Camm, T. F. Luscher, & P. W. Serruys, . *The ESC Textbook of Cardiovascular Medicine*. New York: Oxford University Press, pp. 333-366.
- Hozo, S.P., Djulbegovic, B. & Hozo, I., 2005. Estimating the mean and variance from the median, range, and the size of a sample. *BMC Medical Research Methodology*, vol. 10, pp.1-10.
- Kim, M.C., Kini, A.S. & Fuster, V., 2007. Definitions of acute coronary syndromes. In V. Fuster, V., O'Rourke, R.A., Walsh, R.A., Poole-Wilson, P., *Hurt's The Heart*. New York: McGraw Hill, pp. 1311-1319.
- Kimura, E., 2011. Indonesia in 2010. *Asian Survey*, vol. 51, no.1, pp.186-195.



Krintus, M., Bergmann, K., Sypniewska, G., Sawicki, M., 2010. Comparison of apolipoprotein concentrations and values of apoB : apoAI with traditional lipid measures in women diagnosed with acute coronary syndromes - a preliminary report. *Journal of the International Federation of Clinical Chemistry and Laboratory Medicine*, vol. 21, no.4, pp. 1-7.

Krintus, M., Kozinski, M., Stefanska, A., et al., 2012. Value of C-Reactive Protein as a Risk Factor for Acute Coronary Syndrome: A Comparison with Apolipoprotein Concentrations and Lipid Profile. *Mediators of Inflammation*, vol. 2012, pp.1-10

Lamarche, B., Moorjani, S., Lupien, P.J., et al., 1996. Apolipoprotein A-I and B levels and the risk of ischemic heart disease during a five-year follow-up of men in the quebec cardiovascular study. *Circulation*, vol. 94, no.3 , pp.273-278.

Libby, P., 2010. The pathogenesis, prevention, and treatment of atherosclerosis. In J Loscalzo, ed. *2Harrison's Cardiovascular Medicine*. New York: McGraw Hill, pp. 322-328.

Lima, L.M., Carvalho, G. & Sousa, M.O., 2007. Clinical update apoB / apo A-I ratio and cardiovascular risk prediction. *Arquivos Brasileiros de Cardiologia*, vol. 88, no. 6, pp. e140-e143.

Mackay, J., Mensah, G.A., 2004. *The Atlas of Heart Disease and Stroke*, Geneva: WHO.

Manurung, D., 2006. Lipid profiles of acute coronary syndrome patients hospitalized in ICCU of Cipto Mangunkusumo Hospital. *Acta medica Indonesiana*, vol. 38, no.4, pp.196-20.

Marcovina, S., Packard, C.J., 2006. Measurement and meaning of apolipoprotein AI and apolipoprotein B plasma levels. *Journal of Internal Medicine*, vol. 259, no. 5, pp.437-46.

Maron, D.J., Ridker, P.M., Grundy, S.M., Pearson, T.A., 2008. Preventive strategy for coronary heart disease. In V. Fuster et al., eds. *Hurt's The Heart*. New York: McGraw Hill, pp. 1203-1234.



- Nagano, M., Horio, E., Sato, K., et al., 2014. Comprehensive analysis of intravascular ultrasound and angiographic morphology of culprit lesions between ST-segment elevation myocardial infarction and non-ST-segment elevation acute coronary syndrome. *International Journal of Cardiology*, vol. 171, no. 3, pp.423-430.
- Ninomiya, J.K., L'Italien, G., Criqui, M.H., Whyte, J.L., Gamst, A., Chen, R.S., 2004. Association of the metabolic syndrome with history of myocardial infarction and stroke in the third national health and nutrition examination survey. *Circulation*, vol. 109, pp.42-46.
- Schmidt, C., Wikstrand, J., 2009. High apoB / apoA-I ratio is associated with increased progression rate of carotid artery intima-media thickness in clinically healthy 58-year-old men: Experiences from very long-term follow-up in the AIR study. *Atherosclerosis*, vol. 205, pp.284-289.
- Scirica, B.M., Morrow, D. a, 2012. ST-Elevation myocardial infarction: pathology, pathophysiology, and clinical features. In D. L. Mann et al., *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*. Philadelphia: Elsevier Ireland Ltd, pp. 1068-1094.
- Singh, R., 2010. ApoB / apoAI ratio is statistically the best predictor of myocardial infarction compared to other lipid ratios. *International Journal of Pharma and Bio Sciences*. vol. 1, no. 2, pp.1-8.
- Suradi, R., Siahaan, C.M., Boedjang, R., Sudiyanto., Setyaningsih, I., Soedibjo, S., 2011. Studi kasus-kontrol. In Sudigdo Sastroasmoro & Sofyan Ismael, eds. *Dasar-dasar Metodologi Penelitian Klinis*. Jakarta: Sagung Seto, pp. 147-148.
- Sypniewska, G., Bergmann, K., Krintus, M., Kozinski, M., Kubica, J., 2011. How do apolipoproteins apoB and apoA-I perform in patients with acute coronary syndromes. *Journal of Medical Biochemistry*, vol. 30, no. 3, pp.237-243.



WHO, 2011. The top 10 cause of death. Available at:
<http://www.who.int/mediacentre/factsheets/fs310/en/index1.html> [Accessed April 30, 2014].

Walldius, G., Jungner, I., 2004. Apolipoprotein B and apolipoprotein A-I: risk indicators of coronary heart disease and targets for lipid-modifying therapy. *Journal of Internal Medicine*, vol. 255, no. 2, pp.188-205.

Walldius, G., Jungner, I., 2006. The apoB / apoA-I ratio: a strong , new risk factor for cardiovascular disease and a target for lipid-lowering therapy - a review of the evidence. *Journal of Internal Medicine*, vol. 259, pp.493-519.

Wallenfeldt, K., Bokemark, L., Wikstrand, J., Hulthe, J., Fagerberg, B., 2004. Apolipoprotein B/apolipoprotein A-I in relation to the metabolic syndrome and change in carotid artery intima-media thickness during 3 years in middle-aged men. *Journal of the American Heart Association*, vol. 35, pp.2248-2252.

Yusuf, S., Hawken, S., Ounpuu, S., et al.,2004. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet*, vol. 364, no. 9438 pp.937-52.