

## DAFTAR PUSTAKA

- Akulova, Z., 2012. File: *Tithonia diversifolia*.JPG, [cited 18 November 2014]. Available from: URL: [http://calphotos.berkeley.edu/cgi/img\\_query?enlarge=0000+0000+0312+0738](http://calphotos.berkeley.edu/cgi/img_query?enlarge=0000+0000+0312+0738)
- Arno, A.I., Gauglitz, G.G., Barret, J. P., Jeschke, M.G., 2014. Up-to-date approach to manage keloids and hypertrophic scars: A useful guide. *Burns: journal of the International Society for Burn Injuries*, pp.1-12.
- Bayat, A., Arscott, G., Ollier W.E.R., McGrouther D.A., Ferguson, M.W.J. 2005. Keloid disease: Clinical relevance of single versus multiple site scars. *British Association of Plastic Surgeons* 58:28-37.
- Chagas-paula, D.A., Oliveira, R.B., Silva, F.C., Gobbo-Neto, L., Gasparoto, T.H., Campanelli, A.P., Faccioli, L.H., Da Costa, F.B., 2011. Chlorogenic acids from *Tithonia diversifolia* demonstrate better anti-inflammatory effect than indomethacin and its sesquiterpene lactones. *Journal of Ethnopharmacology*, 136(2), pp.355-362.
- Fried, B, & Sherma, J (eds) 1999, *Thin-Layer Chromatography (4th Edition)*, CRC Press, New York, NY, USA.
- Gauglitz, G.G., Korting, H.C., Pavicic, T., Rucicka, T., Jeschke, M.G., 2011, Hypertrophic scarring and keloids: pathomechanisms and current and emerging treatment strategies. *Mol Med*;17(1-2):113-25.
- Goffin, E., Ziemons, E., De Mol, P., de Madureira Mdo, C., Martins, AP., da Cunha, AP., Philippe, G., Tits, M., Angenot, L., and Frederich, M., 2002, In vitro antiplasmodial activity of *Tithonia diversifolia* and identification of its main active constituent: Tagitinin C, *Planta Medica*; 68(6), 543-545.
- Graham-Brown, R., Burns, T., 2005. *Lecture Notes Dermatologi*. 8<sup>th</sup> ed. Jakarta: Erlangga.

- Gu, J. Q., Gills, J. J., Park, E. J., Mata-Greenwood, E., Hawthorne, M. E., Axelrod, F. Charez, P. I., Fong, H. H., Meththa, R. G., Pezzuto, J. M., Kinghor, J., 2002, Sesquiterpenes from *Tithonia diversifolia* with potential cancer chemopreventive activity. *J Nat Prod*; 65: 532 -36.
- Harting, M., Hicks, M.J., Levy, M.L. 2008. Dermal hypertrophies. Dalam: Wolff K, Goldsmith LA, Katz SI, Gilchrest BA, Paller AS, Leffel DJ, editors. *Fitzpatrick's Dermatology in General Medicine*. 7thed. New York: The McGraw-Hill Companies, 553-4.
- Hochman, B., Locali R.F., Matsuoka, P.K., Ferreira, L.M. 2008. Intralesional Triamcinolone Acetonide for Keloid Treatment:A Systematic Review. *Aesth Plast Surg* 32:705-9.
- Huang, L., Cai, Y.J., Lung, I., Leung, B.C.S., Burd, A., 2013. A study of the combination of triamcinolone and 5-fluorouracil in modulating keloid fibroblasts in vitro\*. *British Journal of Plastic Surgery*, 66(9), pp.e251-e259.
- Hutapea, J.R. 1994. *Inventaris Tanaman Obat Indonesia III. Peneliti Dan Pengembangan Kesehatan R.I.*, Jakarta, pp297-8.
- Jackson, I.T., Bhageshpur, R., DiNick, V., Khan, A., Bhaloo, S., 2001. Investigation of recurrence rates among earlobe keloids utilizing various postoperative therapeutic modalities. *Eur J Plast Surg*. 24(2):88-95.
- Keira, S.M., Ferreira, L.M., Gagnani, A., Duarte, I.S., Santos, I.A., 2004. Experimental model for fibroblast culture 1. *Acta Cir Bras*; 19, pp.11-16.
- Kelly, A.P. 1991. *Keloids and Hipertrophic Scars*. 1<sup>st</sup> ed. New York: McGraw-Hill Inc.
- Liao, M., Lin, W.C., Wen, H.C., Pu, H.F., 2011. *Tithonia diversifolia* and its main active component tagitinin C induce survivin inhibition and G2 / M arrest in human malignant glioblastoma cells. *Fitoterapia*, 82(3), pp.331-341.

- Lin, H.-R., 2012. Sesquiterpene lactones from *Tithonia diversifolia* act as peroxisome proliferator-activated receptor agonists. *Bioorganic & medicinal chemistry letters*, 22(8), pp.2954-8.
- Madureira, M.C., Martins, A.P., Gomes, M., Paiva, J., Cunha, A.P., Rosário, V., 2002. Antimalarial activity of medicinal plants used in traditional medicine in S. Tomé and Príncipe islands. *J. Ethnopharmacol.*, v.81, n.1, p.23- 29.
- Makinde, J.O., 2005. Influence of water extract of Mexican sunflower (*Tithonia diversifolia*) on growth of cowpea (*Vigna unguiculata*)., 4(April), pp.355-360.
- Monica, P.W., 2012. Efek Sitotoksik Ekstrak Etanol Terstandar *T. diversifolia* (Hemsley) A. Gray Terhadap sel Widr dibandingkan 5-Fluorouracil [Skripsi]. Fakultas Kedokteran Universitas Gadjah Mada, Yogyakarta.
- Nabin, C., Baruah., Ram P, Sharma., Madhusudanan, K.P., and Gopalakrishna, T., 1979, Sesquiterpene lactone of *Tithonia diversifolia*. Stereochemistry of the Tagitinins and related compounds. *J.Org.Chem*; 44(11):1831.
- Niessen, F.B., Schalkwijk, J., Vos, H., Timens, W., 2004. Hypertrophic scar formation is associated with an increased number of epidermal Langerhans cells. *J Pathol*;202:121-9.
- Ranti, I., 2014. Efek Antifibrotik Isolat Tagitinin C dari Daun Kembang Bulan [*Tithonia diversifolia* (Hemsley) A. Gray] pada Fibroblas Keloid [Tesis]. Fakultas Kedokteran Universitas Gadjah Mada, Yogyakarta.
- Renawati, E., 2011. Sitotoksitas Ekstrak Etanol Aloe vera Terhadap Sel Fibroblas Sebagai Bahan Medikamen Saluran Akar Secara In Vitro [Skripsi]. Fakultas Kedokteran Gigi Universitas Sumatera Utara, Medan.
- Rittie, L., Fisher, G.J., 2005, Isolation and Culture of Skin Fibroblast dalam J.Varga, D.A.

- Brenner, S.H. Phan (eds): *Fibrosis Research: Methods and Protocols*, Humana Press, Totowa, p. 84.
- Robles, D.T., Berg, D. 2007. Abnormal wound healing: keloids. *Clinics in Dermatology* 25:26-32.
- Saetung, A., Itharat, A., Dechsukum, C., Wattanapiromaskul, C., Keawpradub, N., Ratanasuwan, P., 2005. Cytotoxic activity of Thai medical plants for cancer treatment. *Songklanakarin J. Sci. Technol*, 27 (2): 469-478.
- Sanagi, M.M., 2001. Teknik Pemisahan Dalam Analisis Kimia. Skudai, Malaysia, Penerbit Universiti Teknologi Malaysia (UTM).
- Sharma BK. 2007. *Chromatography*. 5<sup>th</sup> ed. Meerut: GOEL Publishing House.
- Stockert, J.C., Blázquez-Castro, A., Cañete, M., Horobin, R.W., 2012. MTT assay for cell viability: Intracellular localization of the formazan product is in lipid droplets. *Acta histochemica*, 114(8), pp.785-96.
- Takashima, A., 1998, *Current protocol in cell biology : establishment of fibroblast cultures*. Dallas : John Wiley & Sons, Inc.
- Voigt, R., 1995, *Buku Pelajaran Teknologi Farmasi*, Diterjemahkan oleh Soendani N. S., UGM Press, Yogyakarta.
- Wahyuningsih, M.S.H., Sisindari, Murti, Y.B., Sudiby, R.S., 2009, *Potensi herbal terpilih sebagai agen antikanker yang spesifik*. Yogyakarta: LPPM UGM.
- Watson, D.G. 2010. *Analisis Farmasi: Buku Ajar untuk Mahasiswa Farmasi dan Praktisi Kimia Farmasi*. Edisi kedua. Syarif, W.R., penerjemah; Hadinata, A.H., editor. Jakarta: EGC.
- Wijaya, M.A.S., 2012. Efek Ekstrak Etanol Daun Kembang Bulan [*Tithonia diversifolia* (Hemsley) A. Gray] Sebagai Pemacu Respon Proliferasi Limfosit Pada

Mencit [Skripsi]. Fakultas Kedokteran Universitas Gadjah Mada, Yogyakarta.

Yang, M.L.Y.T.C., 2013. Anti-human hepatoma Hep-G2 proliferative, apoptotic, and antimutagenic activity of tagitinin C from *Tithonia diversifolia* leaves., pp.98-106.

Zhao, G.-J., Xi, Z.-X., Chen, W.-S., Li, X., Sun, L., Sun, L.-N., 2012. Chemical constituents from *Tithonia diversifolia* and their chemotaxonomic significance. *Biochemical Systematics and Ecology*, 44, pp.250-254.