

## DAFTAR PUSTAKA

- Bent A.E., Cundiff G.W., Swift S.E., editors. *OSTEGARD'S Urogynecology and Pelvic Floor Dysfunction*. United States of America: Lippincott Williams & Wilkins, 2008.
- Cavkaytar S., Kokanali M.K., Baylas A., Topçu H.O., Laleli B., Taşçı Y. Postpartum urinary retention after vaginal delivery: Assessment of risk factors in a case-control study. *Jurnal of the Turkish-German Gynecological Association*, 2014; 15: 140-3.
- Chitra T.V., Seetha P. Child Birth, Pregnancy and Pelvic Floor Dysfunction. *The Journal of Obstetrics and Gynecology of India* (November–December 2011) 61(6):635–637, 636.
- Cunningham F.G., Leveno K.J., Hauth J.C., Gilstrap III L., Bloom S.L., Wenstrom K.D. *Williams Obstetric 22ND EDITION*. United States of America: McGRAW-HILL, 2005.
- D. Chris, W. Hugh N. *ABC of Urology Second Edition*. United Kingdom: Blackwell Publishing Ltd, 2006.
- DeCherney, A.H., Nathan L., Goodwin T.M., Laufer N., editors. *Current Diagnosis & Treatment Obstetric & Gynecology*, Tenth Edition. United States of America: The McGraw-Hill Companies Inc, 2007.
- Dowling-Castronovo A., Spiro E.L. *Urinary Incontinence Assessment in Older Adults: Part II – Established Urinary Incontinence*. New York University College of Nursing, 2013.
- Elçin B., Özlem G., İskender G., Osman G. The Effects of Stress Urinary Incontinence on the Quality of Life of Turkish Women in the Reproductive Age Group. *Turkish Jurnal of Medical Sciences*, 2012; 42 (5): 845-851.
- Findik R.B., Unluer A.N., Şahin E., Bozkurt Ö.F., Karakaya J., Unsal A. Urinary incontinence in women and its relation with pregnancy, mode of delivery, connective tissue disease and other factors. *Advances in Clinical and Experimental Medicine*, 2012; 21(2): 207-213.
- Foldspang A., Mommsen S., Djurhuus, Jens C. Prevalent urinary incontinence as a correlate of pregnancy, vaginal childbirth, and obstetric techniques, 1999; 89(2): 209-212.
- Fritel X., Fauconnier A., Levet C., Bénifla J. Stress urinary incontinence 4 years after the first delivery: aretrospective cohort survey. *Acta Obstetricia et Gynecologica Scandinavica*, 2004 Oct; 83(10): 941-945.

- Gartland D, Donath S, MacArthur C, Brown S. The onset, recurrence and associated obstetric risk factors for urinary incontinence in the first 18 months after a first birth: an Australian nulliparous cohort study. *BJOG*, 2012; 119: 1361–1369.
- Gibbs R.S., Karlan B.Y., Haney A.F., Nygaard I.E., editors. *Danforth's Obstetric and Gynecology*, 10th Edition. Lippincott Williams & Wilkins, 2008.
- Glazener C, Elders A, MacArthur C, Lancashire R, Herbison P, Hagen S, Dean N, Bain C, Toozs-Hobson P, Richardson K, McDonald A, McPherson G, Wilson D. Childbirth and prolapse: long-term associations with the symptoms and objective measurement of pelvic organ prolapse. *BJOG* 2013; 120: 161–168.
- Kementrian Kesehatan Republik Indonesia. *Buku Saku Pelayanan Kesehatan Ibu di Fasilitas Kesehatan Dasar dan Rujukan*. Jakarta: Dirjen Bina Gizi dan KIA, 2013.
- Kepenekci I, Keskinilic B, Akinsu F, et al. Prevalence of pelvic floor disorders in the female population and the impact of age, mode of delivery, and parity. *Dis. Colon Rectum*. 2011; 54:85–94.
- Kessel V.K., Reed S., Newton S., Meier A., Lentz G. The second stage of labor and stress urinary incontinence. *Am J Obstet Gynecol*. 2001; 184:1571-5.
- Legendre G., Ringa V., Panjo H., Zins M.F.X. Incidence and remission of urinary incontinence at midlife: a cohort study. *BJOG* 2015; 122: 816–823.
- Lemmack G.E., Zimmern P.E. Predictability of urodynamic findings based on the Urogenital Distress Inventory–6 questionnaire. *Urology* 1999; 54: 461-466.
- Lentz G, Lobo R, Gershenson D, Katz V. *Comprehensive Gynecology Sixth Edition*. United States: Mosby 2012.
- Memon H, Handa V.L. Pelvic floor disorders following vaginal or cesarean delivery. *Current Opinion in Obstetrics & Gynecology*, 2012; 24(5): 349–54.
- Oliveira E, Zuliani L.M.M, Ischava J, Albuquerque S.S.R, de Souza A.M.B, Barbosa C.P. Evaluation of Factors Related to the Occurrence of Female Urinary Incontinence, *Revista da Associacao Medica Brasileira*, 2010; 56(6): 688-690.
- Patel D.A, Xu X, Thomason A.D, Ransom S.B, Ivy J.S, DeLancey. Childbirth and pelvic floor dysfunction: An epidemiologic approach to the assessment of

prevention opportunities at delivery. *Am J Obstet Gynecol*, 2006 July; 195(1): 23–28.

Rortveit G, Hannestad YS. Association between mode of delivery and pelvic floor dysfunction. *Tidsskr Nor Lægeforen* 2014; 134: 1848–1852.

S. Budi Iman. Inkontinensia Urin pada Perempuan . *Majalah Kedokteran Indonesia*, 2008; 58(7) : 258-264.

Saadia, Zahira. Effect of Age, Educational Status, Parity and BMI on Development of Urinary Incontinence - a Cross Sectional Study in Saudi Population. *Mater Sociomed*, 2015 Aug; 27(4): 251-254.

Siroky, M. B., Oates, R. D., & Babayan, R. K. *Handbook of urology : diagnosis and therapy*, 2004.

Stephansson O, Sandström A, Petersson G, Wikström A-K, Cnattingius S. Prolonged second stage of labour, maternal infectious disease, urinary retention and other complications in the early postpartum period. *BJOG*, 2015.

Tanagho E.A., McAninch J.W., editors. *Smith General Urology*. United States of America: The McGraw-Hill Companies, Inc 2008.

Uebersax JS, Wyman JF, Shumaker SA, McClish DK, Fantl JA, Continence Program for Women Research Group. Short Forms to Assess Life Quality and Symptom Distress for Urinary Incontinence in Women: The Incontinence Impact Questionnaire and the Urogenital Distress Inventory. *Neurourology and Urodynamics*, 1995; 14: 131-39.