



## REFERENCES

- Alothman, Z.A., 2012, A Review: Fundamental Aspects of Silicate Mesoporous Materials, *Materials*, 5, 2874-2902.
- Anastas, P.T., Kirchhoff, M.M., and Williamson, T.C., 2001, Catalysis as a Foundational Pillar of Green Chemistry, *Appl. Catal. A: Gen*, 221, 3-13.
- Anovitz, L.M., and Cole, D.R., 2015, Characterization and Analysis of Porosity and Pore Structures, *Rev. Mineral. Geochem.*, 80, 61-164.
- Bergen, L.A.H., Roos, G., and Proft, F., 2014, From Thiol to Sulfonic Acid: Modelling the Oxidation Pathway Protein Thiols by Hydrogen Peroxide, *J. of Phys. Chem. A.*, 118, 5078-5084.
- Chen, S., Yokoi, T., Tang, C., Jang, L., Tatsumi, T., Chan J. C., and Cheng, S., 2011, Sulfonic Acid-functionalized Platelet SBA-15 Materials as Efficient Catalysts for Biodiesel Synthesis, *Green Chem.*, 13, 2920.
- Cseri, T., Bekassy, S., Figueras, F., Cseke, E., Menorval, L.C., and Dutartre, R., 1995, Characterization of Clay-Based K Catalysts and Their Application in Friedel-Crafts Alkylation of Aromatics, *Appl. Catal. A: Gen*, 132, 141-155.
- Dacquin, J., Cross, H.E., Brown, D.R., Duren, T., Williams, J.J., Lee, A., and Wilson, K., 2010, Interdependent Lateral Interactions, Hydrophobicity and Acid Strength and Their Influence on the Catalytic Activity of Nanoporous Sulfonic Acid Silicas, *Green Chem.*, 12, 1383-1391.
- Dufaud, D., and Davis, M.E., 2003, Design of Heterogeneous Catalyst via Multiple Active Site Positioning in Organic-Inorganic Hybrid Materials, *J. Am. Chem. Soc.*, 125, 9403-9413
- Das, B., Suneel, K., Vankateswarlu, K., and Ravikanth B., 2008, Sulfonic Acid Functionalized Silica: An Efficient Heterogeneous Catalyst for a Three-Component Synthesis of 1,4-Dihidropyridines under Solvent-Free Conditions, *Chem. Pharm. Bull.*, 56, 366-368.
- Dias, A.S., Pillinger, M., and Valente, A.A., 2004, Dehydration of Xylose into Furfural over Micro-mesoporous Sulfonic Acid Catalyst, *J. Catal.*, 229, 414-423.
- Du, J., Bingying, J., Xingming, K., Xiancheng, Z., and Qingxiang, X., 2002, Enhanced Hydrolysis of Carboxylic Acid Esters Catalyzed by Metallomicelles Made of Cu(II) and Zn(II) Complexes, *J. Colloid Interface Sci.*, 256, 428-434.
- Ferreira, R.V., Pereira, I.L.S., Cavalcante, L.C.D., Gamarra, L.F., Carneiro, S.M., Amaro, E., Fabris, J.D., Domingues, R.Z., and Andrade, A.L., 2009, Synthesis and Characterization of Silica-Coated Nanoparticles of Magnetite, *J. Hyperfine Interact.*, 195, 265-274.



- Goto, T., Sakamoto, Y., and Kamiya, Y., 2009,  $\text{Cs}_{2.5}\text{H}_{0.5}\text{PW}_{12}\text{O}_{40}$  Nanoparticles Fixed on Silica Encapsulating Magnetic Iron Oxide as a Magnetically Separable Water-tolerant Solid Acid, *Chem. Letters*, 38, 7, 736-737.
- Gupta, P., and Paul, S., 2014, Solid Acid: Green Alternatives for Acid Catalysis, *Catalysis Today*, 236, 153-10.
- Hattori, H., and Ono, Y., 2015, *Solid Acid Catalysis from Fundamental to Applications*, Pan Stanford Publishing, USA, 25-47.
- Hong, R., Li, J., Zhang, S., Li, H., Zeng, Y., Ding, J., and Wei, D., 2009, Preparation and Characterization of Silica Coated  $\text{Fe}_3\text{O}_4$  Nanoparticles Used as Precursor of Ferrofluids, *Appl. Surf. Sci.*, 255, 3485-3492.
- Horita, N., Yoshimune, M., Kamiya, Y., and Okuhara, T., 2005.  $\text{Cs}_{2.5}\text{H}_{0.5}\text{PW}_{12}\text{O}_{40}$  Bonded to an Amine Functionalized  $\text{SiO}_2$  as an Excellent Water-tolerant Solid Acid, *Chem. Letters*, 34, 10, 1376-1377.
- Hui, C., Shen, C., Yang, T., Bao, L., Tian, J., Ding, H., Li, C., and Gao, H.J., 2008, Large-Scale  $\text{Fe}_3\text{O}_4$  Nanoparticles Soluble in Water Synthesized by a Facile Method, *J. Phys. Chem. C.*, 112, 11336-11339.
- Inumaru, K., Ishihara, T., Kamiya, Y., Okuhara, T., and Yamanaka, S., 2007, Water-Tolerant, Highly Active Solid Acid Catalysts Composed of the Keggin-Type Polyoxometalate  $\text{H}_3\text{PW}_{12}\text{O}_{40}$  Immobilized in Hydrophobic Nanospace of Organomodified Mesoporous Silica, *Angew. Chem.*, 46, 7625-7628.
- Karimi, B., and Khalkhali, M., 2004, Solid Silica-Based Sulfonic Acid as an Efficient and Recoverable Interphase Catalyst for Selective Tetrahydropyranylation of Alcohols and Phenols, *J. Mol. Catal. A: Chem.*, 232, 113-117.
- Kimura, M., Nakato, T., and Okuhara, T., 1997, Water-Tolerant Solid Acid Catalysts of  $\text{Cs}_{2.5}\text{H}_{0.5}\text{PW}_{12}\text{O}_{40}$  for Hydrolysis of Esters in the Presence of Excess Water, *Appl. Catal A: Gen.*, 165, 227-240.
- Kirchhoff, M.M., 2005, Promoting Sustainability through Green Chemistry, *Conservation and Recycling*, 44, 237-243.
- Kirillov, S.A., 2009, Surface Area and Pore Volume of a System Particles as a Function of Their Size and Packing, *Micropor. and Mesopor. Mat.*, 122, 234-239.
- Lai, D., Deng, L., Li, J., Liao, B., Guo, Q., and Fu, Y., 2011, Hydrolysis of Cellulose into Glucose by Magnetic Solid Acid, *ChemSusChem*, 4, 55-58.
- Liu, S., Zhu, W., Shi, L., Liu, H., Liu, Y., Ni, Y., Li, L., Zhou, H., Xu, S., He, Y., and Liu, Z., 2015, Activity Enhancement of Nafion Resin: Vapor-Phase Carbonylation od Dimethoxymethane over Nafion-Silica Composite, *Appl. Catal. A: Gen.*, 497, 153-159.



- Liu, Y.H., Lin, H., and Mou, C., 2004, Direct Method for Surface Silyl Functionalization of Mesoporous Silica, *Langmuir*, 20, 3231-3239.
- Lopez, D.E., Goodwin, J.G., and Bruce, D.A., 2006, Transesterification of Tiacetin with Methanol on Nafion® Acid Resins, *J. Catal.*, 245, 381-391.
- Marquez, F., Campo, T., Cotto, M., Polanco, R., Roque, R., Fierro, P., Sanz, J.M., Elizalde, E., and Morant, C., 2011, Synthesis and Characterization of Monodisperse Magnetite Hollow Microspheres, *Nano Lett.*, 1, 25-32.
- Materne, T., 2004, *Organosilane Technology in Coating Applications: Review and Perspectives*, Dow Corning Corporation, USA.
- Naeimi, H., and Nazifi, Z.S., 2013, A Highly Efficient nano- $\text{Fe}_3\text{O}_4$  Encapsulated-Silica Particles Bearing Sulfonic Acid Groups as a Solid Acid Catalyst for Synthesis of 1,8-dioxo-octahydroxanthene Derivatives, *J. Nanopart. Res.*, 15, 2026.
- Naggar, A.Y., 2013, Thermal Analysis of the Modified and Unmodified Silica Gels to Estimate their Applicability as Stationary Phase in Gas Chromatography, *JETEAS*, 4, 144-148.
- Niknam, K., and Saberi, D., 2009, Preparation of Sulfuric Acid ([3-(3-silicapropyl)sulfanyl]propyl)ester: A New and Recycable Catalyst for the Formylation and Acetylation of Alcohols under Heterogeneous Conditions, *Appl. Catal. A: Gen*, 366, 220-225.
- Nuryono, N., Rosiati, N.M., Rusdiarso, B., Sakti, S.C.W., and Tanaka, S., 2014, Coating of Magnetite with Mercapto Modified Rice Hull Ash Silica in a One-Pot Process, *SpringerPlus*, 3, 515.
- Okuhara, T., 2002, Water-Tolerant Solid Acid Catalysts, *Chem. Rev.*, 102, 3641-3666.
- Sawant, D.P., Vinu, A., Mirajkar, S.P., Lefebvre, F., Ariga, K., Anandan, S., Mori, T., Nishimura, C., and Halligudi, S.B., 2007, Silicotungstic Acid/Zirconia Immobilized on SBA-15 for Esterifications, *J. Mol. Catal. A: Chem.*, 271, 46-56.
- Sheldon, R.A., and Downing, R.S., 1999, Heterogeneous Catalytic Transformations for Environmentally Friendly Production, *Appl. Catal A: Gen*, 189, 163-183.
- Simonsen M.E., Sonderby, C., Li, Z., Erik, G., and Sogaard, 2009, XPS and FT-IR Investigation of Silicate Polymers, *J. Mater. Sci.*, 44, 2079-2088.
- Singh, T.K., Jain, C.L., Sharma, S.K., and Singh, S.S., 1999, Preparation of Dispersed Silica by Hydrolysis of Tetraethyl Orthosilicate, *Indian J. Eng. & Mat. Sci.*, 6, 349-351.
- Tan, W.L., and Bakar, A., 2006, The Effect of Additives on The Size of  $\text{Fe}_3\text{O}_4$  Particles, *J. Phys. Sci.*, 17, 2, 37-50.



- Wilson, K., Lee, A.F., Macquarrie, D.J., and Clark, J.H., 2002, Structure and Reactivity of Sol-Gel Sulphonic Acid Silicas, *Appl. Catal. A: Gen.*, 228, 127-133.
- Wu, W., He, Q., and Jiang, C., 2008, Magnetic Iron Oxide Nanoparticles: Synthesis and Surface Functionalization Strategies, *Nanoscale Res. Lett.*, 3, 397-415.
- Yamashita, T., and Hayes, P., 2008, Analysis of XPS Spectra of  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$  Ions in Oxide Materials, *J. Appl. Sur. Sci.*, 254, 2441-2449.
- Yang, D., Hu, J., and Fu, S., 2009, Controlled Synthesis of Magnetite-Silica Nanocomposites via a Seeded Sol-Gel Approach, *J. Phys. Chem.*, 113, 7646-7651.
- Yang, Q., Kapoor, M.P., Shirokura, N., Ohashi, M., Inagaki, S., Kondo, J.N., and Domen, K., 2004, Ethane-bridged Hybrid Mesoporous Functionalized Organosilicas with Terminal Sulfonic Groups and Their Catalytic Applications, *J. Mater. Chem.*, 15, 666-673.