

REFERENCES

- Abe, R., 2011, Development of new system for photocatalytic water splitting into H₂ and O₂ under visible light irradiation, *Bull. Chem. Soc. Jpn.*, 84(10), 1000-1030.
- Abe, R., Shinmei, K., Koumura, N., Hara, K. and Ohtani, B., 2013, Visible-light-induced water splitting based on two-step photoexcitation between dye-sensitized layered niobate and tungsten oxide photocatalyst in the presence of a triiodide/iodide shuttle redox mediator, *J. Am. Chem. Soc.*, 135(45), 16872-16884.
- Ansari, M.O., Khan, M.M., Ansari, S.J. and Cho, M.H., 2015, Electrically conductive polyaniline sensitized defective-TiO₂ for improved visible light photocatalytic and photoelectrochemical performance: a synergetic effect, *New J. Chem.*, 39(11), 8381-8388.
- Bagheri, S., Julkapli, N.M. and Hamid, S.B.A., 2014, Titanium dioxide as a catalyst support in heterogeneous catalyst, *Sci. World J.*, 2014, 1-21.
- Barnola, J.M., Pimienta, P., Raynaud, D. and Korotkevich, Y.S., 1991, CO₂-climate relationship as deduce from the Vostok ice core: a re-examination base on a new measurement and on a re-evaluation of the air dating, *Tellus*, 43B, 2, 83-90.
- Caruntu, D., Caruntu, G., Chen, Y., O'Connor, C.J., Goloverda, G. and Kolesnichenko, V.L., 2004, Synthesis of variable-sized nanocrystals of Fe₃O₄ with high surface reactivity, 16(25), 5527-5534.
- Chen, C. and Chen, Y., 2005, Fe₃O₄/TiO₂ core/shell nanoparticles as affinity probes for analysis of phosphopeptides using TiO₂ surface-assisted laser desorption/ionization mass spectrometry, *Anal. Chem.*, 77(18), 5912-5919.
- Beydoun, D., Amal, R., Low, G. and McEvoy, S., 2002, Occurrence and prevention of photodissolution at the phase junction of magnetite and titanium dioxide, *J. Mol. Catal. A: Chem.*, 180(1-2), 193-200.
- Daude, N., Gout, C. and Jouanin, C., 1977, Electronic band structure of titanium dioxide, *Phys. Rev. B*, 15(6), 3229-3235.
- Dimitrijevic, N.M., Vijayan, B.K., Poluektov, O.G., Rajh, T., Gray, K.A., He, H. and Zapol, P., 2011, Role of water and carbonates in photocatalytic transformation of CO₂ to CH₄ on titania, *J. Am. Chem. Soc.*, 133, 3964-3971.

- Farimani, M.H.R., Shahtahmasebi, N., Roknabadi, M.R., Ghows, N., and Kazemi, A., 2014, Study of structural and magnetic properties of superparamagnetic Fe₃O₄/SiO₂ core-shell nanocomposites synthesized with hydrophilic citrate-modified Fe₃O₄ seeds via a sol-gel approach, *Physica E*, 53, 207-216.
- Fletcher, C., Jiang, Y. and Amal, R., 2015, Production of formic acid from CO₂ reduction by means of potassium borohydride at ambient conditions, *Chem. Eng. Sci.*, 137, 301-307.
- Friedlingstein, P., Andrew, R.M., Rogelj, J., Peters, G.P., Canadell, J.G., Knutti, R., Luderer, G., Raupach, M.R., Schaeffer, M., van Vuuren, D.P. and le Quere, C., 2014, Persistent growth of CO₂ emission and implication for reaching climate targets, *Nat. Geosci.*, 7, 2248, 709-715.
- Fu, X., Clark, L.A., Yang, Q. and Anderson, M.A., 1996, Enhanced photocatalytic performance of titania-based binary metal oxides: TiO₂/SiO₂ and TiO₂/ZrO₂, *Environ. Sci. Technol.*, 30(2), 647-653.
- Fu, Y., Sun, D., Chen, Y., Huang, R., Ding, Z., Fu, X. and Li, Z., 2012, An amine-functionalized titanium metal-organic framework photocatalyst with visible-light-induced activity for CO₂ reduction, *Angew. Chem.*, 124, 3420-3423.
- Fujishima, A. and Honda, K., 1972, Electrochemical photolysis of water at a semiconductor electrode, *Nature*, 238, 37-38.
- Gomes, C.D.N., Jacquet, O., Villiers, C., Thuery, P., Ephritikhine, M. and Cantat, T., 2012, A diagonal approach to chemical recycling of carbondioxide: Organocatalytic transformation for the reductive functionalization of CO₂, *Angew. Chem. Int. Ed.*, 51, 187-190.
- Habisreutinger, S.N., Schimdt-Mende, L., Stolarczyk, J.K., 2013, Photocatalytic reduction of CO₂ on TiO₂ and other semiconductors, *Angew. Chem. Int. Ed.*, 52(29), 7372-7408.
- Hariani, P.L., Faizal, M., Ridwan, Marsi and Setyabudidaya, D., 2013, Synthesis and properties of Fe₃O₄ nanoparticles by co-precipitation method to removal procion dye, *IJESD*, 4(3), 336-340.
- Hirayama, J., Kondo, H., Miura, Y., Abe, R. and Kamiya, Y., 2012, Highly effective photocatalytic system comprising semiconductor photocatalyst and supported bimetallic non-photocatalyst for selective reduction of nitrate to nitrogen in water, *Catal. Commun.*, 20, 99-102.

- Hoffmann, M.R., Martin, S.T., Choi, W. and Bahnemann, D.W., 1995, Environmental applications of semiconductor photocatalysis, *Chem. Rev.*, 95, 69-96.
- Ihara, T., Miyoshi, M., Triyama, Y., Marsumato, O. and Sugihara, S., 2003, Visible-light-active titanium dioxide photocatalyst realized by an oxygen-deficient structure and by nitrogen doping, *Appl. Catal. B*, 42(4), 403-409.
- Indermuhle, A., Monnin, E., Stauffer, B. and Stocker, T.F., 2000, Atmospheric CO₂ concentration from 60 to 20 kyr BP from the Taylor Dome ice core, Antarctica, *Geophys. Res. Lett.*, 27, 5, 735-738.
- International Energy Agency, 2016, *World Energy Outlook*, <http://www.worldenergyoutlook.org>.
- Khairy, M. and Zakaria, W., 2014, Effect of metal-doping of TiO₂ nanoparticles on their photocatalytic activities toward removal of organic dyes, *Egypt. J. Pet.*, 23(4), 419-426.
- Kim, K.D., Kim, S.S., Choa, Y. and Kim, H.T., 2007, Formation and surface modification of Fe₃O₄ nanoparticles by co-precipitation and sol-gel method, *J. Ind. Eng. Chem.*, 13(7), 1137-1141.
- Lee, S., Cho, I., Lee, J.H., Kim, D.H., Kim, J.Y., Shin, H., Lee, J., Jung, H.S., Park, N., Kim, K., Ko, M.J. and Hong, K.S., 2010, Two-step sol-gel method-based TiO₂ nanoparticles with uniform morphology and size for efficient photo-energy conversion devices, *Chem. Mater.*, 22(6), 1958-1965.
- Linsebigler, A.L., Lu, G. and Yates, J.T., 1995, Photocatalyst in TiO₂ surfaces: principles, mechanisms, and selected results *Chem. Rev.*, 95(3), 735-785.
- Liu, J., Xu, J., Che, R., Chen, H., Liu, M. and Liu, Z., 2013, Hierarchical Fe₃O₄/TiO₂ yolk-shell microsphere with enhanced microwave-adsorption property, *Chem. Eur. J.*, 19(21), 6746-6752.
- Liu, S., Zhang, H., Xu, Z., Zhong, H. and Jin, H., 2012, Nitrogen-doped carbon xerogel as high active oxygen reduction catalyst for direct methanol alkaline fuel cell, *Int. J. Hydrogen Energy*, 37, 19065-19072.
- Lovell, E.C., Scott, J. and Amal, R., 2015, Ni-SiO₂ catalysts for the Carbon Dioxide Reforming of Methane: Varying Support Properties by Flame Spray Pyrolysis, *Molecules*, 20, 4594-4609.
- McCann, D.M., Lesthaeghe, D., Kletnieks, P.W., Guenther, D.R., Hayman, M.J., Speybroeck, V.V., Waroquier, M. and Haw, J.F., 2008, A complete catalytic cycle for supramolecular methanol-to-olefins conversion by linking theory with experiment, *Angew. Chem. Int. Ed.*, 120, 5257-5260.

- Ma, S.S.K., Maeda, K., Abe, R. and Domen, K., 2012, Visible-light-driven nonsacrificial water oxidation over tungsten trioxide powder modified with two different cocatalyst, *Energy Environ. Sci.*, 5, 8390-8397.
- Mahmood, K., Swain, B.S. and Amassian, A., 2015, Highly efficient hybrid photovoltaics based on hyperbranched three-dimensional TiO₂ electron transporting materials, *Adv. Mater.*, 27(18), 2859-2865.
- Mital, G.S. and Manoj, T., 2011, A review of TiO₂ nanoparticles, *Chin. Sci. Bull.*, 56, 1639-1657.
- Melian-Cabrera, I., Granados, M.L. and Fierro, J.L.G., 2002, Reverse topotactic transformation of a Cu-Zn-Al catalyst during wet Pd impregnation: Relevance for the performance in methanol synthesis from CO₂/H₂ mixtures, *J. Catal.*, 210, 273-284.
- Nakazawa, T., Machida, T., Esumi, K., Tanaka, M., Fujii, Y., Aoki, S. and Watanabe, O., 1993, Measurements of CO₂ and CH₄ concentrations in air in a polar ice core, *J. Glaciol.*, 39, 132, 209-215.
- Olah, G.A., 2005, Beyond oil and gas: the methanol economy, *Angew. Chem. Int. Ed.*, 44, 2636-2639.
- Peterson, A.A. and Norskov, J.K., 2012, Activity descriptors for CO₂ electroreduction to methane on transition-metal catalysts, *J. Phys. Chem. Lett.*, 3(2), 251-258.
- Rakopoulos, C.D. and Kyritsis, D.C., 2001, Comparative second-law analysis of internal combustion engine operation for methane, methanol, and dodecane fuels, *Energy*, 26, 705-722.
- Richardson, I., Thomson, M., Infield, D. and Clifford, C., 2010, Domestic electricity use: A high-resolution energy demand model, *Energy Build.*, 42, 1878-1887.
- Sakai, N., Wang, R., Fujishima, A., Watanabe, T. and Hashimoto, K., 1998, Effect of ultrasonic treatment on highly hydrophilic TiO₂ surfaces, *Langmuir*, 14(20), 5918-5920.
- Sato, S., Morikawa, T., Kajino, T. and Ishitani, O., 2013, A highly efficient mononuclear Iridium complex photocatalyst for CO₂ reduction under visible light, *Angew. Chem.*, 125, 1022-1026.
- Takeda, H., Koike, K., Inoue, H. and Ishitani, O., 2008, Development of an efficient photocatalytic system for CO₂ reduction using Rhenium(I) Complexes based on mechanistic study, *J. Am. Chem. Soc.*, 130, 2023-2031.

- Tilgner, I.C., Fischer, P., Bohnen, F.M., Rehage, H. and Maier, W.F., 1995, Effect of acidic, basic and fluoride-catalyzed sol-gel transition on the preparation of sub-nanostructured silica, *Microporous Mater.*, 5(1-2), 77-90.
- Tomita, O., Otsubo, T., Higashi, M., Ohtani, B. and Abe, R., 2016, Partial oxidation of alcohols on visible-light-responsive WO₃ photocatalyst loaded with palladium oxide cocatalyst, *Catal.*, 6, 1134-1144.
- Vesselli, E., Rizzi, M., de Rogatis, L., Ding, X., Baraldi, A., Comelli, G., Savio, L., Rocca, M., Fornasiero, P., Baldereschi, A. and Peressi, M., 2010, Hydrogen-assisted transformation of CO₂ on Nickel: The role of Formate and Carbonate, *J. Phys. Chem. Lett.*, 1, 402-406.
- Vijayalakshmi, R. and Rajendran, V., 2012, Synthesis and characterization of nano-TiO₂ via different methods, *Arch. Appl. Sci. Res.*, 4(2), 1183-1190.
- Xuan, S., Jiang, W., Gong, X., Hu, Y. and Hen, Z., 2009, Magnetically separable Fe₃O₄/TiO₂ hollow spheres: fabrication and photocatalytic activity, *J. Phys. Chem. C*, 113(2), 553-558.
- Yu, J.C., Yu, J., Ho, W. and Zhang, L., 2001, preparation of highly photocatalytic active nano-sized TiO₂ particles via ultrasonic irradiation, *Chem. Commun.*, 19, 1942-1943.
- Zaleska, A., 2008, Doped-TiO₂: a review, *Recent Pat. Eng.*, 2(3), 157-164.
- Zhang, H. and Banfield, J.F., 2014, Structural characteristics and mechanical and thermodynamic properties of nanocrystalline TiO₂, *Chem. Rev.*, 114, 9613-9644.
- Zhang, S., Li, L., Zhao, S., Sun, Z., Hong, M. and Luo, J., 2015, Hierarchical metal-organic framework nanoflowers for effective CO₂ transformation driven by visible light, *J. Mater. Chem. A*, 3(30), 15764-15768.
- Zhao, Z. and Liu, Q., 2007, Mechanism of higher photocatalytic activity of anatase TiO₂ doped with nitrogen under visible-light irradiation from density functional theory calculation, *J. Phys. D: Appl. Phys.*, 41, 1-10.
- Zhu, C., Zhang, M., Qiao, Y., Xiao, G., Zhang, F. and Chen, Y., 2010, Fe₃O₄/TiO₂ core/shell nanotubes: synthesis and magnetic and electromagnetic wave adsorption characterization, *J. Phys. Chem. C*, 114(39), 16229-16235.