

## DAFTAR PUSTAKA

- Akbar, S., 2011, *Andrographis paniculata*: A Review of Pharmacological Activities and Clinical Effects, *Alternative Medicine Review*, Volume 16, Number 1, page 66-77
- Akowuah, G.A., Zhari, I., Norhayati, I., Sadikun, A., Khamsah, S.M., 2004, Sinensetin, eupatorin, 3'-hydroxy-5,6,7,4'-tetramethoxyflavone and rosmarinic acid contents and antioxidative effect of *Orthosiphon stamineus* from Malaysia, *Food Chemistry* 87, 559-566
- Armstrong, N.A., 2006, *Pharmaceutical Experimental Design and Interpretation*, Second Edition, 84-88, Taylor & Francis Group, LLC, London
- Badan POM RI, 2008, *Direktorat Obat Asli Indonesia*, Jakarta: Badan POM
- Baehr, H.D., dan Stephan, K., 2006, *Heat and Mass Transfer*, 2nd ed, 64-70, Springer, Berlin
- Balasundram N., Sundram, K., Samman, S., 2006, Phenolic compounds in plants and agri-industrial by-products: Antioxidant activity, occurrence, and potential uses, *Food Chemistry*, 191-203
- Bolton, S., dan Bon, C., 2004, *Pharmaceutical Statistic Practical ad Clinical Application 4th ed*, 326-353, 591-601, Marcel Dekker Inc., New York
- Chang, C.C., Yang, M.H., Wem, H.M., Chern, J.C., 2002, Estimation of Total Flavonoid Content in Propolis by Two Complimentary Colorimetric Methods, *Journal of Food and Drug Analysis*, Vol. 10, No.3
- Chao, W.W. dan Lin, B.F., 2010, Isolation and Identification of Bioactive Compounds in *Andrographis paniculata* (*Chuanxinlian*), *Chinese Medicine Journal*. 5:1-15.
- Chao, W.W., Kuo, Y. H., Lin, B. F., 2010, Anti-inflammantory Activity of New Compounds from *Andrographis paniculata* by NF-kB Transactivation Inhibition, *Journal of Agricultural and Food Chemistry*, 508, 2505-2512
- Chebil, L., Humeau, C., Anthoni, J., Dehez, F., Engasser, Jean-Marc, Ghoul, M., 2007, Solubility of Flavonoids in Organic Solvents, *Journal of Chemical and Engineering Data*, Vol. 52, No.5
- Cubas C., Lobo, M.G., Gonzales, M., 2008, Optimization of the extraction on chlorophylls in green beans (*Phaseolus vulgaris* L.) by N,N-dimethylforamide using response surface methodology, *Journal of Food Composition and Analysis*, Vol 21(2), 125-133
- Cujic, N., Savikin, K., Jankovic, T., Pljevljakusic, D., Zdunnic, G., Ibric, S., 2016, Optimization of polyphenols extraction from dried chokeberry using maceration as traditional technique, *Food Chemistry* 194, 135-142

- Departemen Kesehatan, 2000, *Parameter Standar Umum Ekstrak Tumbuhan Obat*, 14-20, Departemen Kesehatan Republik Indonesia, Direktorat Jenderal Pengawasan Obat dan Makanan, Jakarta
- Departemen Kesehatan, 2008, *Farmakope Herbal Indonesia*, Edisi 1, 159-162, Departemen Kesehatan Republik Indonesia, Jakarta
- Dhanani, T., Shah, S., Gajbhiye, N.A., Kumar, S., 2013, Effect of extraction methods on yield, phytochemical constituents and antioxidant activity of *Withania somnifera*, *Arabian Journal of Chemistry* (2017) 10, S1193-S1199
- Durling, N.E., Catchpole, O.J., Grey, J.B., Webby, R.F., Mitchell, K. A., Foo, L. Y., Perry, N.B., 2007, Extraction of phenolics and essential oil from dried sage (*Salvia officinalis*) using ethanol-water mixtures, *Food Chemistry*, 101, 1417-1424
- Everette, J.D., Bryant, Q.M., Green, A.M., Abbey, Y.A., Wangila, G.W., Walker, R.B., 2010, A thorough study of reactivity of various compound classes towards the Folin-Ciocalteu reagent, *J. Agric Food Chem.* 58(14):8139-8144
- Gandjar, I.G. dan Rohman A, 2007, *Kimia Farmasi Analisis*, Pustaka Pelajar, Yogyakarta
- Guo, H., Zhang, Z., Su., Z., Sun., C., Zhang, X., Zhaou, X., Lai., X., Su., Z., Li., Y., Zhan., J. Y., 2016, Enhance anti-tumor activity and reduce toxicity by combination andrographolide and bleomycin in ascitic tumor bearing mice, *European Journal of Pharmacology*, Vol 776, P. 52-63
- Handayani, H., Sriherfyna, F. H., Yunita, 2016, Ekstraksi Antioksidan Daun Sirsak Metode Ultrasonik Bath, *Jurnal pangan dan Agroindustri*, 4(1), 262-272
- Hossain, Md. S., Urbi, Z., Sule, A., Rahman, K. H. M., 2014, *Andrographis paniculata* (Burm. f.) Wall ex Nees: A review of Ethnobotany, Phytochemistry, and Pharmacology, *The scientific World Journal*, Vol 2014
- Huang, W.Y., Cai, Y.Z., & Zhang, Y., 2010, Natural Phenolic Compounds from Medicinal Herbs and Dietary Plants: Potential Use for Cancer Prevention, *Nurt Cancer* 62(1) : 1-20
- Jarukamjorn, K., dan Nemoto, N., 2008, Pharmacological aspects of *Andrographis paniculata* on health and its major diterpenoid constituent andrographolide, *Journal of health science*, 54: 370-381
- Katno, 2008, *Penanganan Pasca Panen Tanaman Obat*, Balai Besar Penelitian dan Pengembangan Tanaman Obat dan Obat Tradisional, Departemen Kesehatan Republik Indonesia, Tawangmangu.

- Kumar, A., J. Dora, A. Sigh, dan R. Tripathi, 2012, A Review on King of Bitter (Kalmegh), *International Journal of Research in Pharmacy and Chemistry*. Vol. 2 (1). P 116-124.
- Kumoro, A.C., dan Hasan, M., 2007, Supercritical carbon dioxide extraction of Andrographolide from *Andrographis paniculata*: Effect of the solvent flow rate, pressure, and temperature, *Chin. J. Chem. Eng.* 15(6) 877-883
- Kumoro, A.C., Hasan. M., Singh, H., 2009, Effect od solvent properties on the soxhlet extraction of diterpenoid lactones from *Andrographis paniculata* leaves, *ScienceAsia* 35: 306-309
- Li, W., Xu, X., Zhang, H., Ma, C., Fong, H., Van Breemen, R., Fitzloff, J., 2007, Secondary metaboltes from *Andrographis paniculata*, *Chem. Pharm. Bull.* 55(3) 455-458
- Luthria, D. L., 2008, Influence of experimental conditions on the extraction of phenolic compounds from parsley (*Petroselinum crispum*) flakes using a pressurized liquid extractor, *Journal of Agriculture and Food Chemistry*, 107, 745-752
- Mahendra, B., 2005, *13 Jenis Tanaman Obat Ampuh*, Penebar Swadaya, Jakarta
- Mandal, V., 2007, Microwave Assisted Extraction– An Innovation and Promising Extraction Tool For Medical Plant Research, *Pharmacognosy Reviews*, Vol. 1, Issue 1, cit. Diantika, F., Sutan, S. M., Yulianingsih, R., 2014, Pengaruh lama ekstraksi dan konsentrasi pelarut etanol terhadap ekstraksi antioksidan biji kakao, *Jurnal teknologi pertanian* Vol 15 No 3 159-164
- Markham, K. R., 1988, *Cara mengidentifikasi flavonoid*, diterjemahkan oleh: Kosasih Padmawinata, Penerbit ITB, Bandung
- Mongkolsilp, S., Pongbupakit, I., Sae-Lee, N., Sitthithaworn, W., 2004, Radical scavenging activity and total phenolic content of medicinal plants used in primary health care, *SWU J Pharm Sci*, Vol. 9 No. 1
- Muchlisah, Fauziah, 2001, *Tanaman Obat Keluarga (Toga)*, Penebar Swadaya. Jakarta
- Nijveldt, R. J., Nood, E. van, Hoorn, EC van, Boelens, P. G., Norren, K. van, Leeuwen, P. AM. van , 2001, Flavonoid: a review of probable mechanism of action and potential applications 1-3, *American Society for Clinical Nutrition*
- Niranjan, A., Tewari, SK., Lehri, A., 2010, Biological Activities of Kalmegh (*Andrographis paniculata* Nees) and its Active Principles-A review, *Indian Journal of Natural Products and Resources*, Vol. 1(2), pp. 125-135
- Nur, A. M., dan Astawan M., 2011, Kapasitas antioksidan bawang dayak (*Eleutherine palmifolia*) dalam bentuk segar, simplisia dan keripik, pada pelarut nonpolar, semipolar dan polar, *Skripsi*, Fakultas teknologi pertanian, Institut pertanian bogor

- Okhuarobo, A., Falodun J. E., Erharuyi, O., Imieje, V., Falodun, A., Langer, P., 2014, Harnessing the medicinal properties of *Andrographis paniculata* for diseases and beyond: a review of its phytochemistry and pharmacology, *Asian Pasific of Tropical Disease*, Vol. 4(3), pp. 213-222
- Pinelo, M., Del Fabbro, P., Manzocco, L., Nunez, M.J., Nicoli, M.C., 2005, Optimization of continuous phenol extraction from *Vitis vinifera* byproducts, *Food Chemistry* 92, 109-117
- Prapanza E. dan Marianto LM, 2003, *Khasiat & Manfaat Sambiloto: Raja Pahit Penakluk Aneka Penyakit*, 4-7, AgroMedia Pustaka.
- Prior, R.L., Wu, X., Schaich, K., 2005, Standardized methods for the determination of antioxidant capacity and phenolics in food and dietary supplement, *J. Agric. Food Chem.*, 53, 4290-4302
- Puspitasari, A., 2016, Optimasi ekstraksi dan penelusuran fraksi aktif herba sambiloto (*Andrographis paniculata* Nees.) dan patikan kebo (*Euphorbia hirta* L.) studi in vitro efek fagositosis makrofag, *Disertasi*, Universitas Gadjah Mada, Yogyakarta
- Quispe Candori, S., Foglio, M.A., Rosa, P.T.V., Meireles, M.A.A., 2008, Obtaining b-caryophyllene from *Cordia verbenacea* de Candolle by supercritical fluid extraction. *J. Supercrit. Fluids* 46, 27–32 cit. Dhanani, T., Shah, S., Gajbhiye, N.A., Kumar, S., 2013, Effect of extraction methods on yield, phytochemical constituents and antioxidant activity of *Withania somnifera*, *Arabian Journal of Chemistry* (2017) 10, S1193-S1199
- Rafi, M., Darusman, L.K., Nurashah, E.S., Syafitri, U.D., 2014, Optimization of extraction conditions for andrographolide using fractional factorial design, *Indonesian J. Pharm.* Vol 25 No. 3: 145-152
- Randhir, R., Lin, Y. T., Shetty, K., 2004, Phenolic, their antimicrobial activity in dark germinated fenugreek sprouts in response to peptide and phytochemical elicitors, *Asia Pasific Journal of Clinical Nutrition*, 13, 295-307
- Rao Y, Koteswara; Vimalamma G, Rao CV, Tzeng YM., 2004, Flavonoids and andrographolides from *Andrographis paniculata*. *Phytochemistry* 2004;65:2317-2321
- Robinson, T., 1995, *Kandungan Organik Tumbuhan Tinggi*, Terjemahan Kosasih Padmawinata, Edisi Keenam, ITB Press, Bandung
- Soeksmanto, A., Hapsari, Y., dan Simanjuntak, P., Kandungan antioksidan pada beberapa bagian tanaman mahkota dewa, *Phaleria macrocarpa* (Scheff) Boerl. (Thymelaceae), *Biodiversitas*, Vol. 8 No. 2 p. 92-95
- Spigno, G., Tramelli, L., Faveri, Dante Marco De, 2007, Effect of extracton time, temperature nd solvent on concentration and antioxidant activity of grape marc phenolics, *Journal of Food Engineering* 81 (2007) 200-208.

- Sultana, B., Anwar, F., dan Ashraf, M., 2009, Effect of extraction solven/technique on the antioxidant activity of selected medicinal plant extracts, *Molecules*, 14, 2167-2180
- Tan, O.W., Tan, C.P., dan Ho, C.W., 2011, Antioxidant properties: Effects of solid-to-solvent ratio on antioxidant compounds and capacities of Pegaga (*Centella asiatica*), *International Food Research Journal* 18: 557-562
- Thoo, Y.Y, Ng, S.Y., Khoo, M. Z., Aida, W. M. Wan., Ho, C. W., 2013, A Binary Solven Extraction System for Phenolic Antioxidant and its Application to the Estimation of Antioxidant Capacity in *Androgrpahis paniculata* extract, *International Food Research Journal*, Vol. 20, Issue 3, p1103-1111. 9p.
- Sin H. N., Yusof, S., Hamid, N. S. A. dan Rahman, R. A., 2006, Optimization of hot water extraction for sapodilla juice using response surface methodology, *Journal of Food Engineering* 74:352-358.
- Wijesekera, R.O.B., 1991, *Medicinal Plant Industry*, 85-98, CRC press, Florida, USA.
- Wong, B.Y., Tan, C.P., Ho, C.W., 2013, Effect of solid-to-solvent ratio on phenolic content and antioxidant capacities of “Dukung Anak” (*Phyllanthus niruri*), *International Food Research Journal* 20(1): 325-330
- Wongkittipong, R., Prat, L., Damronglerd, S., Gourdon, C., 2004, Solid-liquid extraction of andrographolide from plants-experimental study, kinetic reaction and model, *Separation and Purification Technology*, Vol 40 (2), Pages 147-154
- Wu, J., Yu, D., Sun, H., Zhang, Y., Zhang, W., Meng, F., Du, X., 2015, Optimizing the extraction of anti-tumor alkaloids from the stem of *Berberis amurensis* by response surface methodology, *Industrial Crops and Products*, Vol. 69, P. 68-75