

DAFTAR PUSTAKA

- ADA (2014) Diagnosis and classification of diabetes mellitus. *Diabetes Care*, 37(SUPPL.1), hal.81–90.
- Adiwijono dan Ahmad H. (1993) Dislipidemia pada diabetes mellitus tipe ii: patofisiologi dan pendekatan terapi. *Berkala Ilmu Kedokteran*, XXV(4), hal.189–201.
- Alauddin, M., Shirakawa, H., Koseki, T., Kijima, N., Ardiansyah, Budijanto, S., Islam, J., Goto, T. dan Komai, M. (2016) Fermented rice bran supplementation mitigates metabolic syndrome in stroke-prone spontaneously hypertensive rats. *BMC Complementary and Alternative Medicine*, 16(442), hal.1-11.
- Amalan, V., Vijayakumar, N. dan Indumathi, D. (2016) Antidiabetic and antihyperlipidemic activity of p -coumaric acid in diabetic rats , role of pancreatic GLUT 2 : In vivo approach. *Biomedicine dan Pharmacotherapy. Elsevier*, 84, hal.230–236.
- Astawan, M. dan Leomitro, D. (2009) *Khasiat Whole Grain: Makanan Berserat Untuk Hidup Sehat*. Jakarta: PT Gramedia Pustaka Utama
- Bahadoran, Z., Mirmiran, P. dan Azizi, F. (2013) Dietary polyphenols as potential nutraceuticals in management of diabetes : a review. *Journal of Diabetes & Metabolic Disorders*, 12(43), hal.1–9.
- Bantle, J. P. dan Slama, G. S. (2006) *Nutritional Management of Diabetes Mellitus and Dysmetabolic Syndrome*. Swiss: Nestlé Nutrition Institute, Karger.
- Burlando, B. dan Cornara, L. (2014) Therapeutic properties of rice constituents and derivatives (*Oryza sativa L.*): A review update. *Trends in Food Science dan Technology*, 40(1), hal.82–98.
- Camacho, P., Gharib, H., dan Sizemore, G. W. (2012) *Evidence-Based Endocrinology 3rd Edition*. Philadelphia: Lippincott Williams dan Wilkins.
- Chehade, J.M., Gladysz, M. dan Mooradian, A.D. (2013) Dyslipidemia in type 2 diabetes: prevalence, pathophysiology, and management. *Drugs*, 2013(73), hal.327–339.
- Chen, C-W. dan Cheng, H-H. (2006) A rice bran oil diet increases LDL-receptor and HMG-CoA reductase mRNA expressions and insulin sensitivity in rats with streptozotocin/nicotinamide-induced type 2 diabetes. *The Journal of nutrition*, 136(6), hal.1472–1476.
- Chen, Z., Ma, K.Y., Liang, Y., Peng, C., dan Zuo, Y. (2011) Role and classification of cholesterol-lowering functional. *Journal of Functional Foods*, 3(2), hal.61–

69. Tersedia di: <<http://dx.doi.org/10.1016/j.jff.2011.02.003>> [Diakses pada 4 September 2016].

Cheng, H-H., Ma, C-Y., Chou T-W., Chen Y-Y., dan Lai, M-H. (2010) Gamma-oryzanol ameliorates insulin resistance and hyperlipidemia in rats with streptozotocin/nicotinamide-induced type 2 diabetes. *International Journal for Vitamin and Nutrition Research*, 80(1), hal.45–53.

Chithra, P.K., Sindhu, G., Shalini, V., Parvathy, R., Jayalekshmy, A., dan Helen, A. (2015) Dietary Njavara rice bran oil reduces experimentally induced hypercholesterolaemia by regulating genes involved in lipid metabolism. *British Journal of Nutrition*, hal.1207–1219.

Chou, T-W., Ma, C-Y., Cheng, H-H., Chen, Y-Y., dan Lai, M-H. (2009) A rice bran oil diet improves lipid abnormalities and suppress hyperinsulinemic responses in rats with streptozotocin/nicotinamide-induced type 2 diabetes. *Journal of Clinical Biochemistry and Nutrition*, 45(1), hal.29–36. Tersedia di: <<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2704324&tool=pmcentrez&andrendertype=abstract>> [Diakses pada 7 Oktober 2016].

Cicero, A. F. G. dan Gaddi, A. (2001) Rice bran oil and γ -oryzanol in the treatment of hyperlipoproteinaemias and other conditions. *Phytotherapy Research*, 15(March), hal.277–289.

Cicero, A.F.G., Parini, A. dan Rosticci, M. (2015) Nutraceuticals and cholesterol-lowering action. *IJCME*, 6(2015), hal. 1–4.

Daou, C. dan Zhang, H. (2014) Functional and physiological properties of total , soluble , and insoluble dietary fibres derived from defatted rice bran. *J Food Sci Technol*, 51(12), hal.3878–3885.

Doddamani, S., Nikam, S., Nikam, P., dan Dambal, A. (2014.) Activity of lecithin cholesterol acyl transferase and apolipoprotein A-i in newly detected type 2 diabetes mellitus. , 4(14), hal.2653–2660.

Farbstein, D. dan Levy, A.P. (2012) HDL dysfunction in diabetes: causes and possible treatments. *Expert Rev Cardiovasc Ther*, 10(3): 353–61

Faria, S.A.S.C., Bassinello, P.Z., dan Penteado, M.V.C. (2012) Nutritional composition of rice bran submitted to different stabilization procedures. *Brazilian Journal of Pharmaceutical Sciences*, 48, (4), 651-7.

Federer, W.T. (1977) *Experimental Design Theory and Application*, 3rd Ed.. New Delhi: Oxford and IBH Publishing Co. Pvt. Ltd

Fernandez, M. L. dan Webb, D. (2008) The LDL to HDL cholesterol ratio as a valuable tool to evaluate coronary heart disease risk. *J Am Coll Nutr*, 27(1), hal. 1–5.

Garg, N., Agrawal, Y.B., dan Gupta, S. (2014) A study of lipid profile levels in diabetics and non-diabetics taking TC / HDL ratio and LDL / HDL ratio into

consideration. *Journal Indian Academy of Clinical Medicine*, 15(3), hal. 192–195.

Ghatak, S.B. dan Panchal, S.J. (2011) Anti-hyperlipidemic activity of oryzanol, isolated from crude rice bran oil, on Triton WR-1339-induced acute hyperlipidemia in rats. *Brazilian Journal of Pharmacognosy*, 7, hal.10-20.

Handayani, D., Am, A., Soeadmadji, D.W., dan Widodo, M.A. (2003) Enzim lipoprotein lipase suatu alternatif pemeriksaan gangguan metabolisme lemak pada penderita DM tipe 2 in vitro. XIX(2).

Havekes, L.M., Jukema, J.W., Rensen, P.C.N., Kastelein, J.J., Romijn, J.A., Kuipers, F., Princen, H.M.G., dan Van Eck, M. (2009) *Modulation of HDL metabolism*. Thesis, Universiteit Leiden.

Hongu, N., Kitts, D.D, Zawistowski, J., Dossett, C.M., Kope, A., Pope, B.T., dan Buchowski, M.S. (2014) Pigmented rice bran and plant sterol combination reduces serum lipids in overweight and obese adults. *J Am Coll Nutr*, 33(3), hal.231–238.

Hu, G. dan Yu, W. (2013) Binding of cholesterol and bile acid to hemicelluloses from rice bran. *International Journal of Food Sciences and Nutrition*, 64(4), hal.461–566.

International Food Policy Research Institute (2016) *Global Nutrition Report 2016: From Promise to Impact: Ending Malnutrition by 2030*.

Ishii, S., Kamegai, J., Tamura, H., Shimizu, T., Sugihara, H., dan Oikawa, S. (2002) Role of ghrelin in streptozotocin-induced diabetic hyperphagia', *Endocrinology*, 143(12), hal. 4934–4937.

Jung, E.H., Kim, S.R., Hwang, I.K., dan Ha, T.Y. (2007) Hypoglycemic effects of a phenolic acid fraction of rice bran and ferulic acid in C57BL / KsJ- db / db mice. *J. Agric. Food Chem*, 55(24), hal.9800–9804.

Justo, M.L., Rodriguez, R., Claro, C. M., Alvarez, M., Parrado, J., dan Herrera, M.D. (2012) Water-soluble rice bran enzymatic extract attenuates dyslipidemia , hypertension and insulin resistance in obese Zucker rats. *European Journal of Nutrition*.

Kemenkes: Pusat Data dan Informasi (2014) *Situasi dan Analisis Diabetes*.

Kraemer, F.B. dan Shen, W-J. (2006) Hormone-sensitive lipase knockouts. *Nutrition & Metabolism*, 3(12), hal.1–7.

Krentz, A.J. (2003) Lipoprotein abnormalities and their consequences for patients with Type 2 diabetes. *Diabetes, Obesity and Metabolism*, 5, hal.19–27.

Kupski, L., Silvello, M.A.D.C., Fontes, M.R.V., Liima, T.S., Treichel, H., dan Furlong, E.B. (2015) *R . oryzae* cellulases : a new approach to degrading lignocellulosic material. *Journal of Food Biochemistry*. 39(2), hal.1–10.

- Lerma-García, M.J., Herrero-Martínez, J.M., Simó-Alfonso, E.F., Mendonça, C.R.B., dan Ramis-Ramos, G. (2009) Composition, industrial processing and applications of rice bran γ -oryzanol. *Food Chemistry*, 115(2), hal.389–404. Tersedia di: <http://dx.doi.org/10.1016/j.foodchem.2009.01.063>.
- Martins, S., Mussatto, S.I., Martínez-Avila, G., Montañez-Saenz, J., Aguilar, C. N., dan Teixeira, J.A. (2011) Bioactive phenolic compounds: Production and extraction by solid-state fermentation: A review. *Biotechnology Advances*, 29(3), hal.365–373. Tersedia di: <http://dx.doi.org/10.1016/j.biotechadv.2011.01.008> [Diakses pada 9 Oktober 2016].
- Masiello, P., Broca, C., Gross, R., Roye, M., Manteghetti, M., Hillarire-Buys, D., Novelli, M., dan Ribes, G (1998) Development of a new model in adult rats administered streptozotocin and nicotinamide. *Diabetes*, 47(February), hal.224–229.
- Mirmiran, P., Bahadoran, Z. dan Azizi, F. (2014). Functional foods-based diet as a novel dietary approach for management of type 2 diabetes and its complications : A review. *World Journal of Diabetes*, 5(3), hal.267–281.
- Moongngarm, A., Daomukda, N. dan Khumpika, S. (2012) Chemical compositions , phytochemicals , and antioxidant capacity of rice bran , rice bran layer , and rice germ. *APCBEE Procedia*, 2, hal.73–79. Tersedia di: <http://dx.doi.org/10.1016/j.apcbee.2012.06.014> [Diakses pada 14 September 2016].
- Mulyono, A.M.W. (2010) Peran miselium fungi dalam ransum ayam kaya non-starch polysaccharides: suatu kajian pustaka. *Prosiding Seminar Nasional Hari Lingkungan Hidup Se-Dunia 2010*.
- Murray, R.K. Garnner, D.K., dan Rodwell, W. (2009). *Biokimia Harper 27th Ed.* Jakarta: EGC.
- Nair, A. B. dan Jacob, S. (2016) A simple practice guide for dose conversion between animals and human. *Journal of Basic and Clinical Pharmacy*, 7, hal. 27–31.
- NIH (2016) *Explore High Blood Cholesterol*. Tersedia di: <http://www.nhlbi.nih.gov> [Diakses pada 25 November 2016].
- Nimmanapalli, H.D., Kasi, A.D., Devapatla, P., dan Nuttacki, V. (2016) Lipid ratios , atherogenic coefficient and atherogenic index of plasma as parameters in assessing cardiovascular risk in type 2 diabetes mellitus. *International Journal of Research in Medical Sciences*, 4(7), hal. 2863–2869.
- Nursalim, Y. dan Razali, Z. Y. (2007) *Bekatul: Makanan yang Menyehatkan*. Jakarta: PT Agro Media Pustaka
- Oduguwa, O.O., Edema, M.O. dan Ayeni, A.O. (2008) Physico-chemical and microbiological analyses of fermented corn cob, rice bran and cowpea husk

- for use in composite rabbit feed. *Bioresource Technology*, 99(6), hal.1816–1820.
- Ogbera, A.O., Fasanmade, O.A., Chinenye, S., dan Akinlade, A. (2009) Characterization of lipid parameters in diabetes mellitus – a Nigerian report. *International Archives of Medicine*, 7, hal.1–7.
- Ohara, I., Tabuchi, R., dan Onai, K. (2000) Effects of modified rice bran on serum lipids and taste preference in streptozotocin-induced diabetic rats. *Nutrition Research*, 20(1), hal.59–68.
- Okon, U.A., Owo, D.U., Udokang, N.E., Udobang, J.A., dan Ekpenyong, C.E. (2012) Oral administration of aqueous leaf extract of *ocimum gratissimum* ameliorates polyphagia, polydipsia and weight loss in streptozotocin-induced diabetic rats. *American Journal of Medicine and Medical Sciences*, 2(3), hal. 45–49.
- Ozougwu, J., Obimba, K., Belonwu, C., dan Unakalamba, C. (2013) The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *Journal of Physiology and Pathophysiology*, 4(4), hal. 46–57.
- Pandya, H., Lakhani, J.D., Dadhania, J., dan Trivedi, A. (2012) The prevalence and pattern of dyslipidemia among type 2 diabetic patients at rural based hospital. *Indian Journal of Clinical Practice*, 22(12), hal.36–44.
- Pereira, M.A. dan Ludwig, D.S. (2001) Dietary fiber and body-weight regulation: Observations and mechanisms. *Pediatric Clinics of North America*, 48(4), hal. 969–980.
- PERKENI (2015) *Konsensus Pengelolaan dan Pencegahan Diabetes Mellitus Tipe 2*. Jakarta: Perkumpulan Endrokinologi Indonesia.
- Qureshi, A.A., Sami, S.A. dan Khan, F.A. (2002) Effects of stabilized rice bran, its soluble and fiber fractions on blood glucose levels and serum lipid parameters in humans with diabetes mellitus Types I and II. *Journal of Nutritional Biochemistry*, 13, hal.175–187.
- Randhir R. dan Shetty K. (2007) Mung beans processed by solid-state bioconversion improves phenolic content and functionality relevant for diabetes and ulcer management. *Innovations Food Sci Emerg Technol*, 8:197–204.
- Risma, W., Prapti, N.K.G. dan Artawan, I.P. (2014) Pengaruh kombinasi pengaturan pola diet dan senam diabetes terhadap kadar gula darah pada pasien DM tipe II.
- Schwartz, S.L. (2006) Diabetes and dyslipidaemia. *Diabetes, Obesity and Metabolism*, 8, hal.355–364.
- Sengupta, P. (2013) The laboratory rat: relating its age with human's. *International Journal of Preventive Medicine*, 4 (6). hal. 624-630.

- Shao, Y., Xu, F., Sun, X., Bao, J., dan Beta, T. (2014) Identification and quantification of phenolic acids and anthocyanins as antioxidants in bran, embryo and endosperm of white, red and black rice kernels (*Oryza sativa* L.). *Journal of Cereal Science*, 59(2), hal.211–218. Tersedia di: <<http://dx.doi.org/10.1016/j.jcs.2014.01.004>> [Diakses pada 3 Maret 2016].
- Siddiqui, S., Ahsan, H., Rashid, M., dan Siddiqui, W.A. (2013) Protective effects of tocotrienols against lipid-induced nephropathy in experimental type-2 diabetic rats by modulation in TGF- β expression. *Toxicology and Applied Pharmacology*, 273(2), hal.314–324. Tersedia di: <<http://dx.doi.org/10.1016/j.taap.2013.09.004>> [Diakses pada 14 September 2016].
- Sohail, M., Rakha, A., Butt, M., dan Jawad, M. (2016) Rice bran nutraceuticals : A comprehensive review. *Critical Reviews in Food Science and Nutrition*, Taylor & Francis, 8398(March).
- Starzynska-Janiszewska, A., Stodolak, B., dan Jamróz, M. (2008) Antioxidant properties of extracts from fermented and cooked seeds of Polish cultivars of *Lathyrus sativus*. *Food Chemistry*, 109, hal.285–292.
- Sumantha, A., Deepa, P., Sandhya, C., Szakacs, G., Soccol, C.R., dan Pandey, A. (2006) Rice bran as a substrate for proteolytic enzyme production. *Brazilian Archives of Biology and Technology*, 49(5), hal.843–851.
- Tripathi, V. dan Verma, J. (2014) Different models used to induce diabetes: A comprehensive review. *International Journal of Pharmacy and Pharmaceutical Sciences*, 6(6), hal.29–32.
- Tuminah, S. (2009) Peran kolesterol HDL terhadap penyakit kardiovaskuler dan diabetes mellitus. *Gizi Indonesia*, 32(1), hal.69–76.
- Van Linthout, S., Spillmann, F., Schultheiss, H-P., dan Tschöpe, C. (2010) High-density lipoprotein at the interface of type 2 diabetes mellitus and cardiovascular disorders. *Current Pharmaceutical Design*, 16(13), hal.1504–1516. Tersedia di: <https://www.researchgate.net/publication/41668561_High-Density_Lipoprotein_at_the_Interface_of_Type_2_Diabetes_Mellitus_And_Cardiovascular_Disorders> [Diakses pada 8 September 2016].
- Widowati, W. (2008) Potensi antioksidan sebagai antidiabetes. *JKM*, 7(2), hal.1–11.
- Wang, Z.Q., Zuberi, A., Zhang, X.H., Macgowan, J., Qin, J., Ye, X., Son, L., Wu, Q., Lian, K., dan Cefalu, W.T. (2009) Effects of dietary fibers on weight gain, carbohydrate metabolism and gastric ghrelin gene expression in high fat fed mice. *Metabolism*, 56(12), hal. 1635–1642.
- WHO (2016a) *Global Report on Diabetes*. Tersedia di:<http://apps.who.int/iris/bitstream/10665/204871/1/9789241565257_eng.pdf> [Diakses pada 12 September 2016].

WHO (2016b) *Diabetes Mellitus*. Tersedia di:
<<http://www.who.int/ediacentre/factsheets/>> [Diakses pada 3 Maret 2016].

Winter, W.E. dan Signorino, M.R. (2002). *Diabetes Mellitus: Pathophysiology, Etiologies, Complications, Management, and Laboratory Evaluation*. Washington DC: AACC Press.

Xu, Z., Hua, N., dan Godber, J.S. (2001) Antioxidant activity of tocopherols, tocotrienols, and gamma-oryzanol components from rice bran against cholesterol oxidation accelerated by 2,2'-azobis(2-methylpropionamide) dihydrochloride. *J. Agric. Food Chem.*, 49(4), hal.2077-2081

Zhang, H., Jiang, S., Hong, X., dan Huang, J. (2015) Chapter 3: The role of hormones in glucose homeostasis and diabetes treatment. *Recent Advances in Diabetes Treatment*.