

## REFERENCES

- Abu-Ghannam, N. & McKenna, B. (1997). The Application of Peleg's Equation to Model Water Absorption During The Soaking of Red kidney Beans (*Phaseolus vulgaris* L.). *Journal of Food Engineering*, 32, 391-401.
- Anonym<sup>a</sup>.(2016). Life Cycle of a Silkworm.Retrieved March 17<sup>th</sup>, 2017 from <https://everythingsilkworms.com.au/silkworms/life-cycle-of-a-silkworm/>.
- AOAC.(2002). Official Analytical Methods. Washington, DC: Association of Official Analytical Chemists.
- Aramwit, P., Kanokpanont, S., Nakphaeng, T., & Srichana, T. (2010).The Effect of Sericin from Various Extraction Methods on Cell Viability and Collagen Production. *International Journal of Molecular Sciences*, 11, 2200-2211.
- Aramwit, P., Siritientong, T., & Srichana, T. (2012). Potential Application of Silk Sericin, A Natural Protein From Textile Industry by-products. *Waste Management & Research*, 30(3), 217-224.
- Aregbesola, O. A., Ogunsina, B. S., Sofolahan, A. E., & Chime, N.N. (2015). Mathematical Modeling of Thin Layer Drying Characteristics of dika (*Irvingia gabonensis*) Nuts and Kernels. *Nigerian Food Journal*, 33, 83-89.
- Atmosoedarjo, H. S., Katsubrata, J., Kaomini, M., Saleh, W.& Moerdoko, W. (2000). *Sutera Alam Indonesia*. Jakarta: SaranaWana Jaya.
- Bilbao-Sáinz, C., Andres, A. & Fito, P. (2005). Hydration Kinetics of Dried Apples As Affected by Drying Conditions. *Journal of Food Engineering*, 68, 369-376.
- Cacace, J. E. & Mazza, G. (2003). Mass Transfer Process During Extraction of Phenolic Compounds From Milled Berries. *Journal of Food Engineering*, 59 (4), 379-389.
- Chen, X., Lam, K. F., Mak, S. F. & Yeung, K. L. (2011).Precious Metal Recovery by Selective Adsorption Using Biosorbents. *Journal of Hazard Mater*, 186, 902-910.
- Chen, F., Porter, D., & Vollrath, F. (2012).Structure and Physical Properties of Silkworm Cocoons. *Journal of The Royal Society Interface*, 9, 2299-2308.
- Chunningham, S. E., McMinn, W. A. M., Magee, T. R. A., & Richardson, P. S. (2007). Modeling Water Absorption of Pasta During Soaking. *Journal of Food Engineering*, 84, 214-223.

- Cohen, E. & Saguy, L. (1985). Statistical Evaluation of Arrhenius Model and Its Applicability in Prediction of Food Quality Losses. *Journal of Food Processing and Preservation*, 9 (4), 273-290.
- Gupta, D., Agrawal, A., & Rangil, A. (2014). Extraction and Characterization of Silk Sericin. *Indian Journal of Fibre & Textile Research*. 39, pp. 364-372.
- Henderson, S. M. & Pebis, S. (1961). Grain Drying Theory I. Temperature Effect on Drying Coefficient. *Journal of Agricultural Engineering Research*, 6, 169-174.
- Heslot, H. (1998). Artificial Fibrous Proteins: A Review. *Biochimie*, 80, 19-31.
- Hizaji, A. S., Maghsoudlou, Y., & Jafari, S. M. (2010). Application of Peleg Model to Study Effect of Water Temperature and Storage Time on Rehydration Kinetics of Air Dried Potato Cubes. *Latin American Applied Research*, 40, 131-136.
- Inserco<sup>a</sup>. (2013). Statistic: Global Silk Industry. Retrieved March 17<sup>th</sup>, 2017 from <http://inserco.org/en/statistics>.
- Inserco<sup>b</sup>. (2013). Thailand: History. Retrieved March 17<sup>th</sup>, 2017 from <http://inserco.org/en/thailand>.
- Joseph, J. & Raj, S. J. (2012). Therapeutic Applications and Properties of Silk Proteins From *Bombyx mori*. *Frontiers in Life Science*, 6(3-4), 55-60.
- Jumaeri. (1997). *Pengetahuan Barang Tekstil*. Bandung: ITT.
- Kodama, K. (1926). The Preparation and Pshyco-chemical Properties of Sericin. *Biochem*. 20(06), 1208-1222.
- Kurioka, A., Kurioka, F., & Yamazaki, M. (2004). Characterization of Sericin Powder Prepared from Citric Acid-degraded Sericin Polypeptides of the Silkworm, *Bombyx Mori*. *Biosci. Biotechnol. Biochem.*, 68(4), 774-780.
- Ma, M., Hussain, M., Dong, S., & Zhou, W. (2016). Characterization of The Pigment In Naturally Yellow-Colored Domestic Silk. *Journal of Dyes and Pigments*, 124, 6-11.
- Maharaj, V., & Sankat, C. K. (2000). The Rehydration Characteristics and Quality of Dehydrated Dasheen Leaves. *Canadian Agricultural Engineering*, 42(2), 81-85.
- Masahiro, S., Hideyuki, Y. & Norihisa, K. (2000). Consumption of Silk Protein, Sericin Elevates Intestinal Absorption of Zinc, Iron, Magnesium and Calcium in Rats. *Nutr. Res*, 20, 1505-1511.
- Mc Intyre J.E., Daniel, P.N. (1997). *Textile Terms and Definitions 10<sup>th</sup> Edition*. England : The Textile Institute. pp 93.

- Medhekar, A. (2015). Lifecycle of Silkworm (With Diagram). Retrieved March 17<sup>th</sup>, 2017 from <http://www.biologydiscussion.com/animals-2/lifecycle-of-silkworm-with-diagram>.
- Minora, N., & Tsukada, M. (1991). Oxygen permeable membranes, Jpn Kokai Tokkyo Koho Jap 02233128 A2 (to Agency of Ind Aci & Tech, Japan; Ministry of agriculture & Forestry sericulture, Japan). Chem Abstr, 114(6), 45871.
- Moreira, R., Chenlo, F., Chaguri, L. & Fernandes, C. (2008). Water Absorption, Texture, and Color Kinetics of Air-dried Chestnuts During Rehydration. Journal of Food Engineering, 86, 584-594.
- Moss, J. R. & Otten, L. (1989). A Relationship Between Colour Development and Moisture Content During Roasting of Peanuts. Canadian Institute Food Science and Technology Journal, 22, 34-39.
- Mossel, B., Bhandari, B., Arcy, D. & Caffin, N. (2000). Use of An Arrhenius Model To Predict Rheological Behavior in Some Australian Honeys. LWT- Food Science and Technology, 33 (8), 545-552.
- Nikitakis, J., & Breslawee, H. P. (2014). International Cosmetic Ingredient Dictionary and Handbook. 14 ed. Washington DC: Personal Care Products Council.
- Nindhia, T. G. T., Surata, I. W., Knejzlik, Z., Ruml, T., Nindhia, T. S. (2015). New Route in Degumming of Bombyx Mory Silkworm Cocoon for Biomaterial. Journal of Medical and Bioengineering, 4(4), 338-341.
- Padamwar, M. N. & Pawar, A. P. (2004). Silk Sericin and its applications: A review. Journal of Scientific & Industrial Research, 63, 323-329.
- Padamwar, M. N., Pawar, A. P., Daithankar, A. V., & Mahadik, K. R. (2005). Silk Sericin as a Moisturizer : an In Vivo Study. Journal of Cosmetic Dermatology. 4(4), 250-257.
- Panthong, J., Eamchotchawalit, C., & Songsermpong, S. (2015). Effect of Spray Drying Conditions on the Characteristic of Sericin Powder from Eri Silk Boiling Water. LIFE : International Journal of Health and Life-Sciences, 1(1), 151-160.
- Peigler. (1989). A Revision of The Indo-Australian Genus *Attacus*. California: The Lepidoptera Research Foundation, Inc. Beverly Hill.
- Pekic, B., Stojanovic, D., Lepojevic, Z. & Tolic, A. (1988). Investigation of Extraction Kinetics of Glycoside Frim Leaves of *Digitalis lanata* Ehrh. Pharmazeutische Industrie, 50 (8), 984-986.

- Peleg, M. A. (1988). Empirical Model For The Description of Moisture Sorption Curves. *Journal of Food Science*, 53, 1216-1219.
- Phanphanich, M. & Mani, S. (2009). Drying Characteristics of Pine Forest Residues. *BioResources*, 5(1), 108-121.
- Sanjuán, N., Cárcel, J. A., Clemente, G., & Mulet, A. (2001). Modeling of The Rehydration Process of Broccoli Florets. *European Food Research and Technology*, 212(4), 449-453.
- Sasaki, M., Yamada, H., & Kato, N. (2000). Consumption of Silk Protein, Sericin Elevates Intestinal Absorption of Zinc, Iron, Magnesium and Calcium in Rats. *Nutrition Research*. 20(10), 1505-1511.
- Shiozaki, S., Ogata, T. & Hiriuchi, S. (2000). Endogenous Polyamines in The Pericarp and Seed of The Grape Berry During Development and Ripening. *Sci. Hort.*, 83, 33-34.
- Sopade, P. A., & Obekpa, J. A. (1990). Modeling Water Absorption in Soybean, Cowpea, and Peanuts at Three Temperatures Using Peleg's Equation. *Journal of Food Science*, 55, 1084-1087.
- Sopade, P. A., & Kaimur, K. (1999). Application of Peleg's Equation in Desorption Studies of Food Systems: A Case Study with Sago (*Metroxylon sagu rottb.*) Starch. *Drying Technology*, 17, 975-989.
- Stankovic, M. Z., Cakic, M. D., Cvetkovic, D. M. & Veljkovic, V. B. (1994). Kinetics of Extraction of Resinoids From Overground Parts of Sweet Clover (*Melilotus officinalis* L.). *Journal Serbian Chemical Society*, 10 (39), 735-741.
- Takasu, Y., Yamada, H., & Tsubouchi, K. (2002). Extraction and Chromatographic Analysis of Cocoon Sericin of the Silkworm, *Bombyx mori*. *Journal of Insect Biotechnology and Sericology*, 71, 151-156.
- Teramoto, H., Kameda, T., & Yamada, Y. (2008). Preparation of Gel Film from *Bombyx mori* Silk Sericin and Its Characterization as a Wound Dressing. *Bioscience, Biotechnology, and Biochemistry*. 72(12), pp. 3189-3196.
- Teresinha. (2017). Silk Cocoons. Retrieved March 15<sup>th</sup>, 2017 from [http://www.wildfibres.co.uk/html/silk\\_cocoons.html](http://www.wildfibres.co.uk/html/silk_cocoons.html).
- Turhan, M., Sayar, S. & Gunasekaran, S. (2002). Application of Peleg Model to Study Water Absorption in Chickpea During Soaking. *Journal of Food Engineering*, 29, 349-360.
- Wakabayashi, S., & Sugioka, M. (1994). Jap Pat.06-017372A.

- Wangkiat, P. (Feb 19<sup>th</sup>, 2017). Ericulture Reeling Them In. Retrieved March 16<sup>th</sup>, 2017 from <http://www.bangkokpost.com/news/special-reports/1201117/ericulture-reeling-them-in>.
- Watson, E. L. & Bhargava, V. K. (1974). Thin-Layer Drying Studies on Wheat. *Canadian Agricultural Engineering*, 16(1), 18-22.
- Wei, T., Li, M. Z., & Xie, R. J. (2005). Preparation and Structure of Porous Silk Sericin Materials. *Macromolecular Materials and Engineering*, 290, 188-194.
- Xinqing, C., Fung, L. K., Fong, M. S., Kwong, C. W., Nok, N. T., & Lun, Y. K. (2012). Assessment of Sericin Biosorbent for Selective Dye Removal. *Chinese Journal of Chemical Engineering*.20(3), 4246-432.
- Yamada, H., & Matsunaga, A. (1994).Jap Pat.06-017373A.
- Yamada, H., & Yuri, O. (1998). Sericin Coated Powders for Cosmetics. Jpn Kokai Tokyo Koho jap 10226626 A2 (to Noevir Co Ltd, Seiren Co Ltd, Japan). *Chem Abstr*. 129(16), 207-209.
- Yamada, H., Nakao, H., Takasu, Y., Tsubouchi, K. (2001). Preparation of Undegraded Native Molecular Fibroin Solution from Silkworm Cocoons. *Materials Science and Engineering C*.14, 41-46.
- Zhang, Y. Q., Tao, M. L., Shen, W. D., Zhou, Y. Z., Ding, Y., Ma, Y., Zhou, W. L. (2004). Immobilization of L-asparaginase on the Microparticles of the Natural Silk Sericin Protein and Its Characters. *Biomaterials*. 25, 3751-3759.
- Zhaorigetu, S., Sasaki, M., Watanabe, H., & Kato, N. (2001). Supplemental Silk Protein, Sericin, Suppresses Colon Tumorigenesis in 1,2-dimethyldrazine-treated mice by Reducing Oxidative Stress and Cell Proliferation. *Biosci, Biochem, Biochem*,65(10), 2181-2186.