

DAFTAR PUSTAKA

- AlOtaibi SF, Blaser S, MacGregor DL. Neurological complications of kernicterus. *Can J Neurol Sci.* 2005;32(3):311-315.
- Angela SJ, Matsuo T, Gonzalez MRC *et al.* CRIB score, birth weight and gestational age in neonatal mortality risk evaluation. *Rev Saude Publica.* 2003;37(5).
- Bano R, Mushtaq A, Adhi M, Asim N, Afzal N. Incidence and outcome of small for gestational age foetuses: an experience from a secondary care hospital. *JPMA - J Pakistan Med Assoc.* 2013;63(11):1422-1424.
- Bardin C, Zelkowitz P *et al.* Outcomes of small for gestational age Infants Born Before 27 Weeks of Gestation. *J Pediatr.* 2013; 163(1):55-60.
- Bari MI, Ullah MA, Khatun M. Morbidity and mortality of low birth weight baby. *J Teach Assoc.* 2008;21(1):1-5.
- Bartels DB, Schwab F, Geffers C *et al.* Nosocomial infection in small for gestational age newborns with birth weight < 1500 g: a multicentre analysis. *Arch Dis Child Fetal Neonatal Ed.* 2007; 92(6):44-53.
- Bhutani V, Johnson L. Kernicterus: a preventable neonatal brain injury. *J Arab Neonatal Forum.* 2005;2:12-23.
- Boguszewski Margaret CS, Mericq Veronica, Bergada Ignacio, Damiani Durval BA. Latin American Consensus: Children Born Small for Gestational Age. *BMC Pediatr.* 2011;11(66):1471-2431.
- Damman O, Naples M, Bednarek F., *et al.* SNAP-II and SNAPPE-II and the risk of structural and functional brain disorders in extremely low gestational age newborns: the ELGAN study. *Neonatology.* 2010; 97(2):71-82.
- Dey AC, Ahmed FU, Mannan MA, Saha L, Barua CC, Mahmood CB. Small for Gestational Age Babies: Morbidity and Immediate Outcome in a Tertiary Care Hospital - A Prospective Study. *Bangladesh J Child Heal.* 2010;31(1):1-7.
- Dorling JS, Field DJ, Manktelo B. Neonatal disease severity scoring systems review. *Arch Dis Child Fetal Neonatal Ed.* 2005;90:F11-16.
- Ezz-eldin Z, Hamid TA, Youssef MRL. Clinical Risk Index for Babies (CRIB II) Scoring System in Prediction of Mortality in Premature Babies. *Clin Diag Research J.* 2015;9(6): 8-11.
- Gargasz A. Neonatal and Pediatric Parenteral Nutrition. *AACN Adv Crit Care.* 2012;23(4):451-464.



Gagliardi L, Cavazza A, Brunelli A, *et al.* Assessing mortality risk in very low birthweight infants: a comparison of CRIB, CRIB II, and SNAPPE-II. *Arch Dis Child Fetal Neonatal Ed.* 2004; 89:F419-22.

Gleason CA DS. *Avery's Diseases of the Newborn 9th Ed.* Philadelphia: Saunders; 2012

Granovsky S, Reichman B, Lerner-Geva L, *et al.* Mortality and morbidity in preterm small-for-gestational-age infants: A population-based study. *Am J Obstet Gynecol.* 2012;206(2):150.

Hameed NN, Alzubaidi MA, Kadhim SH. Risk Factors of Small for Gestational Age Newborn Babies. *Best Pract Res Clin Obstet Gynaecol.* 2011;10(1).

Havranek J. Polycythemia of the newborn. *Pediatr pro Praxi.* 2013;14(4):240-241.

Itabashi K, Mishina J, Tada H, Sakurai M, Nanri Y, Hirohata Y. Longitudinal follow-up of height up to five years of age in infants born preterm small for gestational age; comparison to full-term small for gestational age infants. *Early Hum Dev.* 2007;83(5):327-333.

Jain A, Aggarwal R, Jeeva Sankar M, Agarwal R, Deorari AK, Paul VK. Hypoglycemia in the newborn. *Indian J Pediatr.* 2010;77(10):1137-1142.

Jolin-dahel K, Ferretti E, Montiveros C, Grenon R, Barrowman N, Jimenez-rivera C. Parenteral Nutrition-Induced Cholestasis in Neonates : Where Does the Problem Lie ? *Gastroenterol Res Pract.* 2013;2013:163632.

Johnson L, Bhutani VK *et al.* Hyperbilirubinemia and kernicterus. *Clin Perinatol.* 2006;33(2):387-410.

Kamath BD, MacGuire ER, McClure EM, Goldenberg RL, Jobe AH. Neonatal Mortality From Respiratory Distress Syndrome: Lessons for Low-Resource Countries. *Pediatrics.* 2011;127(6):1139-1146.

Katz J, Lee ACC, Kozuki N, *et al.* Mortality risk in preterm and small-for-gestational-age infants in low-income and middle-income countries: A pooled country analysis. *Lancet.* 2013;382(9890):417-425.

Katz J, Wu LA, Mullany LC, *et al.* Prevalence of small-for-gestational-age and its mortality risk varies by choice of birth-weight-for-gestation reference population. *PLoS One.* 2014;9(3):1-9.

Kavurt S, Ozcan B, Aydemir O, Bas AY, Demirel N. Risk of retinopathy of prematurity in small for gestational age premature infants. *Indian Pediatr.* 2014;51(10):804-806.

Korejo HB, Bhurgri GR, Bhand S, Qureshi MA, Dahri GM, Chohan RK. Risk Factors for Kernicterus in Neonatal Jaundice. *J Med Sci.* 2010;8(1):6-9.



Kozuki N, Lee ACC, Silveira MF, et al. The associations of parity and maternal age with small-for-gestational-age, preterm, and neonatal and infant mortality: a meta-analysis. *BMC Public Health*. 2013;13. (Suppl 3):S2.

Kurinczuk JJ, Hollowell J, Boyd PA, Oakley L, Brocklehurst P, Gray R. The contribution of congenital anomalies to infant mortality. Inequalities infant Mortal Project briefing Paper 4. *Oxford: National Perinatal Epidemiology Unit, University of Oxford*. 2010;1–13.

Laptook A, Salhab W, Bhaskar B. Admission temperature of low birth weight infants: predictor and associated morbidities. *Pediatrics*. 2007;119:e643-9.

Lee ACC, Katz J, Blencowe H, et al. National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. *Lancet Glob Heal*. 2013;1(1).

Lee Peter A, Chernausek Steven D, Hokken-Koelega Anita CS CP. International Small for Gestational Age Advisory Board Consensus Developmental Conference Statement: Management of Short Children Born Small for Gestational Age, April 24-October 1, 2001. *Pediatrics*. 2003;111(6):1253-1261.

Majeed R, Memon Y, Majeed F, et al. Risk factors of birth asphyxia. *J Ayub Med Coll Abbottabad*. 2007;19(3):67-71

Margaretha A, Drott P AR. Epidemiology and trends of necrotizing enterocolitis in Sweden: 1987-2009. *Am Acad Pediatr*. 2013.

McCowan L, Horgan RP. Risk factors for small for gestational age infants. *Best Pract Res Clin Obstet Gynaecol*. 2009;23(6):779-793.

Mizzi J, Muskat RV. Hypothermia in the early neonatal period: Follow-up study. *Malta Med J*. 2007;19(01):34-37.

Mia RA, Etika R, Harianto A et al. The use of score for neonatal acute physiology perinatal extension II (SNAPPE II) in predicting neonatal outcome in neonatal intensive care unit. *Pediatr Indones*. 2005;45:241-5.