

DAFTAR PUSTAKA

- Bergström, U., Björnstig, U., Stenlund, H., Jonsson, H., & Svensson, O. (2008). Fracture mechanisms and fracture pattern in men and women aged 50 years and older: a study of a 12-year population-based injury register, Umeå, Sweden. *Osteoporosis International*, 19(9), 1267-1273.
- Bhandari, M., Devereaux, P. J., Swiontkowski, M. F., Tornetta, P., Obremskey, W., Koval, K. J., ... Guyatt, G. H. (2003). Internal fixation compared with arthroplasty for displaced fractures of the femoral neck. A meta-analysis. *The Journal of Bone and Joint Surgery. American Volume*, 85-A(9), 1673-1681.
- Bhandari, M., Devereaux, P. J., Tornetta, P., Swiontkowski, M. F., Berry, D. J., Haidukewych, G., ... Guyatt, G. H. (2005). Operative management of displaced femoral neck fractures in elderly patients. An international survey. *The Journal of Bone and Joint Surgery. American Volume*, 87(9), 2122-2130.
- Bhandari, M., Schemitsch, E., Jönsson, A., Zlowodzki, M., & Haidukewych, G. J. (2009). Gamma nails revisited: gamma nails versus compression hip screws in the management of intertrochanteric fractures of the hip:

Bosch, U., Schreiber, T., & Krettek, C. (2002). Reduction and fixation of displaced intracapsular fractures of the proximal femur. *Clinical Orthopaedics and Related Research*, (399), 59-71. Retrieved from <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12011695&retmode=ref&cmd=prlinks\papers2://publication/uuid/48EB1920-E387-4BD4-B17E-F8A770412B5D>

Buchols, R.W., Heckman, J.D., Court-Brown CM., (2006). *Rockwood & Green's Fractures in Adults*, 6th editon. USA: Maryland Composition

Butler, M., Forte, M., & Kane, R. (2009). Treatment of common hip fractures. *Evidence Report/technology Assessment*, (184), 1-85, v. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21433326> \n http://pharmaceuticalapprovals.elsevierbi.com/~media/Images/Publications/Archive/The Gray Sheet/35/032/01350320008/080609_ahrq_hipfracture.pdf

Cawthon, P. M., Ewing, S. K., McCulloch, C. E., Ensrud, K. E., Cauley, J. A., Cummings, S. R., & Orwoll, E. S. (2009). Loss of hip BMD in older men: the osteoporotic

Fractures in men (MrOS) study. *Journal of Bone and Mineral Research*, 24(10), 1728-35.

Cooley MR, Koval KJ. Hip fracture. Epidemiology and risk factors. *Techn Orthop* 2004; 19:104-114.

Cole, Z. a., Dennison, E. M., & Cooper, C. (2008). Osteoporosis epidemiology update. *Current Rheumatology Reports*, 10(2), 92-96.

Compston, J. E., Flahive, J., Hosmer, D. W., Watts, N. B., Siris, E. S., Silverman, S., ... Adachi, J. D. (2014). Relationship of weight, height, and body mass index with fracture risk at different sites in postmenopausal women: The global longitudinal study of osteoporosis in women (GLOW). *Journal of Bone and Mineral Research*, 29(2), 487-493.

Crandall, C. J., Yildiz, V. O., Wactawski-Wende, J., Johnson, K. C., Chen, Z., Going, S. B., ... Cauley, J. a. (2015). Postmenopausal weight change and incidence of fracture: post hoc findings from Women's Health Initiative Observational Study and Clinical Trials. *BMJ (Clinical Research Ed.)*, 350(January), h25.

Cranney, A., Jamal, S. a, Tsang, J. F., Josse, R. G., & Leslie, W. D. (2007). Low bone mineral density and fracture burden in postmenopausal women. *CMAJ*:

l'Association Medicale Canadienne, 177(6), 575-580.

Cummings SR, Melton LJ III. Epidemiology and outcomes of osteoporotic fractures. *Lancet* 2002; 359:1761- 1767.

3. Kanis

Cummings, S. R., Black, D. M., Nevitt, M. C., Browner, W., Cauley, J., Ensrud, K., ... Vogt, T. M. (1993). Bone density at various sites for prediction of hip fractures. The Study of Osteoporotic Fractures Research Group. *Lancet, 341*(8837), 72-75.

De Laet, C. E., van Hout, B. a, Burger, H., Hofman, a, & Pols, H. a. (1997). Bone density and risk of hip fracture in men and women: cross sectional analysis. *BMJ (Clinical Research Ed.), 315*(7102), 221-5.

Retrieved from

<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2127146&tool=pmcentrez&rendertype=abstract>

Dennison E, Cole Z, Cooper C. Diagnosis and epidemiology of osteoporosis. *Curr Opin Rheumatol* 2005; 17:456-461.

Ensrud, K. E. (2013). Epidemiology of fracture risk with advancing age. *Journals of Gerontology - Series A*

1236-1242.

Ensrud, K. E., Ewing, S. K., Stone, K. L., Cauley, J. A., Bowman, P. J., & Cummings, S. R. (2003). Intentional and unintentional weight loss increase bone loss and hip fracture risk in older women. *Journal of the American Geriatrics Society*, 51(0002-8614 (Print)), 1740-1747.

Faulkner, K. A., Cauley, J. A., Studenski, S. A., Landsittel, D. P., Cummings, S. R., Ensrud, K. E., ... Nevitt, M. C. (2009). Lifestyle predicts falls independent of physical risk factors. *Osteoporosis International*, 20(12), 2025-2034.

Garden RS, Preston, E. (1961). Low angle fixation in Fractures of the femoral neck. *Surger*, 101(November), 647. Retrieved from <http://web.jbjs.org.uk/cgi/content/abstract/43-B/4/647>

Gullberg, B., Johnell, O., & Kanis, J. a. (1997). World-wide projections for hip fracture. *Osteoporosis International*, 7(5), 407-413.

Hasenboehler, E. A., Agudelo, J. F., Morgan, S. J., Smith, W. R., Hak, D. J., & Stahel, P. F. (2007). Treatment of complex proximal femoral fractures with

the proximal femur locking compression plate.

Orthopedics, 30(8), 618-23. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/17727017>

Haynes, R. C., Pöll, R. G., Miles, a. W., & Weston, R. B.

(1997). Failure of femoral head fixation: A cadaveric analysis of lag screw cut-out with the gamma locking nail and AO dynamic hip screw. *Injury*, 28(5-6), 337-341.

Heetveld, M. J., Raaymakers, E. L. F. B., Luitse, J. S.

K., Nijhof, M., & Gouma, D. J. (2007). Femoral neck fractures: can physiologic status determine treatment choice? *Clinical Orthopaedics and Related Research*, 461, 203-212.

Hillier TA, Rizzo JH, Pedula KL, Stone KL, Cauley JA,

Bauer DC, Cummings SR (2003) Nulliparity and fracture risk in older women: the study of osteoporotic fractures. *J Bone Miner Res* 18 (5):893-899

Kregor, P. J. (1996). The effect of femoral neck

fractures on femoral head blood flow. *Orthopedics*,

19(12), 1031-6; quiz 1037-8. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/8972521>

Kremer, R., Campbell, P. P., Reinhardt, T., & Gilsanz, V.

(2009). Vitamin D status and its relationship to body

fat, final height, and peak bone mass in young women.

The Journal of Clinical Endocrinology and Metabolism.

Madsen, J. E., Naess, L., Aune, a K., Alho, A., Ekeland, A., & Strømsøe, K. (1998). Dynamic hip screw with trochanteric stabilizing plate in the treatment of unstable proximal femoral fractures: a comparative study with the Gamma nail and compression hip screw. *Journal of Orthopaedic Trauma, 12*(4), 241-248.

Marks, R. (2010). Hip fracture epidemiological trends, outcomes, and risk factors, 1970-2009. *International Journal of General Medicine, 3*, 1-17.

Melton III, L. J. (2003). Epidemiology worldwide. *Endocrinol.Metab Clin.North Am., 32*(0889-8529 (Print)), 1-13, v.

Nguyen TV, Jones G, Sambrook PN, White CP, Kelly PJ, Eisman JA (1995) Effects of estrogen exposure and reproductive factors on bone mineral density and osteoporotic fractures. *J Clin Endocrinol Metab 80* (9):2709-2714.

Orwoll ES. Osteoporosis in men. In: Favus MJ, editor. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. Washington DC: The American Society for Bone and Mineral Research; 2006.

pathogenesis of postmenopausal osteoporosis. *Journal of Bone and Mineral Research: The Official Journal of the American Society for Bone and Mineral Research*, 11(8), 1043-1051.

Premaor, M., Parker, R. a., Cummings, S., Ensrud, K., Cauley, J. a., Lui, L. Y., ... Compston, J. (2013). Predictive value of FRAX for fracture in obese older women. *Journal of Bone and Mineral Research*,

Price, S.A., Lorraine, M.W., (2006). Patofisiologi: Konsep Klinis Proses-Proses Penyakit. Edisi 6 Vol 2, Jakarta: Penerbit Buku Kedokteran ECG

Riggs BL, Wahner HW, Melton LJ, et al. Rate of bone loss in the axial and appendicular skeleton of women: evidence of substantial bone loss prior to menopause. *J Clin Invest*.1986;77:1847-1891

Riggs BL, Khosla S, Melton LJ, 3rd (2002) Sex steroids and the construction and conservation of the adult skeleton. *Endocr Rev* 23 (3):279-302.

Seinsheimer, F. (1978). Subtrochanteric fractures of the femur. *J Bone Joint Surg Am.*, 60(3), 300-6.



HUBUNGAN ANTARA USIA, JENIS KELAMIN, DAN INDEKS MASSA TUBUH DENGAN KEJADIAN HIP FRACTURE PADA PASIEN

RAWAT JALAN POLIKLINIK ORTHOPAEDI RSUP DR. SARDJITO YOGYAKARTA

WILLY IRAWAN, Dr. dr. Puntodewo, M.Kes, Sp.OT(K); dr. Yudha Mathan Sakti, Sp.OT

Universitas Gadjah Mada, 2016 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Shah, A. K., Eissler, J., & Radomisli, T. (2002).

Algorithms for the treatment of femoral neck fractures. *Clinical Orthopaedics and Related Research*, (399), 28-34. Retrieved from <http://eutils.ncbi.nlm.nih.gov/entrez/eutils/elink.fcgi?dbfrom=pubmed&id=12011691&retmode=ref&cmd=prlinks\&npapers2://publication/uuid/2A2CD1ED-D5C8-4CD1-8E7B-2F6566305205>

Shapses, S. A., & Sukumar, D. (2012). Bone metabolism in obesity and weight loss. *Annual Review of Nutrition*, 32, 287-309.

Solomon, L., David, W., Selvadurai, N., (2010). *Apley's System of Orthopaedics and Fractures*. 9th edition, UK: Hodder Arnold, an imprint of Hodder Education, Hacette UK Company.

Tang, X., Liu, G., Kang, J., Hou, Y., Jiang, F., Yuan, W., & Shi, J. (2013). Obesity and risk of hip fracture in adults: a meta-analysis of prospective cohort studies. *PloS One*, 8(4), e55077.

Tarantino, U., Cannata, G., Lecce, D., Celi, M., Cerocchi, I., & Iundusi, R. (2007). Incidence of fragility fractures. *Aging Clin Exp Res*, 19(4 Suppl),

<http://www.ncbi.nlm.nih.gov/pubmed/18182886>

US Department of Health and Human Services. (2004). Bone health and osteoporosis: a report of the Surgeon General. *US Health and Human Services*, 437.

Watson, J. T., Moed, B. R., Cramer, K. E., & Karges, D. E. (1998). Comparison of the compression hip screw with the Medoff sliding plate for intertrochanteric fractures. *Clinical Orthopaedics and Related Research*. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/17900766>

WHO expert consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. *The Lancet*, 2004; 157-163.

WHO. Prevention and Management of Osteoporosis. WHO Technical Report Series. 2003

Youm, T., Koval, K. J., Kummer, F. J., & Zuckerman, J. D. (1999). Do all hip fractures result from a fall? *American Journal of Orthopedics (Belle Mead, N.J.)*, 28(3), 190-4. Retrieved from

<http://www.ncbi.nlm.nih.gov/pubmed/10195844>

Zuckerman, J. D. (1996). Hip fracture. *The New England Journal of Medicine*, 334(23), 1519-1525.