

DAFTAR PUSTAKA

- Andrea RJ. (2008). Nutritional determinants of metabolic syndrome. *J. Nutrigenet Nutrigenomics*, 1: 109-17.
- Anonim. (2005) Guidelines for Data Processing and Analysis of the International Physical Activity Questionnaire (IPAQ)- Short and Long Forms.
- Antonius, O. Salim, L. Sandjaja, B. (2012). Sindroma Metabolik Di Kota Jayapura. *Buletin Penelitian Kesehatan*, 41 (4) hal. 200 – 206
- Azevedo, M. R., Luiza, C., Araújo, P., Reichert, F. F., Siqueira, F. V., Cozzensa, M., & Hallal, P. C. (2007). Gender differences in leisure-time physical activity, 52, 8–15. doi:10.1007/s00038-006-5062-1
- Bantas, K., Koesnanto, H., Moelyono, B. (2007). Ukuran Lingkar Pinggang Optimal untuk Identifikasi Sindrom Metabolik pada Populasi Perkotaan di Indonesia Optimal Waist Circumference for Identification of Subjects with Metabolic Syndrome in Indonesian Urban Population, 3–7.
- Beigh SH, Jain S. (2012). Prevalence of Metabolic Syndrome and Gender Differences. *Bioinformation*, 8 (13): 613-6
- Bloom, D.E., Cafiero, E.T., Jané-Llopis, E., Abrahams-Gessel, S., Bloom, L.R., Fathima, S., Feigl, A.B., Gaziano, T., Mowafi, M., Pandya, A., Prettnner, K., Rosenberg, L., Seligman, B., Stein, A.Z., & Weinstein, C. (2011). The Global Economic Burden of Noncommunicable Diseases. Geneva: World Economic Forum.
- Bosak, K. (2012). Managing Metabolic Syndrome: Focus on Physical Activity. *The Journal for Nurse Practitioners*, 8(3), 206–211. doi:10.1016/j.nurpra.2011.09.001
- Brini, O. El, Akhouayri, O. & Gamal, A., 2014. Prevalence of metabolic syndrome and its components based on a harmonious definition among adults in Morocco. , pp.341–346.
- Bulchan, S. (2012). Metabolic Syndrome and Diabetes: Current Asian Perspective. *Asia Pacific Biotech News* 16(1), pp.11-16.
- Carnethon, M. R, Loria, C, M, Hill, J, O, Sidney, S, Savage, P, J, & Liu, K 2004, 'Risk factors for the metabolic syndrome the coronary artery risk development in young adults (CARDIA) study, 1985–2001', *Diabetes care*, vol. 27, no. 11, pp. 2707-2715
- Caro J, Navarro I, Romero P, Priego MA, Martí'nez-Herva' s S, Real JT. (2013). Metabolic effects of regular physical exercise in healthy population. *Endocri-nol Nutr*; 60:167–72.

- Chen, X., Pang, Z., & Li, K. (2009). Dietary fat, sedentary behaviors and the prevalence of the metabolic syndrome among Qingdao adults. *Nutrition, Metabolism, and Cardiovascular Diseases : NMCD*, 19(1), 27–34. doi:10.1016/j.numecd.2008.01.010
- Cho, E.R., Shin, A., Kim, J., Jee, S.H., Sung, J. (2009). Leisure-time physical activity is associated with a reduced risk for metabolic syndrome. *Ann. Epidemiol.* 19, 784–792.
- Chu, A. H. Y., & Ming, F. (2013). Associations of occupational , transportation , household and leisure-time physical activity patterns with metabolic risk factors among middle-aged adults in a middle-income country. *Preventive Medicine*, 57, S14–S17. doi:10.1016/j.ypmed.2012.12.011
- Chu, A. H. Y., & Moy, F. M. (2014). Journal of Science and Medicine in Sport Association between physical activity and metabolic syndrome among Malay adults in a developing country , Malaysia. *Journal of Science and Medicine in Sport*, 17(2), 195–200. doi:10.1016/j.jsams.2013.04.003
- Colombo, C.M. et al. (2013). Short-term effects of moderate intensity physical activity in patients with metabolic syndrome. *Einstein (São Paulo, Brazil)*, 11(3), pp.324–30. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24136759>.
- Constanian. Cho, N.H. (2014). Diabetes Burden & Prevention in Korea and The Western Pasific Region. *Diabetes Research and Clinical Practice*, S282-S287.
- Cornier MA, Dabelea D, Hernandez TL, Lindstrom LC, Steig AJ, Stob NR. (2008). The Metabolic Syndrome *Endocrine Reviews*: 29(7):777–822 .
- Craig, C.L., Marshall, A.L., Sjostron, M., Bauman, A.E., Booth, M.L., Ainstworth, B.E., Pratt, M., Ekelund, U., Yngve, A., Sallis, J.F., Oja, P. (2003), ‘International physical activity questionnaire : 12-country reliability and validity’, *Medicine and Science in Sports and Exercise*, 35(8), 1381-1395.
- Dalusung-angosta, A., & Gutierrez, A. (2013). Prevalence of Metabolic Syndrome among Filipino-Americans : A Cross-Sectional Study. *Applied Nursing Research*, 26(4), 192–197. doi:10.1016/j.apnr.2013.06.005
- Departemen Kesehatan Republik Indonesia. (2013). Laporan Hasil Riset Kesehatan Dasar Indonesia (Riskesdas).
- Ervin RB. (2009). Prevalence of Metabolic syndrome Among Adults 20 Years of Age and Over, by Sex, Age, Race and Ethnicity, and Body Mass Index: United States 2003-2006. *National Health Statistic Reports*.; 5 (13): 1-7
- Escobedo C, Schargrotsky H, Champagne B, Silva H, Boissonnet CP, Vinueza R. (2009). Prevalence of The Metabolic Syndrome in Latin America and Its Assosiation With Sub-Clinical Carotied Atherosclerosis: The CARMELA Cross Sectional Study *Cardiovascular Diabetology*; 852: 1-9

- Esmailnasab, N., Moradi, G. & Delaveri, A. (2012). Risk Factors of Non-Communicable Diseases and Metabolic. , 41(7), pp.77–85.
- Ford, E.S., Kohl III, H.W., Mokdad, A.H., Ajani, U.A. (2005). Sedentary behavior, physical activity, and the metabolic syndrome among U.S. adults. *Obes. Res.* 13, 608–614.
- Ford, E.S., Li, C. (2006). Physical activity or fitness and the metabolic syndrome. *Expert. Rev. Cardiovasc. Ther.* 4, 897–915.
- Grundy, S. M., Brewer, H. B., Cleeman, J. I., Smith, S. C
- Frugé AD, Byrd SH, Fountain BJ, Cossman JS, Schilling MW, Gerard P. (2015). Nutrition , Metabolism & Cardiovascular Diseases Increased physical activity may be more protective for metabolic syndrome than reduced caloric intake . An analysis of estimated energy balance in U . S . adults : 2007 e 2010 NHANES. *Nutr Metab Cardiovasc Dis* [Internet]. Elsevier B.V;;25(6):535–40. Available from: <http://dx.doi.org/10.1016/j.numecd.2015.03.006>
- ., & Lenfant, C. (2004). Definition of Metabolic Syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on Scientific Issues Related to Definition. *Circulation*, 109(3), 433–438. doi:10.1161/01.CIR.0000111245.75752.C6
- Grundy, S. M. (2008). Metabolic Syndrome Pandemic. doi:10.1161/ATVBAHA.107.151092
- Grundy, S. M. (2016). Metabolic syndrome update. *Trends in Cardiovascular Medicine*, 26(4), 364–373. doi:10.1016/j.tcm.2015.10.004
- Grundy, S. M., Cleeman, J. I., Daniels, S. R., Donato, K. A., Eckel, R. H., Franklin, B. A. (2005). Diagnosis and management of the metabolic syndrome an American Heart Association/National Heart, Lung, and Blood Institute scientific statement. *Circulation*, 112(17), 2735–2752.
- Hajian-Tilaki K, Heidari B. Association between parity, live birth and the risk of obesity in women. *Int J Endocrinol Metab* 2007;4:109–18.
- Hajian-tilaki, K., Heidari, B., Firouzjahi, A., Bagherzadeh, M., & Hajian-tilaki, A. (2014). Diabetes & Metabolic Syndrome : Clinical Research & Reviews Prevalence of metabolic syndrome and the association with socio-demographic characteristics and physical activity in urban population of Iranian adults : A population-based study. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 8(3), 170–176. doi:10.1016/j.dsx.2014.04.012
- Hamilton, M.T., Hamilton, D.G., Zderic, T.W. (2007). Role of low energy expenditure and sitting in obesity, metabolic syndrome, type 2 diabetes, and cardiovascular disease. *Diabetes* 56, 2655–2667.

- Hastert, T. a, Gong, J., Campos, H., & Baylin, A. (2015). Physical activity patterns and metabolic syndrome in Costa Rica. *Preventive Medicine*, 70, 39–45. doi:10.1016/j.ypmed.2014.11.006
- IPAQ Research Committe, 2005, *Guidlines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ) – Short and Long form*, diakses tanggal 6 Februari 2015, from www.ipaq.ki.se.
- Irfan, Ainul. (2004). Pergeseran Usaha Tani dan Perkembangan Desa di Kecamatan Turi Kabupaten Sleman. Tesis, Magister Perencanaan Kota dan Daerah, Universitas Gadjah Mada
- Jafar, N. (2012). Sindroma Metabolik dan Epidemiologi. *Media Gizi Masyarakat Indonesia*.1(2):71-78
- Jalal F, Liputo NI, Susanti N, Oenzil F. (2006). Hubungan Lingkar Pinggang dengan Kadar Glukosa Darah, Trigliserida, dan Tekanan Darah pada Etnis Minang di Kabupaten Padang Pariaman, Sumatera Barat. Universitas Andalas.
- Kamso, S., Dharmayati, P., Lubis, U., Juwita, R., Kurnia, Y., & Besral, R. (n.d.). Prevalensi dan Determinan Sindrom Metabolik pada Kelompok Eksekutif di Jakarta dan Sekitarnya Prevalency and Determinant Metabolic Syndrome on Executive Group in Jakarta and Nearby Areas, 85–90.
- Kassi, E. et al. (2011). Metabolic syndrome: definitions and controversies. *BMC medicine*, 9(1), p.48. Available at: <http://www.biomedcentral.com/1741-7015/9/48>.
- Katzmarzyk, P.T. & Lee, I.-M. (2012). Sedentary behaviour and life expectancy in the USA: a cause-deleted life table analysis. *BMJ open*, 2(4), pp.1–8. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3400064&tool=pmcentrez&rendertype=abstrac>.
- Kaur, J. (2014). A Comprehensive Review on Metabolic Syndrome.
- Kementrian Kesehatan Republik Indonesia. (2014). Profil Kesehatan Indonesia 2013.
- Kemenkes RI. (2012). *Penyakit Tidak Menular Penyebab Kematian Terbanyak di Indonesia*, Available from www.depkes.go.id, diakses tanggal 10 Maret 2015.
- Kemenkes RI. (2013). *Info pengendalian penyakit dan penyehatan lingkungan*, Direktorat Jendral pengendalian penyakit dan penyehatan lingkungan.
- Kim J, Tanabe K, Yokoyama N, Zempo H, Kuno S. (2011). Association between physical activity and metabolic syndrome in middle-aged Japanese: a cross-sectional study. *BMC Public Health*; 11(August):624. <http://dx.doi.org/10.1186/1471-2458-11-624>.
- Kim, Y. (2015). Association of Family Composition and Metabolic Syndrome in Korean Adults Aged over 45 Years Old. *Asian Nursing Research*, 9(4), 349–355. doi:10.1016/j.anr.2015.10.006

- Kurtze, N., Rangul, V., Hustvedt, B. (2008). Reability and validity of the International Physical Activity Questionnaire in the Nord-Trøndelag Health Study (HUNT) population of men, *BMC Medical Research Metodology*.
- Ladabaum, U. Mannalithara, A. Myer, Parvathi A. Singh, Gurkirpal. (2014). Obesity, Abdominal Obesity, Physical Activity, and Caloric Intake in US Adults: 1988 to 2010. *The American Journal of Medicine*, 8 (127) pp 717–727.e12
- Lechleitner M. (2008). Obesity and Metabolic Syndrome in the Elderly: a Mini review. *Gerontology*.; 54: 253-9
- Lemeshow, S., Hosmer, D.W., Klar, J. & Lwanga, S. K. (1997). Besar Sampel dalam Penelitian Kesehatan. Yogyakarta, Gadjah Mada University Press
- Lin H-F, Boden-Albala B, Juo SH. (2005). Heritabilities of the metabolic syndrome and its components in the Northern Manhattan Family Study. *Diabetologia*; 48: 2006-12.
- Medina, C., Barquera, S., Janssen, I. (2013), ‘Validity & reliability of the International Physical Questionnaire among adults in Mexico’, *Rev Panam Salud Publica*, 34(1), 21-28.
- Moniruzzaman M, Zaman MM, Islalm MS, Ahasan HAMN, Kabir H, Yasmin R. (2016). Physical activity levels in Bangladeshi adults : results from STEPS survey 2010. *Public Health* [Internet]. Elsevier Ltd; 2016;4–11. Available from: <http://dx.doi.org/10.1016/j.puhe.2016.02.028>
- Muherdiyantiningsih, Ernawati F, Efendi R, Herman S. (2008). Sindrom Metabolik Pada Orang Dewasa Gemuk di Wilayah Bogor. *Penelitian Gizi dan Makanan*, 31(2) : 75-81
- Notoatmodjo, S. (2012), *Metodologi Penelitian Kesehatan*, Rineka Cipta, Jakarta.
- Oguoma, V. M., Nwose, E. U., Skinner, T. C., Richards, R. S., Digban, K. A., & Onyia, I. C. (2016). Diabetes & Metabolic Syndrome : Clinical Research & Reviews Association of physical activity with metabolic syndrome in a predominantly rural Nigerian population. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 10(1), 13–18. doi:10.1016/j.dsx.2015.08.010
- Orho-Melander, M. (2006). The metabolic syndrome: genetics, lifestyle and ethnicity. *Diabetes Voice*, 51(May), pp.21–24.
- Pavey, T.G. et al. (2011). Effect of exercise referral schemes in primary care on physical activity and improving health outcomes : systematic review and meta-analysis. , pp.1–14.
- Qin, X., Zhang, Y., Cai, Y., He, M., Sun, L., Fu, J., ... Huo, Y. (2013). Prevalence of obesity , abdominal obesity and associated factors in hypertensive adults aged 45 e 75 years. *Clinical Nutrition*, 32(3), 361–367. doi:10.1016/j.clnu.2012.08.005

- Rampal S, Mahadeva S, Guallar E, Bulgiba A, Mohamed R, et al. (2012). Ethnic Differences in the Prevalence of Metabolic Syndrome: Results from a Multi-Ethnic Population-Based Survey in Malaysia. *PLoS ONE* 7(9):
- Rejeki, Sri. (2014). Analisis Tingkat Pendapatan Pedagang Perantara Salak Pondoh di Kecamatan Turi Kabupaten Sleman. Skripsi, Fakultas Pertanian, Universitas Negeri Surakarta
- Ristiananda, D.M. (2011). Hubungan antara status gizi dan pola aktivitas fisik dengan sindroma metabolik pada karyawan RSUP Dr. Sardjito Yogyakarta. Skripsi, Fakultas Kedokteran, Universitas Gadjah Mada.
- Roberts CK, Barnard RJ. (2005). Effects of exercise and diet on chronic disease. *J Appl Physiol*; 98: 3-30.
- Rojas-martínez, R., Aguilar-salinas, C. A., Esp, M., & Jiménez-corona, A. (2012). Prevalence of obesity and metabolic syndrome components in Mexican adults without type 2 diabetes or hypertension, 54(1), 7–13.
- Shahab, A. (2007). Sindrom Metabolik. Media Informasi Ilmu Kesehatan dan Kedokteran. Tersedia di: <http://alwia.com>.
- Sihombing, M. (2015). Hubungan Komponen Sindrom Metabolik Dengan Risiko Diabetes Melitus Tipe 2 Di Lima Kelurahan Kecamatan Bogor Tengah. *Media Litbangkes*, 25(4) hal. 219–226
- Sinclair, Kaimi A., Buchwald, Dedra., Bogart, Andy., Henderson, Jeffrey A. (2011). The Prevalence of Metabolic Syndrome and Associated Risk Factors in Northern Plains and Southwest American Indians. *Diabetes Care* 34(1):pp118–120
- Sisson, S.B., Camhi, S.M., Church, T.S., et al. (2009). Leisure time sedentary behavior, occupational/domestic physical activity, and metabolic syndrome in U.S. men and women. *Metab. Syndr. Relat. Disord.* 7, 529–536.
- S.M.Grundy, J. I. Cleeman, S. R. Daniels. (2005). Diagnosis and management of the metabolic syndrome: an American Heart Association/NationalHeart, Lung, and Blood Institute scientific statement, *Circulation*, vol. 112, no. 17, pp. 2735–2752.
- Soewondo, P., Purnamasari, D., Oemardi, M., & Waspadji, S. (2006). Prevalence of Metabolic Syndrome Using NCEP / ATP III Criteria in Jakarta , Indonesia : The Jakarta Primary Non-communicable Disease Risk Factors Surveillance 2006, 199–203.
- Soewondo P, Purnamasari D, Oemardi M, Waspadji S, Soegondo S. (2010). Prevalence of Metabolic Syndrome Using NCEP/ATP III Criteria in Jakarta, Indonesia: The Jakarta Primary Non communicable Disease Risk Factors Surveillance 2006. *Acta Med Indones-Indones J Intern Med*; 42(IV)

- Sudikno., Herdayati, M., Besral. (2010). Hubungan Aktivitas Fisik Dengan Kejadian Obesitas Pada Orang Dewasa di Indonesia (Analisis Data Riskesdas 2007). *Gizi Indonesia*, 33(1):37-49
- Song, Y. & Joung, H. (2012). A traditional Korean dietary pattern and metabolic syndrome abnormalities. *Nutrition, Metabolism and Cardiovascular Diseases*, 22(5), pp.456–462. Available at: <http://dx.doi.org/10.1016/j.numecd.2010.09.002>.
- Sopiyudin, D. (2012), *Besar sampel dan cara pengambilan sampel dalam penelitian kedokteran dan kesehatan*, Edisi 3, Salemba Medika, Jakarta.
- Sugiyanti, E. (2009). Faktor Resiko Obesitas Sentral pada Orang Dewasa di Sulawesi Utara, Gorontalo. Available from: <http://iirc.ipb.ac.id/jspui/bitstream/12345678/11550/2/109esu.pdf>. Diakses pada 15 Maret 2015
- Susan EB. (2006). Physical activity and the metabolic syndrome in Canada. *Appl. Physiol. Nutr. Metab.* 31: 40-7.
- Tarigan, (2005). Hubungan citra tubuh dengan status obesitas, aktivitas fisik, energi kerja, Tesis, Program Pasca Sarjana Ilmu Kesehatan Masyarakat, Universitas Gadjah Mada.
- Tauler, P., Bennasar-Veny, M., Morales-Asencio, J. M., Lopez-Gonzalez, A. a., Vicente-Herrero, T., De Pedro-Gomez, J., ... Aguilo, A. (2014). Prevalence of premorbid metabolic syndrome in Spanish adult workers using IDF and ATPIII diagnostic criteria: Relationship with cardiovascular risk factors. *PLoS ONE*, 9(2). doi:10.1371/journal.pone.0089281
- Tsigos C. (2002). Hypothalamic pituitary adrenal axis, neuroendocrine factors, and stress. *J. Psychosom Res.*; 53 (4): 865-71.
- UPT Pusat Kesehatan Masyarakat Turi. (2014). Profil Kesehatan Puskesmas Turi 2014, Kabupaten Sleman
- Weiss LA, Pan L, Abney M. (2006). The sex-specific genetic architecture of quantitative traits in humans. *Nat Genet*, 38: 218-22.
- World Health Organization. (2003). *Adherence to long-term therapies evidence for action*, WHO, Switzerland, diakses tanggal 12 Februari 2015, from http://www.who.int/entity/chp/knowledge/publications/adherence_full_report.pdf?ua=1.
- World Health Organization. (2012). Health Topic: Physical Activity, diakses tanggal 20 Februari 2015 from http://www.who.int/topics/physical_activity/en.
- WHO. (2014). Noncommunicable Diseases Country Profiles. Available at http://www.who.int/nmh/publications/ncd_profiles_report.pdf [Access 7 Maret 2015]

- Wiardani, NK., Arsana, IWJ. (2011). Kejadian Sindroma Metabolik Berdasarkan Status Obesitas pada Masyarakat Perkotaan di Denpasar. *Jurnal Ilmu Gizi*, 2(2), pp 129-138.
- Widagdo, Tina Y. (2010). Hubungan Lingkar Pinggang Dengan Beberapa Faktor Risiko Sindrom Metabolik. Skripsi, Fakultas Kedokteran, Universitas Diponegoro Semarang
- Widjaja, F. F., Widjaja, I. R., Astria, Y., Suwita, C. S., & Waspadji, S. (n.d.). Metabolic syndrome and Framingham risk score in obese young adults, 1, 100–106.
- Wiklund P. (2016). The role of physical activity and exercise in obesity and weight management: Time for critical appraisal. *J Sport Heal Sci* [Internet]. Elsevier B.V.;(April):10–3. Available from: <http://dx.doi.org/10.1016/j.jshs.2016.04.001>
- Yunita, R. (2011). Hubungan Pola Aktivitas dengan Sindroma Metabolik pada Peserta GMC Health Center. Skripsi, Fakultas Kedokteran, Universitas Gadjah Mada