

## Daftar Pustaka

- Abeywickrama, K. R. W., Ratnasooriya, W. D., & Amarakoon, A. M. T. (2011). Oral hypoglycaemic, antihyperglycaemic and antidiabetic activities of Sri Lankan Broken Orange Pekoe Fannings (BOPF) grade black tea (*Camellia sinensis* L.) in rats. *Journal of Ethnopharmacology*, 135(2), 278–286.
- ADA. (2014). Standards of medical care in diabetes d 2014. *Diabetes Care*, 37(October 2013), 14–80.
- Adeyemi, D. O., Ukwenya, V. O., Obuotor, E. M., & Adewole, S. O. (2014). Anti-hepatotoxic activities of *Hibiscus sabdariffa* L. in animal model of streptozotocin diabetes-induced liver damage. *BMC Complementary and Alternative Medicine*, 14(277), 1–11.
- Adisakwattana, S., Ruengsamran, T., Kampa, P., & Sompong, W. (2012). In vitro inhibitory effects of plant-based foods and their combinations on intestinal  $\alpha$ -glucosidase and pancreatic  $\alpha$ -amylase. *BMC Complementary and Alternative Medicine*, 12(1), 1–8.
- Alarcon-Aguilar, F. J., Zamilpa, A., Perez-Garcia, M. D., Almanza-Perez, J. C., Romero-Nuñez, E., Campos-Sepulveda, E. A., Vazquez-Carillo, L.I., Roman-Ramos, R. (2007). Effect of *Hibiscus sabdariffa* on obesity in MSG mice. *Journal of Ethnopharmacology*, 114, 66–71.
- Almoosawi, S., Fyfe, L., Ho, C., & Al-Dujaili, E. (2010). The effect of polyphenol-rich dark chocolate on fasting capillary whole blood glucose, total cholesterol, blood pressure and glucocorticoids in healthy overweight and obese subjects. *The British Journal of Nutrition*, 103, 842–850.
- Anton, S. D., Martin, C. K., Han, H., Coulon, S., Cefalu, W. T., Geiselman, P., & Williamson, D. a. (2010). Effects of stevia, aspartame, and sucrose on food intake, satiety, and postprandial glucose and insulin levels. *Appetite*, 55(1), 37–43.
- Aronoff, S. L., Berkowitz, K., Shreiner, B., & Want, L. (2004). Glucose Metabolism and Regulation : Beyond Insulin and Glucagon. *Diabetes Spectrum*, 17, 183–190.
- Avignon, A., Hokayem, M., Bisbal, C., & Lambert, K. (2012). Dietary antioxidants : Do they have a role to play in the ongoing fight against abnormal glucose metabolism ? *Nutrition*, 28(7-8), 715–721.

Aziz, Z., Wong, S. Y., & Chong, N. J. (2013). Effects of *Hibiscus sabdariffa* L. on serum lipids: A systematic review and meta-analysis. *Journal of Ethnopharmacology*, 150(2), 442–450.

Badan Penelitian dan Pengembangan Kesehatan. (2013). *Riset kesehatan dasar 2013* (Vol. 306, pp. 253–255). Jakarta.

Bagian Kependudukan Biro tata Pemerintahan Setda DIY (2015). Profil kependudukan DIY dalam angka. Retrieved May 15, 2016, from [www.kependudukan.jogjaprovo.go.id](http://www.kependudukan.jogjaprovo.go.id)

Baynes, J. W. (1991). Perspectives in diabetes role of oxidative stress in development of complications in diabetes. *Diabetes*, 40(December 1990), 405–412.

Beltrán-Debón, R., Alonso-Villaverde, C., Aragonès, G., Rodríguez-Medina, I., Rull, A., Micol, V., Segura-Carretero, A., Fernández-Gutiérrez, A., Camps, J., Joven, J. (2010). The aqueous extract of *Hibiscus sabdariffa* calices modulates the production of monocyte chemoattractant protein-1 in humans. *Phytomedicine*, 17, 186–191.

Benjamin, S. M., Valdez, R., Geiss, L. S., Rolka, D. B., & Narayan, K. M. V. (2003). Estimated number of adults with prediabetes in the U.S. in 2000. *Diabetes Care*, 26(3), 664–649.

Carakostas, M. C., Curry, L. L., Boileau, A. C., & Brusick, D. J. (2008). Overview : The history, technical function and safety of rebaudioside A, a naturally occurring steviol glycoside, for use in food and beverages. *Food and Chemical Toxicology*, 46, 1–10.

CDC. (2014). Prediabetes. Retrieved January 1, 2015, from <http://www.cdc.gov/diabetes/basics/prediabetes.html>

Chandalia, M., Gard, A., Dieter, L., Bergmann, K. Von, Grundy, S. M., & Brinkley, L. (2000). Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *The New England Journal of Medicine*, 342(19), 1392–1398.

Chiva-blanch, G., Urpi-sarda, M., Ros, E., Valderas-martinez, P., Casas, R., Arranz, S., Guillén, M., Lamuela-Raventós, R. M., Llorach, R., Andres-lacueva, C., Estruch, R. (2013). Effects of red wine polyphenols and alcohol on glucose metabolism and the lipid profile : A randomized clinical trial. *Clinical Nutrition*, 32(2), 200–206.

Christian, K. R., & Jackson, J. C. (2009). Changes in total phenolic and monomeric anthocyanin composition and antioxidant activity of three

varieties of sorrel (*Hibiscus sabdariffa*) during maturity. *Journal of Food Composition and Analysis*, 22(7-8), 663–667.

Curry, L. L., & Roberts, A. (2008). Subchronic toxicity of rebaudioside A. *Food and Chemical Toxicology*, 46, 11–20.

Da-Costa-Rocha, I., Bonnlaender, B., Sievers, H., Pischel, I., & Heinrich, M. (2014). *Hibiscus sabdariffa* L. - A phytochemical and pharmacological review. *Food Chemistry*, 165, 424–443.

Defronzo, R. A., & Abdul-ghani, M. (2011). Assessment and treatment of cardiovascular risk in prediabetes : impaired glucose tolerance and impaired fasting glucose. *The American Journal of Cardiology*, 108(3), 3B–24B.

Defuria, J., Bennett, G., Strissel, K. J., Perfield, J. W. I., Milbury, P. E., Greenberg, A. S., & Obin, M. S. (2009). Dietary blueberry attenuates whole-body insulin resistance in high fat-fed mice by reducing adipocyte death and its inflammatory sequelae. *Journal of Nutrition*, 139, 1510–1516.

Ditjen PP & PL. (2008). *Petunjuk teknis pengukuran faktor risiko diabetes melitus* (p. 9). Jakarta: Departemen Kesehatan RI.

Effendi, A. T., & Waspadji, S. (2011). *Aspek biomolekular diabetes melitus II* (pp. 153–156). Jakarta: Badan Penerbit FKUI.

Ersoy, C., & Imamoglu, S. (2006). Comparison of the obesity risk and related factors in employed and unemployed (housewife) premenopausal urban women. *Diabetes Research and Clinical Practice*, 72, 190–196.

Eshak, E. S., Iso, H., Mizoue, T., Inoue, M., Noda, M., & Tsugane, S. (2013). Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. *Clinical Nutrition*, 32(2), 300–308.

Evans, J. L., Goldfine, I. D., Maddux, B. A., & Grodsky, G. M. (2003). Are oxidative stress-activated signaling pathways mediators of insulin resistance and cell dysfunction? *Diabetes*, 52(1), 1–8.

Gregersen, S., Jeppesen, P. B., Holst, J. J., & Hermansen, K. (2004). Antihyperglycemic effects of stevioside in type 2 diabetic subjects. *Metabolism*, 53(1), 73–76.

Gurrola-Díaz, C. M., García-López, P. M., Sánchez-Enríquez, S., Troyo-Sanromán, R., Andrade-González, I., & Gómez-Leyva, J. F. (2010). Effects of *Hibiscus sabdariffa* extract powder and preventive treatment (diet) on the lipid profiles of patients with metabolic syndrome (MeSy). *Phytomedicine*, 17, 500–505.

- Harrison, A. P., Cooper, R. G., Suliman, M. A., & AlAlami, U. (2009). The Efficacy of karkadeh tea in controlling post-prandial blood glucose levels. *Am. J. Pharm. & Toxicol.* 4(4), 149–155.
- Herranz-López, M., Fernández-Arroyo, S., Pérez-Sanchez, A., Barrajon-Catalán, E., Beltrán-Debón, R., Menéndez, J. A., Alonso-Villaverde, C., Segura-Carretero, A., Joven, J., Micol, V. (2012). Synergism of plant-derived polyphenols in adipogenesis: Perspectives and implications. *Phytomedicine*, 19(3-4), 253–261.
- Herrera-Arellano, A., Flores-Romero, S., Chávez-Soto, M. A., & Tortoriello, J. (2004). Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild to moderate hypertension: A controlled and randomized clinical trial. *Phytomedicine*, 11, 375–382.
- Hosoda, K., Wang, M.-F., Liao, M.-L., Chuang, C.-K., Iha, M., Clevidence, B., & Yamamoto, S. (2003). Antihyperglycemic effect of oolong tea in type 2 diabetes. *Diabetes Care*, 26(June), 1714–1718.
- Hu, F. B., Li, T. Y., Colditz, G. A., Willett, W. C., & Manson, J. E. (2003). Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *American Medical Association*, 289(14), 1785–1791.
- Hu, F. B., Manson, J. E., Stamofor, M. J., Colditz, G., Liu, S., Solomon, C. G., & Willet, W. C. (2001). Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *The New England Journal of Medicine*, 345(11), 790–797.
- Kardika, I. B. W., Herawati, S., & Yasa, I. W. P. S. (2015). *Preanalitik dan interpretasi glukosa darah untuk diagnosis diabetes melitus* (pp-7-8). Bagian Patologi Klinik Fakultas Kedokteran Universitas Udayana Rumah Sakit Umum Pusat Sanglah.
- Kementerian Kesehatan RI. (2014). *Buku studi diet total : survei konsumsi makanan individu Indonesia 2014* (p.143). Jakarta: Balai Penelitian dan Pengembangan Kesehatan.
- Kim, J. K., So, H., Youn, M. J., Kim, H. J., Kim, Y., Park, C., Kim, S.J., Ha, Y. A., Chai, K. Y., Kim, S. M., Kim, K. Y., Park, R. (2007). *Hibiscus sabdariffa* L. water extract inhibits the adipocyte differentiation through the PI3-K and MAPK pathway. *Journal of Ethnopharmacology*, 114, 260–267.
- Larasati, Y. (2010). Pengaruh pemberian seduhan kelopak rosella (*Hibiscus sabdariffa*) dosis bertingkat selama 30 hari terhadap gambaran histologik gaster tikus wistar. *Skripsi*. Universitas Diponegoro, Semarang. hal. 1-16.

- Lemeshow, S., Hosmer, D. W., Klar, J., & Lwanga, S. K. (1990) *Adequacy of sample size in health studies* (First Edit). Wiley
- Lemus-Mondaca, R., Vega-gálvez, A., Zura-Bravo, L., & Ah-Hen, K. (2012). *Stevia rebaudiana* Bertoni, source of a high-potency natural sweetener : A comprehensive review on the biochemical, nutritional and functional aspects. *Food Chemistry*, 132(3), 1121–1132.
- Lin, T.-L., Lin, H.-H., Chen, C.-C., Lin, M.-C., Chou, M.-C., & Wang, C.-J. (2007). *Hibiscus sabdariffa* extract reduces serum cholesterol in men and women. *Nutrition Research*, 27, 140–145.
- Linder, M. C. (1992). *Biokimia nutrisi dan metabolisme dengan pemakaian secara klinis*. (A. Parakkasi, Ed.) (pp. 33–37). Jakarta: UI-Press.
- Mahmoud, Y. I. (2012). Effect of extract of Hibiscus on the ultrastructure of the testis in adult mice. *Acta Histochemica*, 114(4), 342–348.
- Mardiah, Zakaria, F. R., Prangdimurti, E., & Damanik, R. (2015). Anti-inflammatory of purple roselle extract in diabetic rats induced by streptozotocin. *Procedia Food Science*, 3, 182–189.
- Meyer, K. A., Kushi, L. H., Jacobs, D. R., Slavin, J., Sellers, T. A., & Folsom, A. R. (2000). Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr*, 71, 921–30.
- Mohamed, B. B., Sulaiman, A. A., & Dahab, A. A. (2012). Roselle (*Hibiscus sabdariffa* L.) in Sudan, cultivation and their uses. *Bull. Environ. Pharmacol. Life Sci*, 1(May), 48–54.
- Mohamed, S. (2014). Functional foods against metabolic syndrome (obesity, diabetes, hypertension and dyslipidemia) and cardiovascular disease. *Trends in Food Science and Technology*, 35(2), 114–128.
- Mozaffari-Khosravi, H., Jalali-Khanabadi, B., Afkhami-Ardekani, M., Fatehi, F., & Noori-Shadkam, M. (2009). The effects of sour tea (*Hibiscus sabdariffa*) on hypertension in patients with type II diabetes. *Journal of Human Hypertension*, 23, 48–54.
- Mozaffari-Khosravi, H., Ahadi, Z., & Tafti, M. F. (2014). The effect of green tea versus sour tea on insulin resistance, lipids profiles and oxidative stress in patients with type 2 diabetes mellitus : a randomized clinical trial. *Iran J Med Sci*, 39(5), 424–432.
- Murianingrum, M. (2013). Pelepasan varietas baru rosela minuman (*Hibiscus sabdariffa* var. *sabdariffa*). Retrieved January 1, 2015, from

[http://balittas.litbang.pertanian.go.id/ind/index.php?option=com\\_content&view=article&id=279:rosela&catid=4:info-aktual&Itemid=5](http://balittas.litbang.pertanian.go.id/ind/index.php?option=com_content&view=article&id=279:rosela&catid=4:info-aktual&Itemid=5)

Nicholson, W. K., Asao, K., Brancati, F., Coresh, J., Pankow, J. S., & Powe, N. R. (2006). Parity and risk of type 2 diabetes. *Diabetes Care*, 29(11), 2349–2354.

NRCS. (2015). *Hibiscus sabdariffa* L. (roselle). Retrieved January 1, 2015, from <http://plants.usda.gov/core/profile?symbol=hisa2>

Onyenekwe, P. C., Ajani, E. O., Ameh, D. A., & Gamaniel, K. S. (1999). Antihypertensive effect of roselle (*Hibiscus sabdariffa*) calyx infusion in spontaneously hypertensive rats and a comparison of its toxicity with that in Wistar rats. *Cell Biochemistry and Function*, 17(3), 199–206. (Abstract).

Patel, S. (2014). *Hibiscus sabdariffa*: An ideal yet under-exploited candidate for nutraceutical applications. *Biomedicine and Preventive Nutrition*, 4(1), 23–27.

Pemerintah Kecamatan Depok Kabupaten Sleman. (2015). Peta wilayah. Retrieved May 15, 2016, from [www.depokkec.slemankab.go.id/profile/peta-wilayah](http://www.depokkec.slemankab.go.id/profile/peta-wilayah)

Peng, C. H., Chyau, C. C., Chan, K. C., Chan, T. H., Wang, C. J., & Huang, C. N. (2011). *Hibiscus sabdariffa* polyphenolic extract inhibits hyperglycemia, hyperlipidemia, and glycation-oxidative stress while improving insulin resistance. *Journal of Agricultural and Food Chemistry*, 59, 9901–9909.

Pérez-Ramírez, I. F., Castaño-Tostado, E., Ramírez-de León, J. a., Rocha-Guzmán, N. E., & Reynoso-Camacho, R. (2015). Effect of stevia and citric acid on the stability of phenolic compounds and in vitro antioxidant and antidiabetic capacity of a roselle (*Hibiscus sabdariffa* L.) beverage. *Food Chemistry*, 172, 885–892.

Perkumpulan Endokrinologi Indonesia (Perkeni). (2011). *Konsensus pengelolaan dan pencegahan diabetes melitus tipe 2 di Indonesia*. Jakarta: PB. Perkeni.

Prior, R. L., Xianli, W., Liwei, G., Hager, T. J., Hager, A., & Howard, L. R. (2008). Whole berries versus berry anthocyanins : interactions with dietary fat levels in the C57BL / 6J mouse model of obesity. *J. Agric. Food Chem*, 56, 647–653.

Raini, M., & Isnawati, A. (2011). Kajian: khasiat dan keamanan stevia sebagai pemanis pengganti gula. *Media Litbang Kesehatan*, 21(4), 145–156.



- Rehman, A., Setter, S. M., & Vue, M. H. (2011). Drug-induced glucose alterations part 2 : drug-induced hyperglycemia. *Diabetes Spectrum*, 24(4), 234–238.
- Rosemary, Rosidah, & Haro, G. (2014). Antidiabetic effect of roselle calyces extract (*Hibiscus Sabdariffa* L.) in streptozotocin induced mice. *International Journal of PharmTech Research*, 6(5), 1703–1711.
- Rosenberg, D. E., Norman, G. J., Wagner, N., Patrick, K., Calfas, K. J., & Sallis, J. F. (2010). Reliability and validity of the sedentary behavior questionnaire (SBQ) for adults. *Journal of Physical Activity and Health*, 7, 697–705.
- Sabzghabae, A. M., Ataei, E., Kelishadi, R., Ghannadi, A., Soltani, R., Badri, S., & Shirani, S. (2013). Effect of *Hibiscus sabdariffa* calices on dyslipidemia in obese adolescents: a triple-masked randomized controlled trial. *Materia Socio-Medica*, 25(January), 76–9.
- Salas-Salvadó, J., Martínez-González, M. Á., Bulló, M., & Roz, E. (2011). The role of diet in the prevention of type 2 diabetes. *Nutrition, Metabolism and Cardiovascular Diseases*, 21, 832–848.
- Sancho, R. A. S., & Pastore, G. M. (2012). Evaluation of the effects of anthocyanins in type 2 diabetes. *Food Research International*, 46(1), 378–386.
- Sangnate, S., Wattanavijitkul, T., & Sangwatanaroj, S. (2011). Diuretic effects of roselle (*Hibiscus sabdariffa*) – Stevia (*Stevia rebaudiana*) tea compared with hydrochlorothiazide in diabetic patients with hypertension. *Thai Pharm Health Sci*, 6(4), 265–273. (Abstract).
- Sasaki, R., Nishimura, N., Hoshino, H., Isa, Y., Kadowaki, M., Ichi, T., Tanaka, A., Nishimuri, S., Fukuda, I., Ashida, H., Horio, F., Tsuda, T. (2007). Cyanidin 3-glucoside ameliorates hyperglycemia and insulin sensitivity due to downregulation of retinol binding protein 4 expression in diabetic mice. *Biochemical Pharmacology*, 74, 1619–1627.
- Sharma, R., Yadav, R., & Manivannan, E. (2012). Study of effect of *Stevia rebaudiana* bertonii on oxidative stress in type-2 diabetic rat models. *Biomedicine & Aging Pathology*, 2(3), 126–131.
- Shaw, J. E., Sicree, R. A., & Zimmet, P. Z. (2010). Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Research and Clinical Practices*, 87, 4–14.

- Sinaga, E. (2012). Pengaruh pemberian susu kedelai terhadap kadar glukosa darah puasa pada wanita prediabetes. *Skripsi*. Universitas Diponegoro, Semarang. hal. 1-41.
- Sindi, H. A., Marshall, L. J., & Morgan, M. R. A. (2014). Comparative chemical and biochemical analysis of extracts of *Hibiscus sabdariffa*. *Food Chemistry*, 164, 23–29.
- Soewondo, P., & Pramono, L. (2011). Prevalence, characteristics, and predictors of pre-diabetes in Indonesia. *Medical Journal of Indonesia*, 25(4), 238–243.
- Stull, A. J., Cash, K. C., Johnson, W. D., Champagne, C. M., & Cefalu, W. T. (2010). Bioactives in blueberries improve insulin sensitivity in obese, insulin-resistant men and women. *Journal of Nutrition*, 140(4), 1764–1768.
- Tabák, A. G., Herder, C., Rathmann, W., Brunner, E. J., & Kivimäki, M. (2012). Prediabetes: a high-risk state for diabetes development. *The Lancet*, 379(9833), 2279–2290.
- Tang, W., Li, S., Liu, Y., Huang, M.-T., & Ho, C.-T. (2013). Anti-diabetic activity of chemically profiled green tea and black tea extracts in a type 2 diabetes mice model via different mechanisms. *Journal of Functional Foods*, 5(4), 1784–1793.
- Vue, M. H., & Setter, S. M. (2011). Drug-induced glucose alterations part 1 : drug-induced hypoglycemia. *Diabetes Spectrum*, 24(3), 171–177.
- Wautier, J. L., Boulanger, E., & Wautier, M. P. (2006). Postprandial hyperglycemia alters inflammatory and hemostatic parameters. *Diabetes Metabolism*, 32, 2S34–2S36.
- Whiting, D. R., Guariguata, L., Weil, C., & Shaw, J. (2011). IDF diabetes atlas : global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Research and Clinical Practice*, 94(3), 311–321.
- Wilson, R. D., & Islam, M. S. (2012). Fructose-fed streptozotocin-injected rat: an alternative model for type 2 diabetes. *Pharmacological Reports : PR*, 64(1), 129–39.