



Daftar Pustaka

- Abeywickrama, K. R. W., Ratnasooriya, W. D., & Amarakoon, A. M. T. (2011). Oral hypoglycaemic, antihyperglycaemic and antidiabetic activities of Sri Lankan Broken Orange Pekoe Fannings (BOPF) grade black tea (*Camellia sinensis* L.) in rats. *Journal of Ethnopharmacology*, 135(2), 278–286.
- ADA. (2014). Standards of medical care in diabetes d 2014. *Diabetes Care*, 37(October 2013), 14–80.
- Adeyemi, D. O., Ukwanya, V. O., Obuotor, E. M., & Adewole, S. O. (2014). Anti-hepatotoxic activities of *Hibiscus sabdariffa* L. in animal model of streptozotocin diabetes-induced liver damage. *BMC Complementary and Alternative Medicine*, 14(277), 1–11.
- Adisakwattana, S., Ruengsamran, T., Kampa, P., & Sompong, W. (2012). In vitro inhibitory effects of plant-based foods and their combinations on intestinal α -glucosidase and pancreatic α -amylase. *BMC Complementary and Alternative Medicine*, 12(1), 1–8.
- Alarcon-Aguilar, F. J., Zamilpa, A., Perez-Garcia, M. D., Almanza-Perez, J. C., Romero-Nuñez, E., Campos-Sepulveda, E. A., Vazquez-Carillo, L.I., Roman-Ramos, R. (2007). Effect of *Hibiscus sabdariffa* on obesity in MSG mice. *Journal of Ethnopharmacology*, 114, 66–71.
- Almoosawi, S., Fyfe, L., Ho, C., & Al-Dujaili, E. (2010). The effect of polyphenol-rich dark chocolate on fasting capillary whole blood glucose, total cholesterol, blood pressure and glucocorticoids in healthy overweight and obese subjects. *The British Journal of Nutrition*, 103, 842–850.
- Anton, S. D., Martin, C. K., Han, H., Coulon, S., Cefalu, W. T., Geiselman, P., & Williamson, D. a. (2010). Effects of stevia, aspartame, and sucrose on food intake, satiety, and postprandial glucose and insulin levels. *Appetite*, 55(1), 37–43.
- Aronoff, S. L., Berkowitz, K., Shreiner, B., & Want, L. (2004). Glucose Metabolism and Regulation : Beyond Insulin and Glucagon. *Diabetes Spectrum*, 17, 183–190.
- Avignon, A., Hokayem, M., Bisbal, C., & Lambert, K. (2012). Dietary antioxidants : Do they have a role to play in the ongoing fight against abnormal glucose metabolism ? *Nutrition*, 28(7-8), 715–721.



Aziz, Z., Wong, S. Y., & Chong, N. J. (2013). Effects of *Hibiscus sabdariffa* L. on serum lipids: A systematic review and meta-analysis. *Journal of Ethnopharmacology*, 150(2), 442–450.

Badan Penelitian dan Pengembangan Kesehatan. (2013). *Riset kesehatan dasar 2013* (Vol. 306, pp. 253–255). Jakarta.

Bagian Kependudukan Biro tata Pemerintahan Setda DIY (2015). Profil kependudukan DIY dalam angka. Retrieved May 15, 2016, from www.kependudukan.jogjaprov.go.id

Baynes, J. W. (1991). Perspectives in diabetes role of oxidative stress in development of complications in diabetes. *Diabetes*, 40(December 1990), 405–412.

Beltrán-Debón, R., Alonso-Villaverde, C., Aragonès, G., Rodríguez-Medina, I., Rull, A., Micol, V., Segura-Carretero, A., Fernández-Gutiérrez, A., Camps, J., Joven, J. (2010). The aqueous extract of *Hibiscus sabdariffa* calices modulates the production of monocyte chemoattractant protein-1 in humans. *Phytomedicine*, 17, 186–191.

Benjamin, S. M., Valdez, R., Geiss, L. S., Rolka, D. B., & Narayan, K. M. V. (2003). Estimated number of adults with prediabetes in the U.S. in 2000. *Diabetes Care*, 26(3), 664–649.

Carakostas, M. C., Curry, L. L., Boileau, A. C., & Brusick, D. J. (2008). Overview : The history, technical function and safety of rebaudioside A, a naturally occurring steviol glycoside, for use in food and beverages. *Food and Chemical Toxicology*, 46, 1–10.

CDC. (2014). Prediabetes. Retrieved January 1, 2015, from <http://www.cdc.gov/diabetes/basics/prediabetes.html>

Chandalia, M., Gard, A., Dieter, L., Bergmann, K. Von, Grundy, S. M., & Brinkley, L. (2000). Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *The New England Journal of Medicine*, 342(19), 1392–1398.

Chiva-blanch, G., Urpi-sarda, M., Ros, E., Valderas-martinez, P., Casas, R., Arranz, S., Guillén, M., Lamuela-Raventós, R. M., Llorach, R., Andres-lacueva, C., Estruch, R. (2013). Effects of red wine polyphenols and alcohol on glucose metabolism and the lipid profile : A randomized clinical trial. *Clinical Nutrition*, 32(2), 200–206.

Christian, K. R., & Jackson, J. C. (2009). Changes in total phenolic and monomeric anthocyanin composition and antioxidant activity of three



- varieties of sorrel (*Hibiscus sabdariffa*) during maturity. *Journal of Food Composition and Analysis*, 22(7-8), 663–667.
- Curry, L. L., & Roberts, A. (2008). Subchronic toxicity of rebaudioside A. *Food and Chemical Toxicology*, 46, 11–20.
- Da-Costa-Rocha, I., Bonnlaender, B., Sievers, H., Pischel, I., & Heinrich, M. (2014). *Hibiscus sabdariffa* L. - A phytochemical and pharmacological review. *Food Chemistry*, 165, 424–443.
- Defronzo, R. A., & Abdul-ghani, M. (2011). Assessment and treatment of cardiovascular risk in prediabetes : impaired glucose tolerance and impaired fasting glucose. *The American Journal of Cardiology*, 108(3), 3B–24B.
- Defuria, J., Bennett, G., Strissel, K. J., Perfield, J. W. I., Milbury, P. E., Greenberg, A. S., & Obin, M. S. (2009). Dietary blueberry attenuates whole-body insulin resistance in high fat-fed mice by reducing adipocyte death and its inflammatory sequelae. *Journal of Nutrition*, 139, 1510–1516.
- Ditjen PP & PL. (2008). *Petunjuk teknis pengukuran faktor risiko diabetes melitus* (p. 9). Jakarta: Departemen Kesehatan RI.
- Effendi, A. T., & Waspadji, S. (2011). *Aspek biomolekular diabetes melitus II* (pp. 153–156). Jakarta: Badan Penerbit FKUI.
- Ersoy, C., & Imamoglu, S. (2006). Comparison of the obesity risk and related factors in employed and unemployed (housewife) premenopausal urban women. *Diabetes Research and Clinical Practice*, 72, 190–196.
- Eshak, E. S., Iso, H., Mizoue, T., Inoue, M., Noda, M., & Tsugane, S. (2013). Soft drink, 100% fruit juice, and vegetable juice intakes and risk of diabetes mellitus. *Clinical Nutrition*, 32(2), 300–308.
- Evans, J. L., Goldfine, I. D., Maddux, B. A., & Grodsky, G. M. (2003). Are oxidative stress-activated signaling pathways mediators of insulin resistance and sell dysfunction? *Diabetes*, 52(1), 1–8.
- Gregersen, S., Jeppesen, P. B., Holst, J. J., & Hermansen, K. (2004). Antihyperglycemic effects of stevioside in type 2 diabetic subjects. *Metabolism*, 53(1), 73–76.
- Gurrola-Díaz, C. M., García-López, P. M., Sánchez-Enríquez, S., Troyo-Sanromán, R., Andrade-González, I., & Gómez-Leyva, J. F. (2010). Effects of *Hibiscus sabdariffa* extract powder and preventive treatment (diet) on the lipid profiles of patients with metabolic syndrome (MeSy). *Phytomedicine*, 17, 500–505.



Harrison, A. P., Cooper, R. G., Suliman, M. A., & AlAlami, U. (2009). The Efficacy of karkadeh tea in controlling post-prandial blood glucose levels. *Am. J. Pharm. & Toxicol.* 4(4), 149–155.

Herranz-López, M., Fernández-Arroyo, S., Pérez-Sánchez, A., Barrajón-Catalán, E., Beltrán-Debón, R., Menéndez, J. A., Alonso-Villaverde, C., Segura-Carretero, A., Joven, J., Micol, V. (2012). Synergism of plant-derived polyphenols in adipogenesis: Perspectives and implications. *Phytomedicine*, 19(3-4), 253–261.

Herrera-Arellano, A., Flores-Romero, S., Chávez-Soto, M. A., & Tortoriello, J. (2004). Effectiveness and tolerability of a standardized extract from *Hibiscus sabdariffa* in patients with mild to moderate hypertension: A controlled and randomized clinical trial. *Phytomedicine*, 11, 375–382.

Hosoda, K., Wang, M.-F., Liao, M.-L., Chuang, C.-K., Iha, M., Clevidence, B., & Yamamoto, S. (2003). Antihyperglycemic effect of oolong tea in type 2 diabetes. *Diabetes Care*, 26(June), 1714–1718.

Hu, F. B., Li, T. Y., Colditz, G. A., Willett, W. C., & Manson, J. E. (2003). Television watching and other sedentary behaviors in relation to risk of obesity and type 2 diabetes mellitus in women. *American Medical Association*, 289(14), 1785–1791.

Hu, F. B., Manson, J. E., Stamper, M. J., Colditz, G., Liu, S., Solomon, C. G., & Willet, W. C. (2001). Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *The New England Journal of Medicine*, 345(11), 790–797.

Kardika, I. B. W., Herawati, S., & Yasa, I. W. P. S. (2015). *Preanalitik dan interpretasi glukosa darah untuk diagnosis diabetes melitus* (pp-7-8). Bagian Patologi Klinik Fakultas Kedokteran Universitas Udayana Rumah Sakit Umum Pusat Sanglah.

Kementerian Kesehatan RI. (2014). *Buku studi diet total : survei konsumsi makanan individu Indonesia 2014* (p.143). Jakarta: Balai Penelitian dan Pengembangan Kesehatan.

Kim, J. K., So, H., Youn, M. J., Kim, H. J., Kim, Y., Park, C., Kim, S.J., Ha, Y. A., Chai, K. Y., Kim, S. M., Kim, K. Y., Park, R. (2007). *Hibiscus sabdariffa* L. water extract inhibits the adipocyte differentiation through the PI3-K and MAPK pathway. *Journal of Ethnopharmacology*, 114, 260–267.

Larasati, Y. (2010). Pengaruh pemberian seduhan kelopak rosella (*Hibiscus sabdariffa*) dosis bertingkat selama 30 hari terhadap gambaran histologik gaster tikus wistar. *Skripsi*. Universitas Diponegoro, Semarang. hal. 1-16.



Lemeshow, S., Hosmer, D. W., Klar, J., & Lwanga, S. K. (1990) *Adequacy of sample size in health studies* (First Edit). Wiley

Lemus-Mondaca, R., Vega-gálvez, A., Zura-Bravo, L., & Ah-Hen, K. (2012). *Stevia rebaudiana* Bertoni, source of a high-potency natural sweetener : A comprehensive review on the biochemical, nutritional and functional aspects. *Food Chemistry*, 132(3), 1121–1132.

Lin, T.-L., Lin, H.-H., Chen, C.-C., Lin, M.-C., Chou, M.-C., & Wang, C.-J. (2007). *Hibiscus sabdariffa* extract reduces serum cholesterol in men and women. *Nutrition Research*, 27, 140–145.

Linder, M. C. (1992). *Biokimia nutrisi dan metabolisme dengan pemakaian secara klinis*. (A. Parakkasi, Ed.) (pp. 33–37). Jakarta: UI-Press.

Mahmoud, Y. I. (2012). Effect of extract of Hibiscus on the ultrastructure of the testis in adult mice. *Acta Histochemica*, 114(4), 342–348.

Mardiah, Zakaria, F. R., Prangdimurti, E., & Damanik, R. (2015). Anti-inflammatory of purple roselle extract in diabetic rats induced by streptozotocin. *Procedia Food Science*, 3, 182–189.

Meyer, K. A., Kushi, L. H., Jacobs, D. R., Slavin, J., Sellers, T. A., & Folsom, A. R. (2000). Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am J Clin Nutr*, 71, 921–30.

Mohamed, B. B., Sulaiman, A. A., & Dahab, A. A. (2012). Roselle (*Hibiscus sabdariffa* L.) in Sudan, cultivation and their uses. *Bull. Environ. Pharmacol. Life Sci*, 1(May), 48–54.

Mohamed, S. (2014). Functional foods against metabolic syndrome (obesity, diabetes, hypertension and dyslipidemia) and cardiovascular disease. *Trends in Food Science and Technology*, 35(2), 114–128.

Mozaffari-Khosravi, H., Jalali-Khanabadi, B., Afkhami-Ardekani, M., Fatehi, F., & Noori-Shadkam, M. (2009). The effects of sour tea (*Hibiscus sabdariffa*) on hypertension in patients with type II diabetes. *Journal of Human Hypertension*, 23, 48–54.

Mozaffari-Khosravi, H., Ahadi, Z., & Tafti, M. F. (2014). The effect of green tea versus sour tea on insulin resistance, lipids profiles and oxidative stress in patients with type 2 diabetes mellitus : a randomized clinical trial. *Iran J Med Sci*, 39(5), 424–432.

Murianingrum, M. (2013). Pelepasan varietas baru rosela minuman (*Hibiscus sabdariffa* var. *sabdariffa*). Retrieved January 1, 2015, from



http://balittas.litbang.pertanian.go.id/ind/index.php?option=com_content&view=article&id=279:rosela&catid=4:info-aktual&Itemid=5

Nicholson, W. K., Asao, K., Brancati, F., Coresh, J., Pankow, J. S., & Powe, N. R. (2006). Parity and risk of type 2 diabetes. *Diabetes Care*, 29(11), 2349–2354.

NRCS. (2015). *Hibiscus sabdariffa* L. (roselle). Retrieved January 1, 2015, from <http://plants.usda.gov/core/profile?symbol=hisa2>

Onyenekwe, P. C., Ajani, E. O., Ameh, D. A., & Gamaniel, K. S. (1999). Antihypertensive effect of roselle (*Hibiscus sabdariffa*) calyx infusion in spontaneously hypertensive rats and a comparison of its toxicity with that in Wistar rats. *Cell Biochemistry and Function*, 17(3), 199–206. (Abstract).

Patel, S. (2014). *Hibiscus sabdariffa*: An ideal yet under-exploited candidate for nutraceutical applications. *Biomedicine and Preventive Nutrition*, 4(1), 23–27.

Pemerintah Kecamatan Depok Kabupaten Sleman. (2015). Peta wilayah. Retrieved May 15, 2016, from www.depokkec.sleman.go.id/profile/peta-wilayah

Peng, C. H., Chyau, C. C., Chan, K. C., Chan, T. H., Wang, C. J., & Huang, C. N. (2011). *Hibiscus sabdariffa* polyphenolic extract inhibits hyperglycemia, hyperlipidemia, and glycation-oxidative stress while improving insulin resistance. *Journal of Agricultural and Food Chemistry*, 59, 9901–9909.

Pérez-Ramírez, I. F., Castaño-Tostado, E., Ramírez-de León, J. a., Rocha-Guzmán, N. E., & Reynoso-Camacho, R. (2015). Effect of stevia and citric acid on the stability of phenolic compounds and in vitro antioxidant and antidiabetic capacity of a roselle (*Hibiscus sabdariffa* L.) beverage. *Food Chemistry*, 172, 885–892.

Perkumpulan Endokrinologi Indonesia (Perkeni). (2011). *Konsesus pengelolaan dan pencegahan diabetes melitus tipe 2 di Indonesia*. Jakarta: PB. Perkeni.

Prior, R. L., Xianli, W., Liwei, G., Hager, T. J., Hager, A., & Howard, L. R. (2008). Whole berries versus berry anthocyanins : interactions with dietary fat levels in the C57BL / 6J mouse model of obesity. *J. Agric. Food Chem*, 56, 647–653.

Raini, M., & Isnawati, A. (2011). Kajian: khasiat dan keamanan stevia sebagai pemanis pengganti gula. *Media Litbang Kesehatan*, 21(4), 145–156.



- Rehman, A., Setter, S. M., & Vue, M. H. (2011). Drug-induced glucose alterations part 2 : drug-induced hyperglycemia. *Diabetes Spectrum*, 24(4), 234–238.
- Rosemary, Rosidah, & Haro, G. (2014). Antidiabetic effect of roselle calyces extract (*Hibiscus Sabdariffa L.*) in streptozotocin induced mice. *International Journal of PharmTech Research*, 6(5), 1703–1711.
- Rosenberg, D. E., Norman, G. J., Wagner, N., Patrick, K., Calfas, K. J., & Sallis, J. F. (2010). Reliability and validity of the sedentary behavior questionnaire (SBQ) for adults. *Journal of Physical Activity and Health*, 7, 697–705.
- Sabzghabaee, A. M., Ataei, E., Kelishadi, R., Ghannadi, A., Soltani, R., Badri, S., & Shirani, S. (2013). Effect of *Hibiscus sabdariffa* calices on dyslipidemia in obese adolescents: a triple-masked randomized controlled trial. *Materia Socio-Medica*, 25(January), 76–9.
- Salas-Salvadó, J., Martínez-González, M. Á., Bulló, M., & Roz, E. (2011). The role of diet in the prevention of type 2 diabetes. *Nutrition, Metabolism and Cardiovascular Diseases*, 21, 832–848.
- Sancho, R. A. S., & Pastore, G. M. (2012). Evaluation of the effects of anthocyanins in type 2 diabetes. *Food Research International*, 46(1), 378–386.
- Sangnate, S., Wattanavijitkul, T., & Sangwatanaroj, S. (2011). Diuretic effects of roselle (*Hibiscus sabdariffa*) – Stevia (*Stevia rebaudiana*) tea compared with hydrochlorothiazide in diabetic patients with hypertension. *Thai Pharm Health Sci*, 6(4), 265–273. (Abstract).
- Sasaki, R., Nishimura, N., Hoshino, H., Isa, Y., Kadokawa, M., Ichi, T., Tanaka, A., Nushimuri, S., Fukuda, I., Ashida, H., Horio, F., Tsuda, T. (2007). Cyanidin 3-glucoside ameliorates hyperglycemia and insulin sensitivity due to downregulation of retinol binding protein 4 expression in diabetic mice. *Biochemical Pharmacology*, 74, 1619–1627.
- Sharma, R., Yadav, R., & Manivannan, E. (2012). Study of effect of *Stevia rebaudiana bertoni* on oxidative stress in type-2 diabetic rat models. *Biomedicine & Aging Pathology*, 2(3), 126–131.
- Shaw, J. E., Sicree, R. A., & Zimmet, P. Z. (2010). Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Research and Clinical Practices*, 87, 4–14.



Sinaga, E. (2012). Pengaruh pemberian susu kedelai terhadap kadar glukosa darah puasa pada wanita prediabetes. *Skripsi*. Universitas Diponegoro, Semarang. hal. 1-41.

Sindi, H. A., Marshall, L. J., & Morgan, M. R. A. (2014). Comparative chemical and biochemical analysis of extracts of *Hibiscus sabdariffa*. *Food Chemistry*, 164, 23–29.

Soewondo, P., & Pramono, L. (2011). Prevalence, characteristics, and predictors of pre-diabetes in Indonesia. *Medical Journal of Indonesia*, 25(4), 238–243.

Stull, A. J., Cash, K. C., Johnson, W. D., Champagne, C. M., & Cefalu, W. T. (2010). Bioactives in blueberries improve insulin sensitivity in obese, insulin-resistant men and women. *Journal of Nutrition*, 140(4), 1764–1768.

Tabák, A. G., Herder, C., Rathmann, W., Brunner, E. J., & Kivimäki, M. (2012). Prediabetes: a high-risk state for diabetes development. *The Lancet*, 379(9833), 2279–2290.

Tang, W., Li, S., Liu, Y., Huang, M.-T., & Ho, C.-T. (2013). Anti-diabetic activity of chemically profiled green tea and black tea extracts in a type 2 diabetes mice model via different mechanisms. *Journal of Functional Foods*, 5(4), 1784–1793.

Vue, M. H., & Setter, S. M. (2011). Drug-induced glucose alterations part 1 : drug-induced hypoglycemia. *Diabetes Spectrum*, 24(3), 171–177.

Wautier, J. L., Boulanger, E., & Wautier, M. P. (2006). Postprandial hyperglycemia alters inflammatory and hemostatic parameters. *Diabetes Metabolism*, 32, 2S34–2S36.

Whiting, D. R., Guariguata, L., Weil, C., & Shaw, J. (2011). IDF diabetes atlas : global estimates of the prevalence of diabetes for 2011 and 2030. *Diabetes Research and Clinical Practice*, 94(3), 311–321.

Wilson, R. D., & Islam, M. S. (2012). Fructose-fed streptozotocin-injected rat: an alternative model for type 2 diabetes. *Pharmacological Reports : PR*, 64(1), 129–39.