

## REFERENCES

- Ahsan Ejaz, A., Dass, B., Kambhampati, G., Ejaz, N., Maroz, N., Dhatt, G., Arif, A., Faldu, C., Lanaspas, M., Shah, G. and Johnson, R. (2012). Lowering serum uric acid to prevent acute kidney injury. *Medical Hypotheses*, 78(6), pp.796-799.
- Akira, S. and Takeda, K. (2004). Toll-like receptor signalling. *Nat Rev Immunol*, 4(7), pp.499-511.
- Ann Johnson, C., Levey, A., Coresh, J., Levin, A., Lau, J. and Eknoyan, G. (2016). Clinical Practice Guidelines for Chronic Kidney Disease in Adults: Part II. Glomerular Filtration Rate, Proteinuria, and Other Markers - American Family Physician. [online] Aafp.org. Available at: <http://www.aafp.org/afp/2004/0915/p1091.html> [Accessed 23 Mar. 2016].
- Arumugam, T., Okun, E., Tang, S., Thundyil, J., Taylor, S. and Woodruff, T. (2009). TOLL-LIKE RECEPTORS IN ISCHEMIA-REPERFUSION INJURY. *Shock*, 32(1), pp.4-16.
- Barrett, K. and Ganong, W. (2010). *Ganong's review of medical physiology*. New York: McGraw-Hill Medical.
- Basile, D., Anderson, M. and Sutton, T. (2012). Pathophysiology of Acute Kidney Injury. *Comprehensive Physiology*.
- Beck, L. and Salant, D. (2014). Membranous nephropathy: from models to man. *Journal of Clinical Investigation*, 124(6), pp.2307-2314.
- Bonventre, J. (2014). Primary proximal tubule injury leads to epithelial cell cycle arrest, fibrosis, vascular rarefaction, and glomerulosclerosis. *Kidney International Supplements*, 4(1), pp.39-44.
- Bonventre, J. and Yang, L. (2011). Cellular pathophysiology of ischemic acute kidney injury. *Journal of Clinical Investigation*, 121(11), pp.4210-4221.
- Bonventre, J., Vaidya, V., Schmouder, R., Feig, P. and Dieterle, F. (2010). Next-generation biomarkers for detecting kidney toxicity. *Nat Biotechnol*, 28(5), pp.436-440.

- Boutayeb, A. (2006). The double burden of communicable and non-communicable diseases in developing countries. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 100(3), pp.191-199.
- Coresh, J., Astor, B., McQuillan, G., Kusek, J., Greene, T., Van Lente, F. and Levey, A. (2002). Calibration and random variation of the serum creatinine assay as critical elements of using equations to estimate glomerular filtration rate. *American Journal of Kidney Disease*, 39(5), pp.920-929.
- COUTINHO, T., TURNER, S., PEYSER, P., BIELAK, L., SHEEDYII, P. and KULLO, I. (2007). Associations of Serum Uric Acid With Markers of Inflammation, Metabolic Syndrome, and Subclinical Coronary Atherosclerosis. *American Journal of Hypertension*, 20(1), pp.83-89.
- Ejerblad, E. (2006). Obesity and Risk for Chronic Renal Failure. *Journal of the American Society of Nephrology*, 17(6), pp.1695-1702.
- Eroschenko, V. and Fiore, M. (2008). *DiFiore's atlas of histology with functional correlations*. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins.
- Feneis, H., Dauber, W. (2000). "Pocket atlas of human anatomy based on the international nomenclature". Stuttgart : Thieme.
- Foundation, N. (2002). *K/DOQI clinical practice guidelines for chronic kidney disease: evaluation, classification, and stratification*. - PubMed - NCBI. [online] Ncbi.nlm.nih.gov. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11904577> [Accessed 1 Dec. 2015].
- Hall, J. and Guyton, A. (2011). *Guyton & Hall physiology review*. Philadelphia: Elsevier Saunders.
- Han, H., Lim, M., Lee, Y., Lee, J., Yang, I. and Taub, M. (2006). Uric acid inhibits renal proximal tubule cell proliferation via at least two signaling pathways involving PKC, MAPK, cPLA2, and NF-kB. *AJP: Renal Physiology*, 292(1), pp.F373-F381.
- Johnson, R., Nakagawa, T., Sanchez-Lozada, L., Shafiu, M., Sundaram, S., Le, M., Ishimoto, T., Sautin, Y. and Lanasa, M. (2013). Sugar, Uric Acid, and the Etiology of Diabetes and Obesity. *Diabetes*, 62(10), pp.3307-3315.

- Kang, D. (2002). A Role for Uric Acid in the Progression of Renal Disease. *Journal of the American Society of Nephrology*, 13(12), pp.2888-2897.
- Kawasaki, T. and Kawai, T. (2014). Toll-Like Receptor Signaling Pathways. *Front. Immunol.*, 5.
- Khosla, U., Zharikov, S., Finch, J., Nakagawa, T., Roncal, C., Mu, W., Krotova, K., Block, E., Prabhakar, S. and Johnson, R. (2005). Hyperuricemia induces endothelial dysfunction. *Kidney International*, 67(5), pp.1739-1742.
- Kuehnelt, W. (2003). *Color atlas of cytology, histology, and microscopic anatomy*. Stuttgart: Thieme.
- Liu, Y 2006, 'Renal fibrosis: new insights into the pathogenesis and therapeutics', *Kidney international*, vol. 69, no. 2, pp. 213-217.
- Longo, D. and Harrison, T. (2011). *Harrison's principles of internal medicine*. Maidenhead: McGraw-Hill.
- López-Novoa, J., Rodríguez-Peña, A., Ortiz, A., Martínez-Salgado, C. and López Hernández, F. (2011). Etiopathology of chronic tubular, glomerular and renovascular nephropathies: Clinical implications. *Journal of Translational Medicine*, 9(1), p.13.
- Martin, LJ. (2015). *Uric acid - blood : MedlinePlus Medical Encyclopedia*. [online] Nlm.nih.gov. Available at: <https://www.nlm.nih.gov/medlineplus/ency/article/003476.htm> [Accessed 1 Dec. 2015].
- Mazzali, M., Kanellis, J., Han, L., Feng, L., Xia, Y., Chen, Q., Kang, D., Gordon, K., Watanabe, S., Nakagawa, T., Lan, H., and Johnson, R. (2002). Hyperuricemia induces a primary renal arteriopathy in rats by a blood pressure-independent mechanism. *American Journal of Physiology - Renal Physiology*, 282(6), pp.F991-F997.
- Moreth, K., Frey, H., Hubo, M., Zeng-Brouwers, J., Nastase, M., Hsieh, L., Hacen, R., Pfeilschifter, J., Iozzo, R. and Schaefer, L. (2014). Biglycan-triggered TLR-2- and TLR-4-signaling exacerbates the pathophysiology of ischemic acute kidney injury. *Matrix Biology*, 35, pp.143-151.
- Murray, R. (2009). *Harper's illustrated biochemistry*. New York: McGraw-Hill Medical.

- Nacak, H., van Diepen, M., de Goeij, M., Rotmans, J., and Dekker, F. (2014). Uric acid: association with rate of renal function decline and time until start of dialysis in incident pre-dialysis patients. *BMC Nephrol*, 15(1), p.91.
- Netea, M., Kullberg, B., Blok, W., Netea, R. and Van der Meer, J. (1997). The role of hyperuricemia in the increased cytokine production after lipopolysaccharide challenge in neutropenic mice. *Blood Journal*, 89(2), pp.577 - 582.
- Obermayr, R., Temml, C., Gutjahr, G., Knechtelsdorfer, M., Oberbauer, R. and Klauser-Braun, R. (2008). Elevated Uric Acid Increases the Risk for Kidney Disease. *Journal of the American Society of Nephrology*, 19(12), pp.2407-2413.
- Perrone, RD, Madias, NE & Levey, AS. (1992). Serum creatinine as an index of renal function: New insights into old concepts, *Clinical Chemistry*, vol. 38, no. 10, pp. 1933-1953.
- Piccinini A. and Midwood, K. (2010). DAMPening Inflammation by Modulating TLR Signalling. *Mediators of Inflammation*, 2010, pp.1-21.
- Prasad Sah, O. and Qing, Y. (2015). Associations between Hyperuricemia and Chronic Kidney Disease: A Review. *Nephro-Urology Monthly*, 7(3).
- Pulskens, W., Rampanelli, E., Teske, G., Butter, L., Claessen, N., Luijck, I., van der Poll, T., Florquin, S. and Leemans, J. (2010). TLR4 Promotes Fibrosis but Attenuates Tubular Damage in Progressive Renal Injury. *Journal of the American Society of Nephrology*, 21(8), pp.1299-1308.
- Rosin, D. and Okusa, M. (2011). Dangers Within: DAMP Responses to Damage and Cell Death in Kidney Disease. *Journal of the American Society of Nephrology*, 22(3), pp.416-425.
- Saladin, K., Widmaier, E., Raff, H., Strang, K. and Kardong, K. (2011). *Evolutionary human anatomy and physiology*. Boston: McGraw-Hill Higher Education.
- Samra, M. (2012). False Estimates of Elevated Creatinine. *permj*, 16(2).

- Sanchez-Lozada, L., Tapia, E., Lopez-Molina, R., Nepomuceno, T., Soto, V., Avila-Casado, C., Nakagawa, T., Johnson, R., Herrera-Acosta, J. and Franco, M. (2006). Effects of acute and chronic L-arginine treatment in experimental hyperuricemia. *AJP: Renal Physiology*, 292(4), pp.F1238-F1244.
- Sanchez-Lozada, L., Soto, V., Tapia, E., Avila-Casado, C., Sautin, Y., Nakagawa, T., Franco, M., Rodriguez-Iturbe, B., and Johnson, R. (2008). Role of oxidative stress in the renal abnormalities induced by experimental hyperuricemia. *AJP: Renal Physiology*, 294(4), pp.F1134-F1141.
- Smeets, B., Boor, P., Dijkman, H., Sharma, S., Jirak, P., Mooren, F., Berger, K., Bornemann, J., Gelman, I., Floege, J., van der Vlag, J., Wetzels, J. and Moeller, M. (2013). Proximal tubular cells contain a phenotypically distinct, scattered cell population involved in tubular regeneration. *J. Pathol.*, 229(5), pp.645-659.
- Stevens, L., Coresh, J., Greene, T. and Levey, A. (2006). Assessing Kidney Function "Measured and Estimated Glomerular Filtration Rate. *New England Journal of Medicine*, 354(23), pp.2473-2483.
- The National Kidney Foundation. (2015). Global Facts: About Kidney Disease. [online] Available at: <https://www.kidney.org/kidneydisease/global-facts-about-kidney-disease> [Accessed 23 Mar. 2016].
- Tortora, G. and Derrickson, B. (2012). *Principles of anatomy & physiology*. Hoboken, NJ: Wiley.
- Traynor, J., Mactier, R., Geddes, C. and Fox, J. (2006). How to measure renal function in clinical practice. *BMJ*, 333(7571), pp.733-737.
- Wang, J., Zhou, Y., Xu, M., Rong, R., Guo, Y. and Zhu, T. (2011). Urinary Metabolomics in Monitoring Acute Tubular Injury of Renal Allografts: A Preliminary Report. *Transplantation Proceedings*, 43(10), pp.3738-3742.
- Wu, H., Chen, G., Wyburn, K., Yin, J., Bertolino, P., Eris, J., Alexander, S., Sharland, A. and Chadban, S. (2007). TLR4 activation mediates kidney ischemia/reperfusion injury. *Journal of Clinical Investigation*, 117(10), pp.2847-2859.

- Xi, Y., Shao, F., Bai, X., Cai, G., Lv, Y. and Chen, X. (2014). Changes in the Expression of the Toll-Like Receptor System in the Aging Rat Kidneys. *PLoS ONE*, 9(5), p.e96351.
- Yamamoto, M. and Takeda, K. (2010). Current Views of Toll-Like Receptor Signaling Pathways. *Gastroenterology Research and Practice*, 2010, pp.1-8.
- Zhang, B., Ramesh, G., Uematsu, S., Akira, S. and Reeves, W. (2008). TLR4 Signaling Mediates Inflammation and Tissue Injury in Nephrotoxicity. *Journal of the American Society of Nephrology*, 19(5), pp.923-932.