



## REFERENCES

- Abrams, M., Hook, S., & Ramachandran, B. (1999). *ASTER User Handbook (Ver.2)*. California: Jet Propulsion Laboratory, California Institute of Technology.
- Armstrong, R. A. (1993). Remote sensing of submerged vegetation canopies for biomass estimation. *International Journal of Remote Sensing*, 14, 10-16.
- Asner, G. P., Scurlock, J. M., & Hicke, J. A. (2003). Global synthesis of leaf area index observations: Implications for ecological and remote sensing studies. *Global Ecology and Biogeography*, 12, 191 – 205.
- As-syakur, R., Osawa, T., & Adnyana, I. W. (2010). Medium Spatial Resolution Satellite Imagery to Estimate Gross Primary Production in an Urban Area. *Remote Sensing*, 2, 1496-1507.
- Bajracharya, S. (2008). *Community Carbon Forestry: Remote Sensing of Forest Carbon and Forest Degradation in Nepal*. The Netherlands: International Institute for Geo-information Science and Earth Observatio, ITC.
- Birth, G. S., & McVey, G. (1968). Measuring the color of growing turf with a reflectance spectrophotometer. *Agronomy Journal*, 66, 640-643.
- Björk, M., Short, F., Mcleod, E., & Beer, S. (2008). *Managing seagrasses for resilience to climate change*. Gland, Switzerland: IUCN.
- Boonsong, K., Piyatiratitivorakul, S., & Patanaponpaiboon, P. (2003). Potential use of mangrove plantation as constructed wetland for municipal wastewater treatment. *Water Science and Technology*, 48, 257-266.
- Borum, J. (2003). *Global climate change and the future of seagrass meadows*. Copenhagen: Freshwater Biological Laboratory, University of Copenhagen.
- Bouillon, S., Borges, A., Castañeda-Moya, E., Diele, K., Dittmar, T., Duke, N., et al. (2008). Mangrove production and carbon sinks: A revision of global Budget estimates. *Global Biogeochemical Cycles*, 22.
- Broge, N. H., & Leblanc, E. (2000). Comparing prediction power and stability of broadband and hyperspectral vegetation indices for estimation of green leaf area index and canopy chlorophyll density. *Remote Sensing of Environment*, 76, 156-172.
- Bukata, R. P., Jerome, J. H., Kondratyev, K. Y., & Pozdnyakov, D. V. (1995). *Optical Properties and Remote Sensing of Inland and Coastal Waters*. New York: CRC.
- Burdige, D. G., Zimmerman, R. C., & Hu, X. (2008). Rates of carbonate dissolution in permeable sediments estimated from pore-water profiles: the role of seagrasses. *Limnology Oceanography*, 53, 549-565.
- Capolsini, P., Andrefouet, S., Rion, C., & Payri, C. (2003). A Comparison of Landsat ETM+, SPOT HRV, IKONOS, ASTER and airborne MASTER Data for Coral Reef Habitat Mapping in South Pacific Island. *Canadian Journal of Remote Sensing*, 29 (2), 187-200.
- Cebrian, J., & Duarte, C. M. (1996). Plant growth-rate dependence of detrital carbon storage in ecosystems. *Science*, 268, 1606-1608.



- Ceccato, P., Flasse, S., Tarantola, S., Jacquemond, S., & Gregoire, J. M. (2001). Detecting vegetation water content using reflectance in the optical domain. *Remote Sensing of Environment*, 77, 22–33.
- Charphy-Roubaud, C., & Sournia, A. (1990). The comparative estimation of phytoplanktonic microphytobenthic production in the oceans. *Mar. Micr. Food Webs*, 4, 31–57.
- Chave, J., Andalo, C., Brown, S., Cairns, M. A., Chambers, J. Q., Eamus, D., et al. (2005). Tree allometry and improved estimation of carbon stocks and balance in tropical forests. *Oecologia*, 145, 87–99.
- Chavez Jr, P. (1996). Image-based atmospheric corrections — Revisited and improved. *Photogramm. Eng. Remote Sensing*, 62, 1025–1036.
- Chavez, P., Berlin, G., & Mitchell, W. (1977). Computer Enhancement Techniques of Landsat MSS Digital Images for Landuse/Landcover Assessments. *Remote Sensing of Earth Resources*, 6, 259.
- Chen, C. T. (2009). Reconciling Opposing Views on Carbon Cycling in the Coastal Ocean: Continental Shelves as Sinks and Near-shore Ecosystems as Sources of Atmospheric CO<sub>2</sub>. *Deep Sea Research II* 2009, 56, 578–590.
- Chen, J. M., Pavlic, G., Brown, L., Cihlar, J., Leblanc, S., White, H., et al. (2002). Derivation and validation of Canada-wide coarse-resolution leaf area index maps using high resolution satellite imagery and ground measurement. *Remote Sensing of Environment*, 80, 165–184.
- Chen, X., Vierling, L., & Deering, D. (2005). A simple and effective radiometric correction method to improve landscape change detection across sensors and across time. *Remote Sensing of Environment*, 98, 63 – 79.
- Chen, X., Vierling, L., Rowell, E., & DeFelice, T. (2004). Using lidar and effective LAI data to evaluate IKONOS and Landsat 7 ETM+ vegetation cover estimates in a ponderosa pine forest. *Remote Sensing of Environment*, 91, 14–26.
- Clough, B. F., & Scott, K. (1989). Allometric relationships for estimating aboveground biomass in six mangrove species. *Forest Ecology Management*, 27, 117–127.
- Coles, R. G., McKenzie, L. J., Rasheed, M. A., Mellors, J. E., Taylor, H., Dew, K., et al. (2007). *Status and Trends of Seagrass Habitats in the Great Barrier Reef World Heritage Area. Report to the Marine and Tropical Sciences Research Facility*. Cairns: Reef and Rainforest Research Centre Ltd.
- Comley, B. w., & McGuinness, K. A. (2005). Above- and below-ground biomass, and allometry of four common northern Australian mangroves. *Australian Journal of Botany*, 53, 431–436.
- Dekker, A., Brando, V., Anstee, J., Fyfe, S. K., Malthus, T., & Karpouzli, E. (2006). Remote sensing of seagrass systems: Use of spaceborne and airborne systems. In A. Larkum, R. Orth, C. Duarte, & C. Duarte (Ed.), *Seagrasses: Biology, Ecology and Conservation* (p. 347–359). Dordrecht: Springer.
- Delbart, N., Kergoat, L., Toan, T. L., Lhermitte, J., & Picard, G. (2005). Determination of phenological dates in boreal regions using normalized difference water index. *Remote Sensing of Environment*, 97, 26–38.



- Dennison, W. (1990). Shoot density. Chap. 9. In: Seagrass research methods. In R. Phillips, C. McRoy, & C. P. McRoy (Ed.), *Monographs on Oceanographic Methodology* (pp. 61-63). Paris: UNESCO.
- Dierssen, H. M., Zimmerman, R. C., Drake, L. A., & Burdige, D. (2010). Benthic ecology from space: optics and net primary production in seagrass and benthic algae across the Great Bahama Bank. *Marine Ecology Progress Series* , 411, 1-15.
- Duarte, C. M., Middelburg, J., & Caraco, N. (2005). Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences* , 2, 1-8.
- Edwards, A. (2004). Assessing Mangrove LAI using CASI Airborne Imagery. *Bilko Tutorial on the Applications of satellite and airborne image data to coastal management* . UNESCO.
- Edwards, A. (2004). Geometric Correction of an Aerial Image of South Caicos Island. *Bilko Tutorial on the Applications of satellite and airborne image data to coastal management* . Paris: UNESCO.
- Eitel, J., Long, D., Gessler, P., & Hunt, E. (2008). Combined spectral index to improve ground-based estimates of nitrogen status in dryland wheat. *Agronomy Journal* , 100, 1694-1702.
- Eveleth, R. (2010). *Seagrass: A Potential Carbon Sink (Report ENVR 102)*. UNEP.
- FAO. (2010). *Global Forest Resources Assessment 2010 - Main Report*. Rome: FAO.
- FAO. (2007). *The world's mangroves 1980-2005. FAO Forestry Paper 153*. Rome: Food and Agriculture Organization of the United Nations.
- Fatoyinbo, T. E., & Armstrong, A. H. (2010). Remote Characterization of Biomass Measurements: Case Study of Mangrove Forests. In M. Momba, & F. Bux (Eds.), *Biomass*. Croatia: Sciyo.
- Fatoyinbo, T., Simard, M., & Washington-Allen, R. A. (2008). Landscape scale height, biomass and carbon estimation of mangrove forests with Shuttle Radar Topography Mission elevation data. *Journal for Geophysical Research-Biogeosciences* , 113.
- Fourqurean, J. (2014, October 21). Estimates of Seagrass Carbon Storage - Indonesia. Rio Grande, Brasil: The Blue Carbon Initiative, International Scientific Working Group Meeting.
- Fourqurean, J. W., Duarte, C. M., Kennedy, H., Marbà, N., Holmer, M., Mateo, M. A., et al. (2012). Seagrass ecosystems as a globally significant carbon stock. *Nature Geoscience* , 5, 505-509.
- Frazer, G. W., Canham, C. D., & Lertzman, K. P. (1999). *Gap Light Analyzer (GLA), Version 2.0: Imaging software to extract canopy structure and gap light transmission indices from true-color fisheye photographs, users manual and program documentation*. Burnaby, British Columbia - Millbrook, New York: Simon Fraser University & the Institute of Ecosystem Studies.
- Fyfe, S. K. (2003). Spatial and temporal variation in spectral reflectance: Are seagrass species spectrally distinct? *Limnology and Oceanography* , 48 (1, part 2), 464-479.
- Fyfe, S. K., & Dekker, A. G. (2001). Seagrass species: are they spectrally distinct? *Proceedings IEEE 2001 International Geoscience and Remote Sensing*, 2746, pp. 2740-2742. Piscataway, NJ.



- Gao, B. C. (1996). NDWI - A normalized difference water index for remote sensing of vegetation liquid water from space. *Remote Sensing of Environment* , 58, 257-266.
- Giri, C., Tieszen, L. L., Zhu, Z., Singh, A., Loveland, T., Masek, J., et al. (2010). Status and Distribution of Mangrove Forest of the World using Earth Observation Satellite Data. *Global Ecology and Biogeography* , 1-6.
- Gitelson, A. A. (2003). Wide Dynamic Range Vegetation Index for Remote Quantification of Biophysical Characteristics of Vegetation. *Plant Physiology* , 161, 165–173.
- Gitelson, A. A., Kaufman, Y. J., Stark, R., & Rundquist, D. (2002). Novel algorithms for remote estimation of vegetation fraction. *Remote Sensing of Environment* , 80, 76–87.
- Gobron, N. (2008). *Leaf Area Index*.
- Goodman, J. A., Lee, Z., & Ustin, S. L. (2008). Influence of Atmospheric and Sea-Surface Corrections on Retrieval of Bottom Depth and Reflectance Using a Semi-Analytical Model: A Case Study in Kaneohe Bay, Hawaii. *Appl. Opt* , 47, 1-11.
- Gosa, A. G. (2009). *Remote sensing of leaf area index: enhanced retrieval from close-range and remotely sensed optical observations*. Helsinki: Hansaprint – Hansa Direct 2009.
- Green, E. P., & Mumby, P. J. (2000). Mapping Mangroves. In A. J. Edwards (Ed.), *Remote Sensing Handbook for Tropical Coastal Management* (p. 183–192). Paris: UNESCO.
- Green, E. P., & Short, F. T. (Eds.). (2003). *World Atlas of Seagrasses*. Los Angeles: University of California Press.
- Green, E. P., Mumby, P. J., Edwards, A. J., & Clark, C. D. (2000). *Remote Sensing Handbook for Tropical Coastal Management*. *Coastal Management Sourcebooks 3*. (A. J. Edwards, Ed.) Paris: UNESCO.
- Griffin, M. K., & Burke, H. K. (2003). Comparison of hyperspectral data for atmospheric Effects. *Lincoln Laboratory Journal* , 14, 29–54.
- Gu, Y., Hunt, E., Wardlow, B., Basara, J., Brown, J., & Verdin, J. (2008). Evaluation of MODIS NDVI and NDWI for vegetation drought monitoring using Oklahoma Mesonet soil moisture data. *Geophysical Research Letters* , 35.
- Hadjimitsis, D., Papadavid, G., Agapiou, A., Themistocleous, K., Hadjimitsis, M., Retalis, A., et al. (2010). Atmospheric correction for satellite remotely sensed data intended for agricultural applications: impact on vegetation indices. *Nat. Hazards Earth Syst. Sci* , 10, 89–95.
- Hedley, J. D., Harborne, A. R., & Mumby, P. J. (2005). Simple and Robust Removal of Sunlight for Mapping Shallow-Water Benthos. *Journal of Remote Sensing* , 26 (10), 2107-2112.
- Hedley, J. D., Roelfsema, C. M., Phinn, S. R., & Mumby, P. J. (2012). Environmental and sensor limitations in optical remote sensing of coral reefs: implications for monitoring and sensor design. *Remote Sensing* , 4, 271-302.
- Heumann, B. W. (2011). An Object-Based Classification of Mangroves Using a Hybrid Decision Tree—Support Vector Machine Approach. *Remote Sensing* , 3, 2440-2460.
- Hochberg, E., Andrefouet, S., & Tyler, M. (2003). Sea Surface Correction of High Spatial Resolution Ikonos Images to Improve Bottom Mapping in Near-Shore Environments. *IEEE Trans. Geosci. Remote Sens* , 41, 1724-1729.



- Huete, A. R., & Liu, H. Q. (1994). An Error and Sensitivity Analysis of the Atmospheric and Soil-Correcting Variants of the NDVI for the MODIS-EOS. *IEEE Transaction on Geoscience and Remote Sensing* , 32 (4), 897-905.
- Huete, A. R., Hua, G., Qi, J., Chehbouni, A., & Ven Leeuwem, W. J. (1992). Normalization of Multidirectional Red and Near-infrared Reflectances with the SAVI. *Remote Sensing of Environment* , 40, 1-20.
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment* , 83, 195–213.
- Inoue, Y., Peñuelas, J., Miyata, A., & Mano, M. (2008). Normalized Difference Spectral Indices for Estimating Photosynthetic Hyperspectral and CO<sub>2</sub> Flux Measurements in Rice. *Remote Sensing of Environment* , 112, 156 – 172.
- IPCC. (2003). *IPCC Good Practice Guidance for Land Use, Land-Use Change and Forestry Prepared by the National Greenhouse Gas Inventories Programme*. (J. Penman, M. Gytarsky, & T. Hiraishi, Eds.) Hayama: IGES.
- IPCC. (2006). *IPCC Guidelines for National Greenhouse Gas Inventories*. Japan: IPCC.
- Janzen, H. H. (2004). Carbon cycling in earth systems – a soil science perspective. *Agriculture ecosystems and Environment* , 104, 399-417.
- Jardine, S. L., & Siikamaki, J. V. (2014). A Global Predictive Model of Carbon in Mangrove Soils. *Environmental Research Letters* , 9, 1-9.
- Jennerjahn, T. C., & Ittekkot, V. (2002). Relevance of mangroves for the production and deposition of organic matter along tropical continental margins. *Die Naturwissenschaften* , 89 (1), 23-30.
- Jensen, J. R. (2006). *Remote Sensing of the Environment. An Earth Resource Perspective*. (K. E. Clarke, Ed.) Prentice Hall.
- Jiang, Z., Huete, A. R., Didan, K., & Miura, R. (2008). Development of a two-band enhanced vegetation index without a blue band. *Remote Sensing of the Environment* , 112 (10), 3833-3845.
- Jupp, D. L. (1988). Background and extensions to depth of penetration (DOP) mapping in shallow coastal waters. *Proceedings of the Symposium on Remote Sensing of the Coastal Zone*. Gold Coast, Queensland.
- Kamal, M., & Phinn, S. (2011). Hyperspectral Data for Mangrove Species Mapping: A Comparison of Pixel-Based and Object-Based Approach. *Remote Sensing* , 3, 2222-2242.
- Kapp, G., Agus, F., Hairiah, K., Verlarde, S., & Noordwijk, M. v. (2011). Carbon Measurement [workshop presentation]. REDD+.
- Karnieli, A., Kaufman, Y. J., Remer, L., & Wald, A. (2001). AFRI: Aerosol Free Vegetation Index. *Remote Sensing of Environment* , 77, 10-21.
- Kaufman, Y. J., & Tanre, D. (1992). Atmospherically Resistant Vegetation Index (ARVI) for EOS-MODIS. *IEEE Transaction on Geoscience and Remote Sensing* , 30 (2), 261-270.



- Kay, S., D, H. J., & Lavender, S. (2009). Sun Glint Correction of High and Low Spatial Resolution Images of Aquatic Scenes: a Review of Methods for Visible and Near-Infrared Wavelengths. *Remote Sensing* , 1, 697-730.
- Kiage, L. M., & Walker, N. D. (2008). Using NDVI from MODIS to Monitor Duckweed Bloom in Lake Maracaibo, Venezuela. *Water Resource Management* .
- Knudby, A., & Nordlund, L. (2011). Remote sensing of seagrass in a patchy multi-species environment. *International Journal of Remote Sensing* , 32 (8), 2227-2244.
- Knyazikhin, Y., Glassy, J., L, P. J., Tian, Y., Lotsch, A., Zhang, Y., et al. (1999). *MODIS Leaf Area Index (LAI) and Fraction of Photosynthetically Active Radiation Absorbed by Vegetation (FPAR) Product (MOD15) Algorithm Theoretical Basis Document*. NASA.
- Komiyama, A., Ong, J., & Pongpam, S. (2008). Allometry, biomass, and productivity of mangrove forests: A review. *Aquatic Botany* , 89, 128-137.
- Komiyama, A., Pongpam, S., & Kato, S. (2005). Common allometric equations for estimating the tree weight of mangroves. *Journal of Tropical Ecology* , 21, 471-477.
- Kotchenova, S. Y., Vermote, E. F., Matarrese, R., & Klemm Jr, F. J. (2006). Validation of a vector version of the 6S radiative transfer code for atmospheric correction of satellite data. Part I: Path Radiance. *Applied Optics* , 45 (26), 6726-6774.
- Kruse, F. A. (2004). Comparison of ATREM, ACORN, and FLAASH atmospheric corrections using low-altitude AVIRIS data of Boulder, CO. *Summaries of 13th JPL Airborne Geoscience Workshop*. Pasadena, CA: Jet Propulsion Lab.
- Kuenzer, C., Bluemel, A., Gebhardt, S., Vo Quoc, T., & Dech, S. (2011). Remote Sensing of Mangrove Ecosystems: A Review. *Remote Sensing* , 3, 878-928.
- Laffoley, D., & Grimsditch, G. (2009). *The management of natural coastal carbon sinks*. (G. Grimsditch, Ed.) Gland, Switzerland: IUCN.
- Lathrop, R. G., Montesano, P., & Haag, S. (2006). A Multi-scale Segmentation Approach to Mapping Seagrass Habitats Using Airborne Digital Camera Imagery. *Photogrammetric Engineering & Remote Sensing* , 72 (6), 665-675.
- Leprieur, C., Kerr, Y. H., & Pichon, J. M. (1996). Critical assessment of vegetation indices from AVHRR in a semi-arid environment. *International Journal of Remote Sensing* , 17, 2549-2563.
- Li, A., Deng, W., Liang, S., & Huang, C. (2010). Investigation on the Patterns of Global Vegetation Change Using a Satellite-Sensed Vegetation Index. *Remote Sensing* , 2, 1530-1548.
- Li, X., Gar-on Yeh, A., Wang, S., Liu, K., Liu, X., Qian, J., et al. (2007). Regression and analytical models for estimating mangrove wetland biomass in South China using Radarsat images. *International Journal of Remote Sensing* , 28 (24), 5567-5582.
- Liu, G. R., Liang, C. K., Kuo, T. H., Lin, T. H., & Huang, S. J. (2004). Comparison of the NDVI, ARVI and AFRI Vegetation Index, Along with Their Relations with the AOD Using SPOT 4 Vegetation Data. *TAO* , 15 (1), 15-31.
- Lotze, H. K., Lenihan, H. S., Bourque, B. J., Bradbury, R. H., Cooke, R. G., Kay, M. C., et al. (2006). Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science* , 312, 1806-1809.



- Lucas, R. M., Mitchell, A. L., Rosenqvist, A., Proisy, C., Melius, A., & Ticehurst, C. (2007). The potential of L-band SAR for quantifying mangrove characteristics and change: case studies from the tropics. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 17, 245.
- Lyon, J. G., Yuan, D., Lunetta, R. S., & Elvidge, C. D. (1998). A Change Detection Experiment Using Vegetation Indices. *Photogrammetric Engineering & Remote Sensing*, 64 (2), 143-150.
- Lyons, M. B., Phinn, S. R., & Roelfsema, C. M. (2012). Long term land cover and seagrass mapping using Landsat and object-based image analysis from 1972 to 2010 in the coastal environment of South East Queensland, Australia. *ISPRS Journal of Photogrammetry and Remote Sensing*, 71, 34-46.
- Lyons, M., Phinn, S., & Roelfsema, C. (2011). Integrating Quickbird Multi-Spectral Satellite and Field Data: Mapping Bathymetry, Seagrass Cover, Seagrass Species and Change in Moreton Bay, Australia in 2004 and 2007. *Remote sensing*, 3, 42-64.
- Lyzenga, D. R. (1978). Passive Remote-Sensing Techniques for Mapping Water Depth and Bottom Features. *Applied Optics*, 17, 379-383.
- Lyzenga, D., Malinas, N., & Tanis, F. (2006). Multispectral Bathymetry Using a Simple Physically Based Algorithm. *IEEE Trans. Geosci. Remote Sens*, 44, 2251-2259.
- Manila, M. R. (2007, December). Forest Canopy Density Mapping for Forest Climate Change Mitigation, FLR and SFM: Philippine Experience. Nusa Dua, Bali, Indonesia: Paper Presented at ITTO Side Event at COP 13 of UNFCCC on Forest Day.
- Mateo, M. A., Cebrian, J., Dunton, K., & Mutchler, T. (2006). Carbon flux in seagrass ecosystems. In W. D. Larkum, R. J. Orth, & C. M. Duarte (Eds.), *Seagrasses: Biology, Ecology and Conservation* (pp. 567-593). Springer.
- Mateo, M. A., Romero, J., Pérez, Littler, M. M., & Littler, D. S. (1997). Dynamics of Millenary Organic Deposits Resulting from the Growth of the Mediterranean Seagrass *Posidonia oceanica*. *Estuarine, Coastal and Shelf Science*, 44, 103-110.
- McCoy, R. M. (2005). *Field Methods in Remote Sensing*. New York: The Guilford Press.
- McKenzie, L., Finkbeiner, M. A., & Kirkman, H. (2001). Seagrass Mapping Methods. In F. T. Short, & R. G. Coles, *Global Seagrass Research Methods*. Amsterdam: Elsevier.
- Meyer, C. A., & Pu, R. (2011). Seagrass resource assessment using remote sensing methods in St. Joseph Sound and Clearwater Harbor, Florida, USA. *Environmental Monitoring Assessment*, 184 (2), 1131-1143.
- Mobley, C. D. (1994). *Light and water: Radiative transfer in natural waters*. Sand Diego: Academic Press.
- Mougin, E., Proisy, C., Marty, G., Fromard, F., Puig, H., Betoulle, J. L., et al. (1999). Multifrequency and multipolarisation radar backscattering from mangrove forests. *IEEE Transactions on Geoscience and Remote Sensing*, 37, 94-102.
- Mumby, P. J., & Green, E. (2000). Field survey: Building the link between image and reality. In A. J. Edwards (Ed.), *Remote Sensing Handbook for Tropical Coastal Management* (pp. 57-66). Paris: UNESCO.



- Mumby, P. J., Green, E. P., Edwards, A. J., & Clark, C. D. (1997). Measurement of seagrass standing crop using satellite and digital airborne remote sensing. *Marine Ecology Progress Series*, 159, 51–60.
- Murdiyarso, D., Donato, D., Kauffman, J., Kurnianto, S., Stidham, M., & Kanninen, M. (2009). *Carbon storage in mangrove and peatland ecosystem: A preliminary account from plots in Indonesia*. Center for International Forestry Research: Bogor.
- Murdiyarso, D., van Noordwijk, M., Wasrin, U., Tomich, T., & Gillison, A. (2002). Environmental benefits and sustainable land-use in Jambi transect, Sumatra, Indonesia. *Journal of Vegetation Science*, 13, 429–438.
- Murphy, B., & Cowra. (2008, November). How do we estimate soil carbon levels? Making best use of existing science and knowledge [workshop presentation]. Department of Environment & Climate Change NSW.
- Nagelkerken, I., Blaber, S., Bouillon, S., Green, P., Haywood, Kirton, L. G., et al. (2008). The habitat function of mangroves for terrestrial and marine fauna: a review. *Aquatic Botany*, 89, 155-185.
- Nellemann, C., Corcoran, E., Duarte, C. M., Valdes, L., De Young, C., Fonseca, L., et al. (2009). *Blue Carbon. A Rapid Response Assessment*. (G. Grimsditch, Ed.) United Nations Environment Programme, GRID-Arendal.
- Oliver, J. K., Berkelmans, R., & Eakin, C. M. (2009). Coral Bleaching in Space and Time. In M. H. Van Oppen, & J. M. Lough (Eds.), *Coral Bleaching: Patterns, Processes, Causes and Consequences* (p. 178). Berlin: Ecological Studies 205, Springer.
- Ong, J. E. (2002). The Hidden Costs of Mangrove Services: Use of Mangroves for Shrimp Aquaculture. *International Science Roundtable for the Media*. Bali, Indonesia: ICSU, IGBP, IHDP, WCRP, DIVERSITAS, START.
- Ong, J. E., Gong, W. K., & Wong, C. H. (2004). Allometry and partitioning of the mangrove, *Rhizophora apiculata*. *Forest Ecology Management*, 188, 395–408.
- Oohata, S., & Shinozaki, K. (1979). A statistical model of plant form - further analysis of the pipe model theory. *Japanese Journal of Ecology*, 29, 323–335.
- Orth, R., Carruthers, T., Dennison, W., Duarte, C., Fourqurean, J., Heck Jr., K., et al. (2006). A Global Crisis for Seagrass Ecosystems. *Bioscience*, 56 (12), 987-996.
- Palacios, S. L., & Zimmerman, R. C. (2007). Response of eelgrass *Zostera marina* to CO<sub>2</sub> enrichment: possible impacts of climate change and potential for remediation of coastal habitats. *Marine Ecology Progress Series*, 344, 1-13.
- Parwati, E., Kartika, T., Indarto, J., A.K. Dyah, F., Nur, M., & Kartasmita, M. (2007). Kajian Hubungan Antara Laju Perubahan TSS (Total Suspended Solid) Dengan Penutup / Penggunaan Lahan Di Wilayah Pesisir Kabupaten Berau, Kalimantan Timur. *Proceeding Geo-Marine Research Forum 2007*, (pp. 323-338).
- Pasqualini, V., Pergent-Martini, C., Pergent, G., Agreil, M., Skoufas, G., Sourbes, L., et al. (2005). Use of SPOT 5 for mapping seagrasses: An application to *Posidonia oceanica*. *Remote Sensing of Environment*, 94, 39-45.



- Pendleton, L., Donato, D., Murray, B., Crooks, S., Jenkins, W., Sifleet, S., et al. (2012). Estimating Global “Blue Carbon” Emissions from Conversion and Degradation of Vegetated Coastal Ecosystems. *PlosOne* , 7 (9).
- Pergent, G., Romero, J., Pergent-Martini, C., Mateo, M. A., & Boudouresque, C. F. (1994). Primary production, stocks and fluxes in the Mediterranean seagrass *Posidonia oceanica*. *Marine Ecology Progress Series* , 106, 139- 146.
- Phinn, S. R., Roelfsema, C. M., Brando, V., & Anstee, J. (2008). Mapping seagrass species, cover and biomass in shallow waters: An assessment of satellite multi-spectral and airborne hyper-spectral imaging systems in Moreton Bay (Australia). *Remote Sensing of Environment* , 112, 3413–3425.
- Phinn, S., & Dekker, A. G. (2004). *An integrated remote sensing approach for adaptive Management of complex coastal waters: The Moreton Bay Case Study*. Queensland: Cooperative Research Centre for Coastal Zone, Estuary and Waterway Management.
- Pinty, B., & Verstraete, M. M. (1992). GEMI: A non-linear index to monitor global vegetation from satellites. *Vegetatio* , 101, 15-20.
- Potitsep, S., Nasahara, N. K., & Suzuki, R. (2010). Comparison of LAI estimation based vegetation index and MODIS LAI for gross primary productivity estimation, deciduous broadleaf forest in Japan. *Geophysical Research Abstracts* , 12.
- Poungparn, S., Komiyama, A., Jintana, V., Piriyaota, S., Sangtiew, T., Tanapermpool, P., et al. (2001). A quantitative analysis on the root system of a mangrove, *Xylocarpus granatum* Koenig. *Tropics* , 12, 35–42.
- Proisy, C., Mougin, E., Fromand, F., Trichon, V., & Karam, M. A. (2002). On the influence of canopy structure on the polarimetric radar response from mangrove forest. *International Journal of Remote Sensing* , 23, 4197–4210.
- Proisy, C., Mougin, E., Fromard, F., & Karam, M. A. (2000). Interpretation of polarimetric radar signatures of mangrove forests. *Remote Sensing of Environment* , 71, 56-66.
- Pu, R., Bell, S., Levy, K. H., & Meyer, C. (2010). Mapping detailed seagrass habitats using satellite imagery. *Proceedings of IGARSS 2010*. Honolulu, USA: IGARSS.
- Purkis, S. J., & Pasterkamp, R. (2004). Integrating in situ Reef-top Reflectance Spectra with Landsat Tm Imagery to Aid Shallow-Tropical Benthic Habitat Mapping. *Coral Reefs* , 23, 5–20.
- Rich, P. M. (1990). Characterizing plant canopies with hemispherical photographs. In G. N. S., & J. M. Norman (Eds.), *Instrumentation for studying vegetation canopies for remote sensing in optical and thermal infrared regions* (pp. 13-29). Remote Sensing Reviews.
- Richter, R., & Schlaepfer, D. (2002). Geo-atmospheric processing of airborne imaging spectrometry data, Part 2: atmospheric/topographic correction. *International Journal of Remote Sensin* , 23 (13), 2631–2649.
- Roelfsema, C. M., Joyce, K. E., & Phinn, S. R. (2004). Evaluation of benthic survey techniques for validating remotely sensed images of coral reefs. *Proceedings of the 12th International Coral Reefs Symposium*. Okinawa, Japan: International Coral Reef Society.



- Roelfsema, C. M., Phinn, S. R., Dennison, W. C., Dekker, A. G., & Brando, V. E. (2006). Monitoring toxic cyanobacteria *Lyngbya majuscula* (Gomont) in Moreton Bay, Australia by integrating satellite image data and field mapping. *Harmful Algae*, 5, 45-56.
- Roelfsema, C. M., Phinn, S. R., Udy, N., & Maxwell, P. (2009). An Integrated Field and Remote Sensing Approach for Mapping Seagrass Cover, Moreton Bay, Australia. *Spatial Science*, 54 (1), 45-62.
- Roelfsema, C., & Phinn, S. (2010). Integrating field data with high spatial resolution multispectral satellite imagery for calibration and validation of coral reef benthic community maps. *Journal of Applied Remote Sensing*, 4, 1-28.
- Rouse, J. W., Haas, R. H., Schell, J. A., & Deering, D. W. (1973). Monitoring vegetation systems in the Great Plains with ERTS. *Third ERTS Symposium. SP-351 I*, pp. 309- 317. NASA.
- Sagawa, T., Mikami, A., Komatsu, T., Kosaka, N., Kosaki, A., Miyazaki, S., et al. (2008). Technical Note. Mapping seagrass beds using IKONOS satellite image and side scan sonar measurements: a Japanese case study. *International Journal of Remote Sensing*, 29 (1), 281-291.
- Sagawa, T., Mikami, A., Komatsu, T., Kosaka, N., Kosaki, A., Miyazaki, S., et al. (2008). Technical Note. Mapping seagrass beds using IKONOS satellite image and side scan sonar measurements: a Japanese case study. *International Journal of Remote Sensing*, 29 (1), 281-291.
- Saputro, G. B., Hartini, S., Sukardjo, S., Susanto, A., & Poniman, A. (2009). *Peta Mangroves Indonesia*. Bogor: BAKOSURTANAL.
- Schlerf, M., Atzberger, C., & Hill, J. (2005). Remote Sensing of Forest Biophysical Variables Using Hymap Imaging Spectrometer Data. *Remote Sensing of Environment*, 95, 177-194.
- Seth, G. (2007). *How to calculate the amount of CO2 sequestered in a tree per year*. Retrieved 11 10, 2010, from <http://www.plant-trees.org>: <http://www.plant-trees.org/resources/Calculating%20CO2%20Sequestration%20by%20Trees.pdf>
- Shinozaki, K., Yoda, K., Hozumi, K., & Kira, T. (1964). A quantitative analysis of plant form—the pipe model theory. II. Further evidence of the theory and its application in forest ecology. *Japanese Journal of Ecology*, 14, 133-139.
- Short, F. T., & G, C. R. (2001). *Global Seagrass Research Methods*. (R. G. Coles, Ed.) Amsterdam: Elsevier Science B.V.
- Simard, M., Rivera-Monroy, V., Mancera-Pineda, J., Castañeda-Moya, E., & Twilley, R. (2008). A systematic method for 3D mapping of mangrove forests based on Shuttle Radar Topography Mission elevation data, ICESat/GLAS waveforms and field data: Application to Ciénaga Grande de Santa Marta, Colombia. *Remote Sensing of Environment, Earth Observations for Terrestrial Biodiversity and Ecosystems Special Issue*, 112 (5), 2131-2144.
- Simard, M., Zhang, K., Rivera-Monroy, V. H., Ross, M. S., Ruiz, P. L., Castañeda-Moya, E., et al. (2006). Mapping Height and Biomass of Mangrove Forests in Everglades National Park with SRTM Elevation Data. *Photogrammetric Engineering & Remote Sensing*, 72 (3), 299–311.
- Singer, R. (2013). *Geolocation Accuracy of WorldView Products*. Digital Globe.



- Spalding, J. (2001, October 7). *Absolute Global Cooling Through Seagrass*. Retrieved September 19, 2010, from The Ocean Foundation: <http://www.oceanfdn.org/index.php?ht=a/GetDocumentAction/i/186>
- Stenberg, P., Rautiainen, M., Manninen, T., Voipio, P., & Smolander, H. (2004). Reduced simple ratio better than NDVI for estimating LAI in Finnish pine and spruce stands. 2004. *Silva Fennica* , 38 (1), 3–14.
- Strong, A. E., Arzayus, F., Skirving, W., & Heron, S. F. (2006). Identifying Coral Bleaching Remotely via Coral Reef Watch - Improved Integration and Implications for Changing Climate. (J. T. Phinney, Ed.) *Coastal and Estuarine Studies* , 61, 163-180.
- Swain, P. H., & Davis, S. M. (1978). *Quantitative Approach of Remote Sensing*. Washington: McGraw-Hill.
- Thenkabail, P. S., Stucky, N., Griscom, B. W., Ashton, M. S., Enclona, E., Diels, J., et al. (2002). Biomass Estimations and Carbon Stock Calculations in the Oil Palm Plantations of African Derived Savannas using IKONOS Data. *FIEOS 2002 Conference Proceedings*. ISPRS.
- Thom, R. (2001). *Carbon Sinks in Nearshore Marine Vegetated Ecosystems* . Pacific Northwest Laboratory, Marine Science Laboratory.
- Updike, T., & Comp, C. (2010). *Radiometric Use of WorldView-2 Imagery*. Longmont, Colorado: DigitalGlobe®.
- Urbanski, J. A., Mazur, A., & Janas, U. (2009). Object-oriented classification of QuickBird data for mapping seagrass spatial structure. *International Journal of Oceanography and Hydrobiology* , 38 (1), 27-43.
- Vaiphasa, C., Skidmore, A. K., & de Boer, W. F. (2006). A post-classifier for mangrove mapping using ecological data,. *PandRS* , 61 (1), 1-10.
- Van Der Meer, F., Bakker, W., Scholte, K., Skidmore, A., De Jong, S., Clevers, J., et al. (2000). Vegetation Indices, Above Ground Biomass Estimates and The Red Edge From MERIS. *International Archives of Photogrammetry and Remote Sensing* , XXXIII, Part B7.
- Vichkovitten, T. (1998). Biomass, Growth and Productivity of Seagrass; *Enhalus acoroides* (Linn. f) in Khung Kraben Bay, Chanthaburi, Thailand. *Kasetsart J. (Nat. Sci)* , 32, 109-115.
- Viña, A., & Gitelson, A. A. (2011). Sensitivity to Foliar Anthocyanin Content of Vegetation Indices Using Green Reflectance. *IEEE Geoscience and Remote Sensing Letters* , 8 (3), 463-467.
- Viña, A., & Gitelson, A. A. (2011). Sensitivity to Foliar Anthocyanin Content of Vegetation Indices Using Green Reflectance. *IEEE Geoscience and Remote Sensing Letters* , 8 (3), 463-467.
- Viña, A., Gitelson, A. A., Rundquist, D. C., Keydan, G., Leavitt, B., & Schepers, J. (2004). Monitoring maize (*Zea mays* L.) phenology with remote sensing. *Agronomy Journal* , 96 (4), 1139–1147.
- Wamunyima, S. (2005). *Estimating Fresh Grass Biomass at Landscape Level Using Hyperspectral Remote Sensing*. Enschede, The Netherlands: International Institute for Geo-Information Science and Earth Observation (ITC).
- Wang, M., & Shi, W. (2008). Satellite-observed algae blooms in China's Lake Taihu. *EOS Trans Am Geophys Union* , 89, 201–202.



- Wicaksono, P. (2010). *Integrated Model of Water Column Correction Technique for Improving Satellite-Based Benthic Habitat Mapping*. Yogyakarta: Gadjah Mada University.
- Wicaksono, P. (2008). Perbandingan Kemampuan Citra Landsat 7 ETM+ dan ASTER dalam Pemetaan Kondisi Kesehatan Terumbu Karang. Studi Kasus Pulau Menjangan Besar dan Menjangan Kecil, Kepulauan Karimunjawa. *Proceedings of PIT MAPIN XVII*. Bandung, Indonesia: MAPIN.
- Wicaksono, P., & Hafizt, M. (2013). Mapping seagrass from space: Addressing the complexity of seagrass LAI mapping. *European Journal of Remote Sensing* , 46, 18-39.
- Wicaksono, P., Hartono, & Danoedoro, P. (2011). Mapping Seagrass as a Natural Carbon Sink: A Remote Sensing Approach. *Proceedings of International Seminar on Marine. "Sustainable Marine and Coastal Resource Management in Coral Triangle Initiative (CTI) Region"*. Denpasar, Bali, Indonesia.
- Wong, K. K., & Fung, T. (2010). *Leaf Area Index Modeling of Mangrove based on Hyperspectral Remote Sensing and in-situ Hemispherical Photography in Maipo Ramsar Site of Hong Kong*. Hong Kong: Department of Geography and Resource Management, The Chinese University of Hong Kong.
- Wu, Y., Chung, A., Tam, N. F., Pi, N., & Wong, M. H. (2008). Constructed mangrove wetlands as secondary treatment system for municipal wastewater. *Ecology Engineering* , 34, 137-146.
- Xiao, X., Boles, S., Frohling, S., Salas, W., Moore III, B., Li, C., et al. (2002). Observation of flooding and rice transplanting of paddy rice fields at the site to landscape scales in China using VEGETATION sensor data. *International Journal of Remote Sensing* , 23, 3009-3022.
- Yang, D., & Yang, C. (2009). Detection of seagrass distribution changes from 1991 to 2006 in Xincun Bay, Hainan, with Satellite Remote Sensing. *Sensor* , 9, 830-844.
- Yang, D., Yang, Y., Yang, C., Zhao, J., & Sun, Z. (2011). Detection of seagrass in optical shallow water with Quickbird in the Xincun Bay, Hainan province, China. *IET Image Processing* , 5 (5), 363-368.
- Yohe, G. W., Lasco, R. D., Ahmad, Q. K., Arnell, N. W., Cohen, S. J., Hope, C., et al. (2007). Perspectives on climate. In M. Parry, O. Canziani, J. Palutikof, P. van der Linden, & C. Hanson (Eds.), *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* (pp. 811-841). Cambridge, UK: Cambridge University Press.

Act No. 4/2011

Act No. 27/2007

PP No. 8/2013

Kepmen No. 200/2004

Kepmen No. 201/2004

SK Menhut No. 78/Kpts-II/1999

SK Dirjen PHKA NoSK.79/IV/Set-3/2005