

DAFTAR PUSTAKA

- AAP, A. A. (2004; 114). Clinical Practice Guideline: Management of Hyperbilirubinemia in the Newborn Infant 35 or More Weeks of Gestation. *Pediatrics*, 297-316.
- ABM, T. A. (2010). ABM Clinical Protocol #22: Guidelines for Management of Jaundice in the Breastfeeding Infant Equal to or Greater Than 35 Weeks' Gestation. *Breastfeeding Medicine*; Vol. 5; No. 2, 87-93.
- Acosta-Torres, S. M., Torres-Espina, M. T., Colina-Araujo, J. A., & Colina-Chourio, J. A. (2012). Usefulness of The Kramer's Index in the Diagnosis. *Invest Clin*; Vol. 53 (2), 148 - 156.
- Aladag, N., Filiz, T. M., Topsever, P., & Gourpelioglu, a. S. (2006). Parents Knowledge and Behaviour Concerning Sunning Their Babies: A cross-Sectional, Descriptive Study. *BMC Pediatrics*; Vol. 6; 27, 1-6.
- Almeida, M. F., & Draque, C. M. (2007). Neonatal Jaundice and Breastfeeding. *NeoReviews - American of Academy Pediatrics*; Vol. 9; No. 7, 282-289.
- Bertini, G., Dani, C., Tronchin, M., & Rubaltelli, a. F. (2001). Is Breastfeeding Really Favoring Early Neonatal Jaundice? *Pediatrics*; Vol. 107, e41-e45.
- Dennerly, P. A., Seidman, D. S., & Stevenson, D. K. (2001). Neonatal Hyperbilirubinemia. *N Engl J Med*; Vol. 344; No. 8, 583-590.
- Depkes. (2004). Tatalaksana Ikterus Neonatorum. *HTA Indonesia*, 1-23.
- Depkes. (2008). *Buku Bagan Manajemen Terpadu Balita Sakit (MTBS)*. Jakarta: Departemen Kesehatan Republik Indonesia.
- Gartner, L. M. (2001; Vol. 21). Breastfeeding and Jaundice. *Journal of Perinatology*, s25-s29.
- Gourley, G. R., Li, Z., Kreamer, B. L., & Kosorok, a. M. (2005). A Controlled, Randomized, Double-Blind Trial of Prophylaxis Against Jaundice Among Breastfed Newborns. *Pediatrics*; Vol. 166: 2, 384-393.
- Harrison, S., Buttner, P., & Nowak, a. M. (2005). Maternal Belief About the Reputed Therapeutic Uses of Sun Exposure in Infancy and the Postpartum Period. *Australian Midwifery*; Vol. 18: 2, 22-28.
- HTA, H. T. (2012). *Management of Neonatal Hyperbilirubinemia*. Malaysia: Health Technology Assesment Unit Medical Development Division Ministry of Malaysia.
- JNPK-KR, J. N.-K. (2008). *Asuhan Persalinan Normal & Inisiasi Menyusu Dini*. Jakarta: JNPK - KR Departemen Kesehatan Republik Indonesia.
- Lauer, B. J., & Spector, N. D. (2011). Hyperbilirubinemia in the Newborn. *American Academy of Pediatrics*; Vol. 32; No. 8, 341-349.
- Lin, Y.-Y., Tsao, P.-N., Hsieh, W.-S., Chen, C.-Y., & Chou, a. H.-C. (2008). The Impact of Breast-Feeding on Early Neonatal Jaundice. *Clinical Neonatology*, 31-35.
- Maisels, M. J. (2010). Screening an Early Posnatal Management Strategies to Prevent Hazardous Hyperbilirubinemia in Newborn of 35 or More Weeks of Gestation. *Seminars in Fetal & Neonatal Medicine* 15 (pp. 129 - 135). Oakland, USA: Elsevier.

- Maisels, M. J., Bhutani, V. K., Bogen, D., Newman, T. B., Stark, A. R., & Watchko, a. J. (2009). Hyperbilirubinemia in The Newborn Infant \geq 35 Weeks Gestation: An Update With Clarifications. *Pediatrics*, 1192-1199.
- Nursanti, Ida. (2011). *Pengaruh Kecukupan ASI terhadap Risiko Terjadinya Ikterus Neonatorum di Yogyakarta*. Yogyakarta: Universitas Gadjah Mada.
- Pediatrics, A. A. (2004; Vol. 114: 1). Management of Hyperbilirubinnemia in the Newborn Infant 35 or More Weeks of Gestation. *Pediatrics*, 297-316.
- Smithermen, H., Stark, A. R., & and Bhutani, V. K. (2006; Vol. 11: 3). Early Recognition of Neonatal Hyperbilirubinemia and Its Emergent Management. *Semin Fetal Neonatal Med*, 214-224.
- Suparyanto. (2011, Juli 07). Inisiasi Menyusu Dini (IMD). <http://dr-suparyanto.blogspot.com/2011/07/inisiasi-menyusu-dini-imd.html>.
- Suradi, R., & Letupeirissa, D. (2013, August 23). Air Susu Ibu dan Ikterus. *Buku Bedah ASI IDAI*.
- Suradi, R., Situmeang, E. H., & Tambunan, d. T. (2001; Vol. 41). The Association of Neonatal Jaundice and Breast - Feeding. *Paediatrica Indonesiana*, 69-75.
- Suresh, G. K., & Clark, R. E. (2004). Cost-effectiveness of strategies that are intended to prevent kernicterus in newborn infants. *Pediatrics*; 114, 917-24.
- Surjono, A. (1995; 27). Hiperbilirubinemia pada Neonatus: Pendekatan Kadar Bilirubin Bebas. *Berkala Ilmu Kedokteran*, 43-46.
- Tikmani, S. S., Warraich, H. J., Abbasi, F., Rizvi, A., Darmstadt, G. L., & Zaidi, a. A. (2010). Incidence of Neonatal Hyperbilirubinemia: A Population-Based. *Tropical Medicine and International Health*; Vol. 15 No. 5, 502–507 .
- UNICEF. (2007). *Breast Crawl - Initiation of Breastfeeding by Breast Crawl (Insert with Breast Crawl Video) 1st Edition*. Mumbai; India: UNICEF Maharashtra.
- Utami, R. (2008). *Inisiasi Menyusu Dini Plus ASI Eksklusif*. Jakara: Pustaka Bunda.
- WHO. (2003). *Managing Newborn Problems: A Guide for Doctors, Nurses, and Midwives*. Geneva: Departement of Reproductive Health and Research, WHO.
- Yang, W.-C., Zhao, L.-L., Li, Y.-C., Chen, C.-H., Chang, Y.-J., & Fu, Y.-C. (2013; Vol. 13). Bodyweight Loss in Predicting Neonatal Hyperbilirubinemia 72 Hours after Birth in Term Newborn Infants. *BMC Pediatrics*, 145-151.