

## LIST OF REFERENCES

---

- Abdullah, A., Nassr, S., & Ghaleeb, A. (2013). Landsat ETM-7 for lineament mapping using automatic extraction technique in the SW part of Taiz area, Yemen, *13*(3).
- Ali, A., & Pour, A. (2014). Lithological mapping and hydrothermal alteration using Landsat 8 data: a case study in ariab mining district, red sea hills, Sudan. *International Journal of Basic and Applied Sciences*, *3*(3), 199–208. <http://doi.org/10.14419/ijbas.v3i3.2821>
- Alvarez-Berríos, N. L., & Mitchell Aide, T. (2015). Global demand for gold is another threat for tropical forests. *Environmental Research Letters*, *10*(1), 14006. <http://doi.org/10.1088/1748-9326/10/1/014006>
- Aspinall, C. (2001). Small-scale mining in Indonesia, (79).
- Australian Government Geoscience Australia. (n.d.). 2.4 Iron oxide copper-gold mineral systemtAustralia. Retrieved from <http://www.ga.gov.au/data-pubs/data-and-publications-search/publications/critical-commodities-for-a-high-tech-world/iron-oxide-copper-gold>
- Berk, a., Bernstein, L. S., Anderson, G. P., Acharya, P. K., Robertson, D. C., Chetwynd, J. H., & Adler-Golden, S. M. (1998). MODTRAN cloud and multiple scattering upgrades with application to AVIRIS. *Remote Sensing of Environment*, *65*(3), 367–375. [http://doi.org/10.1016/S0034-4257\(98\)00045-5](http://doi.org/10.1016/S0034-4257(98)00045-5)
- Billings, W. D. (1950). Vegetation and plant growth as affected by chemically altered rocks in the Western Great Basin. *Ecology*. Retrieved from [http://www.jstor.org/stable/1931361?seq=7#page\\_scan\\_tab\\_contents](http://www.jstor.org/stable/1931361?seq=7#page_scan_tab_contents)
- Billings, W. D. (1952). The environmental complex in relation to plant growth and distribution. *The Quarterly Review of Biology*. <http://doi.org/10.1086/399022>
- Blaschke, T., Lang, S., Lorup, E., Strobl, J., & Zeil, P. (2002). Object-oriented image processing in an integrated GIS / remote sensing environment and perspectives for environmental applications, 555–570.
- Bodruddoza Mia, M., & Fujimitsu, Y. (2012). Mapping hydrothermal altered mineral deposits using Landsat 7 ETM+ image in and around Kuju volcano, Kyushu, Japan. *Journal of Earth System Science*, *121*(4), 1049–1057. <http://doi.org/10.1007/s12040-012-0211-9>

- Brown, S. (1997). Estimating Biomass and Biomass Change of Tropical Forests: a Primer. (FAO Forestry Paper). Retrieved April 10, 2015, from <http://www.fao.org/docrep/w4095e/w4095e00.HTM>
- Brown, S., Iverson, L. R., Prasad, A., & Liu, D. (1993). Geographical distributions of carbon in biomass and soils of tropical Asian forests. *Geocarto International*, 8(4), 45–59. <http://doi.org/10.1080/10106049309354429>
- Caran, S. C., Jr, C. M. W., & Thompson, E. J. (1982). Lineament analysis and inference of geologic structure : examples from the Balcones Ouachita trend of Texas, XXXI.
- Center of Soil Research. (1982). Indonesia soil classification (in Indonesia).
- Chanan, M. (2012). Estimation of above ground carbon on Teak plantation (*Tectona grandis* Linn.F)( in RPH Sengguruh BKPH Sengguruh KPH Malang Perhutani East Java) (in Indonesia), 7(C), 61–73.
- Charou, E., Stefouli, M., Dimitrakopoulos, D., Vasilou, E., & Mavrantza, O. D. (2010). Using remote sensing to assess impact of mining activities on land and water resources. *Mine Water and the Environment*, 29(1), 45–52. <http://doi.org/10.1007/s10230-010-0098-0>
- Chevrel, S., Bourguignon, A., Cottard, F., & Itard, Y. (2005). Exploitation of ASTER imagery in mining-related environmental management.
- Chupezi, T., Ingram, V., & Schure, J. (2009). Impacts of artisanal gold and diamond mining on livelihoods and the environment in the Sangha Tri-National Park landscape. Retrieved from [http://www.cifor.org/publications/pdf\\_files/books/bchupezi0901.pdf](http://www.cifor.org/publications/pdf_files/books/bchupezi0901.pdf)
- Colaborative Partnership on Forest (CPF). (2014). Assessment and monitoring of forest degradation. Retrieved August 11, 2014, from <http://www.cpfweb.org/73032/en/>
- Communication and Information Agency. (2009). Environmental agency monitor the Banyuwangi mining exploration. Retrieved April 20, 2015, from <http://kominfo.jatimprov.go.id/watch/18195>
- Condit, R. (2008). Methods for estimating aboveground biomass of forest and replacement vegetation in the tropics. *Center for Tropical Forest Science Research Manual*, (August), 73 pages.

- Decree of Minister of Environmental. (2004). Decree of Minister of Environmental of 51 of sea water standard (in Indonesia).
- Definiens. (2011). *eCognition Developer 8.7 reference book*. Munchen Germany.
- Energy and Mineral Resources Agency. (2013). Strategic minerals in Districts (in Indonesia). Retrieved from <http://webmap.psdg.bgl.esdm.go.id/geosain/neraca-mineral-strategis.php?mode=administrasi>
- Energy and Mineral Resources Departement. (2009). 2009 Mining colloquium (in Indonesia) (pp. 978–979).
- Erk, A. Van. (2011). Application of remote sensing for ecosystem services monitoring in tropical forest conservation, (September).
- FAO. (2003). Deforestation and forest degradation factors. Retrieved August 10, 2014, from <http://www.fao.org/docrep/article/wfc/xii/ms12a-e.htm>
- FAO. (2011). *Assessing forest degradation towards the development of globally applicable guidelines*.
- Gabr, S., Ghulam, A., & Kusky, T. (2010). Detecting areas of high-potential gold mineralization using ASTER data. *Ore Geology Reviews*, 38(1-2), 59–69. <http://doi.org/10.1016/j.oregeorev.2010.05.007>
- Gao, Y., Mas, J. F., Niemeyer, I., Marpu, P. R., & Palacio, J. L. (2003). Object-based image analysis for mapping land-cover in a forest area, (8701).
- Ge, L., Chang, H.-C., & Rizos, C. (2007). Mine subsidence monitoring using multi-source satellite SAR images. *Photogrammetric Engineering & Remote Sensing*, 73(3), 259–266. <http://doi.org/10.14358/PERS.73.3.259>
- Geology voice: Potency of gold mineral in Malang Regency (in Indonesia). (2014). Retrieved December 1, 2014, from <http://suarageologi.blogspot.com/2014/05/potensi-mineral-emas-di-kabupaten-malang.html>
- Gibbs, H. K., Brown, S., Niles, J. O., & Foley, J. a. (2007). Monitoring and estimating tropical forest carbon stocks: making REDD a reality. *Environmental Research Letters*, 2(4), 045023. <http://doi.org/10.1088/1748-9326/2/4/045023>
- Hashim, M., Ahmad, S., Johari, M. A., & Pour, A. B. (2013). Automatic lineament extraction in a heavily vegetated region using Landsat Enhanced

- Thematic Mapper ( ETM + ) imagery. *Advances in Space Research*, 51(5), 874–890. <http://doi.org/10.1016/j.asr.2012.10.004>
- Hedenquist, J. W., & Lowenstern, J. B. (1994). The role of magmas in the formation of hydrothermal ore deposits.
- Hellman & Schofield Pty Ltd. (2011). *Tujuh Bukit Project Report on Mineral Resources , Technical Report for Interpid Mines Limited.*
- Hidayat, M. Y. (2006). *Evaluation of land suitability of Paraserianthes falcataria L Nielsen in some slope units (case study in Cipatat Bandung West Java ) (in Indonesia).*
- IDL. (2009). ENVI atmospheric correction module: QUAC and FLAASH user's guide. *Module Version.* Retrieved from <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:ENVI+Atmospheric+Correction+Module:+QUAC+and+FLAASH+user's+guide#0>
- Igneous Rocks. (n.d.). Retrieved April 19, 2015, from <http://www.galleries.com/rocks/igneous.htm>
- Indonesia Investments. (2014). Gold mining in Indonesia - Indonesian gold production consumption. Retrieved August 15, 2014, from <http://www.indonesia-investments.com/doing-business/commodities/gold/item167>
- Indonesia Nasional Standard of 7724. (2011). Ground based forest carbon accounting (in Indonesia).
- IPCC. (2006). *Guidelines for national greenhouse gas inventories Volume 4 Agriculture, Forestry and Other Land Use.* Retrieved from <http://www.ipcc-nggip.iges.or.jp/public/2006gl/vol4.html>
- ITT Visual Information Solutions. (n.d.). ENVI Tutorial - Advanced Hyperspectral Analysis.
- Jaya, I. N. . (2006). *Practical guideline of remote sensing.*
- Jordan, G., & Schott, B. (2005). Application of wavelet analysis to the study of spatial pattern of morphotectonic lineaments in digital terrain models. A case study. *Remote Sensing of Environment*, 94, 31–38. <http://doi.org/10.1016/j.rse.2004.08.013>
- Karna, Y. K. (2012). *Mapping above ground carbon using worldview satellite image and lidar data in relationship with tree diversity of forests.* University

of Twente Faculty of Geo-Information and Earth Observation (ITC).  
Retrieved from [http://www.itc.nl/library/papers\\_2012/msc/nrm/karna.pdf](http://www.itc.nl/library/papers_2012/msc/nrm/karna.pdf)

- Kocal, A., Duzgun, H. S., Karpuz, C., Technologies, G. I., Mapping, D., & Imagery, H. (n.d.). Discontinuity mapping with automatic lineament extraction from high resolution satellite imagery.
- Kujjo, C. P. (2010). Application of remote sensing for gold exploration in the Nuba Mountains , Sudan, (May).
- Lee, J. Y., & Warner, T. A. (2014). Image classification with a region based approach in high spatial resolution imagery.
- Li, W., Guo, Q., Jakubowski, M. K., & Kelly, M. (2012). A new method for segmenting individual trees from the Lidar point cloud. *Photogrammetric Engineering and Remote Sensing*, 78, 75–84.
- Liu, L., Zhou, J., Jiang, D., Zhuang, D., Mansaray, L. R., & Zhang, B. (2013). Targeting mineral resources with remote sensing and field data in the Xiemisitai area, West Junggar, Xinjiang, China. *Remote Sensing*, 5(7), 3156–3171. <http://doi.org/10.3390/rs5073156>
- Lowe, S. (2014). Can mining be part of a low-carbon, low-deforestation development strategy? the case of Guyana. *Environment Magazine*. Retrieved from <http://www.environmentmagazine.org/Archives/Back Issues/2014/May-June 2014/mining-full.html>
- Minister of Energy and Mineral Resources. (2014). Decree of Minister of Energy and Mineral Resources No. 1204 K/30/MEM/2014 of mining area in Java and Bali (in Indonesia).
- Ministry of Forestry of Republic of Indonesia. Technical guideline of carbon estimation of national forest inventory (in Indonesia) (2013).
- Musa, H. D., & Jiya, S. N. (2011). An assessment of mining activities impact on vegetation in bukuru Jos Plateau State Nigeria using Normalized Differential Vegetation Index ( NDVI ). *Journal of Sustainable Development*, 4(6), 150–159. <http://doi.org/10.5539/jsd.v4n6p150>
- Najafi, C. (2011). Gold formation, extraction, and refining in the Sierra Nevada Mountain Region, 1–15.
- Natural Gold Indicators (Part 2) | RareGoldNuggets.com. (2014). Retrieved April 14, 2015, from <http://raregoldnuggets.com/blog/?p=750>

- Nguyen, T. N. (2010). *Estimation and mapping of above ground biomass for the assessment and mapping of carbon stocks in tropical forest using SAR data : a case study in Afram headwaters forest, Ghana. University of Twente Faculty of Geo-Information and Earth Observation (ITC)*. Retrieved from <http://adlib.itc.nl/detail.aspx?parentpreref=>
- Patterson, M. E. (2003). Hydrothermal alterations and the formation of metal ores in the Sierra Nevada's. Retrieved April 19, 2015, from <http://www.indiana.edu/~sierra/papers/2003/Patterson.html>
- Perhutani. (2014). *Brief review of illegal gold mining of KPH South Banyuwangi.pdf*.
- Perkins, T., Adler-golden, S., Matthew, M., Berk, A., Anderson, G., Gardner, J., & Felde, G. (2005). Retrieval of atmospheric properties from hyper- and multi-spectral imagery with the FLAASH atmospheric correction algorithm. *Proc. of SPIE Vol 5979 5979E-1*, 5979, 1–11. <http://doi.org/10.1117/12.626526>
- Placer deposit. (2010). Retrieved from <http://www.indiana.edu/~sierra/papers/2010/cross-najafi.html>
- Pour, a B., & Hashim, M. (2014). Alteration mineral mapping using ETM+ and hyperion remote sensing data at Bau Gold Field, Sarawak, Malaysia. *IOP Conference Series: Earth and Environmental Science*, 18, 012149. <http://doi.org/10.1088/1755-1315/18/1/012149>
- Pour, A. B., & Hashim, M. (2014). Hydrothermal alteration mapping using Landsat-8 data, Sar Cheshmeh copper mining district, SE Iran. *Journal of Taibah University for Science*. <http://doi.org/10.1016/j.seizure.2014.02.015>
- Pour, A. B., Hashim, M., & Marghany, M. (2014). Exploration of gold mineralization in a tropical region using Earth Observing-1 (EO1) and JERS-1 SAR data: A case study from Bau gold field, Sarawak, Malaysia. *Arabian Journal of Geosciences*, 7(6), 2393–2406. <http://doi.org/10.1007/s12517-013-0969-3>
- Raina, A. K., & Gupta, M. K. (2009). Soil and vegetation studies in relation to parent material of Garhwal Himalayas, Uttarakhand India.
- Raj, D. P. P. (2005). *Ground improvement techniques (PB)*. Firewall Media. Retrieved from <https://books.google.com/books?id=cDGIhh7ttMcC&pgis=1>
- Ramadan, T. M., & Abdel Fattah, M. F. (2010). Characterization of gold mineralization in Garin Hawal area, Kebbi State, NW Nigeria, using remote



- sensing. *Egyptian Journal of Remote Sensing and Space Science*, 13(2), 153–163. <http://doi.org/10.1016/j.ejrs.2009.08.001>
- Rigina, O. (2002). Environmental impact assessment of the mining and concentration activities in the Kola Peninsula, Russia by multirate remote sensing. *Environmental Monitoring and Assessment*, 75(1), 11–31. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15900663>
- Sabins, F. F. (1999). Remote sensing for mineral exploration. *Ore Geology Reviews*, 14(3-4), 157–183. [http://doi.org/10.1016/S0169-1368\(99\)00007-4](http://doi.org/10.1016/S0169-1368(99)00007-4)
- Sartohadi, J., Sianturi, R. S., Rahmadana, A. D. W., Maritimo, F., Wacano, D., Munawaroh, ... Pratiwi, E. S. (2014). *Landscape of Ijen Volcano and its surrounding*. Yogyakarta.
- Schnitzler, M. (2010). *Calculation of Carbon Capture in Tropical Reforestation Projects La Gamba , Costa Rica ( Rainforest of the Austrians )*.
- Shanks III, W. C. P. (2012). *Hydrothermal alteration. Volcanic Massive Sulphide Occurrence Model*.
- Shrestha, S. K. (2011). *Carbon stock estimation using very high resolution satellite imagery and individual crown segmentation ( a case study of broadleaved and needle leaved forest of Dolakha , Nepal )*.
- Singh, H., & Huat, B. B. . (2012). *Handbook of tropical residual soils engineering*. CRC Press. Retrieved from <https://books.google.com/books?id=UmWPqahaECIC&pgis=1>
- Sreejesh, K. K., Thomas, T. P., Rugmini, P., Prasanth, K. M., & Kripa, P. K. (2013). Carbon sequestration potential of Teak ( *Tectona grandis* ) plantations in Kerala, 2, 167–170.
- Sukamar, M., Venkatesan, N., & Babu, C. N. K. (2014). A Review of Various Lineament Detection Techniques for high resolution Satellite Images. *International Journal of Advanced Research in Computer Science and Software Engineering*, 4(3), 72–78.
- Susintowati. (2012). *Mercury bioacumulation and communitystructure of gastropoda in ex-area of traditional gold tailing in Lampon river, Banyuwangi, East Java (in Indonesia)*. Universitas Gadjah Mada. Retrieved from [http://etd.ugm.ac.id/index.php?mod=penelitian\\_detail&sub=PenelitianDetail&act=view&typ=html&buku\\_id=57729&obyek\\_id=4](http://etd.ugm.ac.id/index.php?mod=penelitian_detail&sub=PenelitianDetail&act=view&typ=html&buku_id=57729&obyek_id=4)

- Tom-Dery, D., Dagben, Z. J., & Cobbina, S. J. (2012). Effect of illegal small-scale mining operations on vegetation cover of arid, 4(6), 674–679.
- Wardani, S. F. Y. (2013). *Estimation of carbon stock changes in above ground woody biomass due to volcano pyroclastic flow and pyroclastic surge : the case of Merapi Volcano National Park. University of Twente Faculty of Geo-Information and Earth Observation (ITC)*. Retrieved from [http://www.itc.nl/library/papers\\_2013/msc/aes/wardani.pdf](http://www.itc.nl/library/papers_2013/msc/aes/wardani.pdf)
- Woldai, T. (2001). Application of remotely sensed data and GIS in assessing the impact of mining activities on the environment, 75–84.
- Woldai, T., & Fabbri, A. G. (1998). The impact of mining on the environment.- a case study from the Tharsis-Lagunazo mining area, Province Huelva, SW Spain.
- Wysocki, D. A., Schoeneberger, P. J., Hirmas, D. R., & LaGarry, H. E. (2011). Geomorphology of soil landscapes. In *Handbook of Soil Sciences* (pp. 1–26). <http://doi.org/doi:10.1201/b11267-35>