

CHAPTER VI

REFERENCES

- Aditya, R., 2010, *Association Between The History Of Prenatal And Perinatal Events With Severe Profound Sensorineural Hearing loss In Children In RSUP Dr. Kariadi Semarang*, pp.1-22
- Blum NJ, Baron MA. Speech and Language Disorders.,1997, In: SchwartzMV, ed. *Pediatric Primary Care: a problem oriented approach*. St. Louis: Mosby :845-900
- Downey, D., Mraz, R., Knott, J., Knutson,C.,Holte, L., Van Dyke, D. 2002. Diagnosis and evaluation of children who are not talking.*Inf Young Children*; 15(2):38-48
- First, L.R., Palfrey, J.S.1994. The infant or young child with development delay.*N Engl j Med*; 330(7): 478-83
- Hediger, M.L.,Overpeck, M.D., Ruan, W.J., Troendle, JF. 2002. Birthweight and Gestational Age Effects on Motor and Social Development. *Paediatrics and Perinatal Epidemiology*;16
- Khalil, K.A., El-Amrawy, S.M., Ibrahim, A.G., El-Jeimy, N.A., Greiw, A.E. 1995. Pattern of growth and development of premature children at the age of two and three years in Alexandria, Egypt (part 1). 1(2): 176-85. Available at: <http://www.emro.who.int/publication/emjh>.
- Leung, A.,Kao, C.P.1999.Evaluation and management of the child with speech delay. *Am Fam Physician*; 59:3121-3128,3135
- Roizen NJ. 2003. Nongenetic causes of hearing loss. *Mental retardation and developmental disabilities research reviews* 9: 120-7
- Beata AS, Ioana AN., 2008, Etiopahtogeny of congenital and early-onset hearing loss; detection and early

intervention methos in infants and children. J
Clin med 3(1):45-51

Nlm.nih.gov, (2015). *Neonatal sepsis: MedlinePlus Medical Encyclopedia*. [online] Available at:
<http://www.nlm.nih.gov/medlineplus/ency/article/007303.htm> [Accessed 10 Jan. 2015].

Board, A. (2013). Neonatal sepsis. *A.D.A.M.*. [online] Available at:
<http://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0004557/> [Accessed 10 Jan. 2015].

Mmspf.msdonline.com.br, (2015). *Sensorineural Hearing Loss*. [online] Available at:
<http://mmspf.msdonline.com.br/ebooks/HeadNeckSurgeriyOtolaryngology/sid682242.html> [Accessed 10 Jan. 2015].

Perera, R., Glasziou, P., Heneghan, C., McLellan, J., Williamson, I. (1996). Autoinflation for hearing loss associated with otitis media with effusion. *Cochrane Database of Systematic Reviews*.

Apgar BJ, Greenberg G, Yen G., 2005, Prevention of group B streptococcal disease in the newborn, *Am Fam Physician*; 71(5):903-910

Nlm.nih.gov, (2015). *Sensorineural deafness: MedlinePlus Medical Encyclopedia*. [online] Available at:
<http://www.nlm.nih.gov/medlineplus/ency/article/003291.htm> [Accessed 14 Jan. 2015].

America, H. (2015). Types, Causes and Treatment | Hearing Loss Association of America. [online] [Hearingloss.org](http://www.hearingloss.org). Available at:
<http://www.hearingloss.org/content/types-causes-and-treatment> [Accessed 14 Jan. 2015].

Asha.org, (2015). *Sensorineural Hearing Loss*. [online] Available at:
<http://www.asha.org/public/hearing/Sensorineural-Hearing-Loss/> [Accessed 14 Jan. 2015].

- Nichd.nih.gov, (2015). Antibiotic regimens for suspected early neonatal sepsis. [online] Available at: https://www.nichd.nih.gov/cochrane_data/mtitimilae_01/mtitimilae_01.html [Accessed 14 Jan. 2015].
- WHO (2008). The global burden of disease: 2004 update (online.August) e.d. Geneva, Switzerland: *World Health Organization*. P.35
- Yoshinaga-Itano C., 2004. Levels of evidence: universal newborn hearing screening (UNHS) and early detection and intervention system (EHDI). *J Commun and Disord* 37:451-465
- Dunmed A.D, et. al., 2006, Profound bilateral sensorineural hearing loss in Nigerian children, *J Deaf Stud Deaf Educ*, 12 ed., pp. 112-18.
- Schachtele, S., Mutnal, M., Schleiss, M. Lokensgard, J. (2011). Cytomegalovirus-induced sensorineural hearing loss with persistent cochlear inflammation in neonatal mice. *Journal of NeuroVirology*, 17(3), pp.201-211.
- Nishiyama, N., Hardie, N. and Shepherd, R. (2000). Neonatal sensorineural hearing loss affects neurone size in cat auditory midbrain. *Hearing Research*, 140(1-2), pp.18-22.
- Fujioka, M., Okano, H. and Ogawa, K. (2014). Inflammatory and immune responses in the cochlea: potential therapeutic targets for sensorineural hearing loss. *Front. Pharmacol.*, 5.