

DAFTAR PUSTAKA

- Albino, S., Tabb, K. M., Requena, D., Egoavil, M., Pineros-Leano, M. F., Zunt, J. R., & García, P. J. (2014). Perceptions and acceptability of short message services technology to improve treatment adherence amongst tuberculosis patients in Peru: a Focus Group Study. *PloS one*, 9(5), e95770. doi:10.1371/journal.pone.0095770
- Baranoski, Elizabeth Meuser, Helene Hardya, Elizabeth F. Closson, Matthew J. Mimiagacde, Steven A. Safrende, Pushwaz Virkf, Rowena Lukf, P. R. S. & V. S. K. (2013). Patient and provider perspectives on cellular phone-based technology to improve HIV treatment adherence. *AIDS Care: Psychological and Socio-medical Aspects of AIDS/HIV*, Volume 26, (Issue 1), pages 26–32. doi:DOI:10.1080/09540121.2013.802282
- Barclay, E. (2009). Text messages could hasten tuberculosis drug compliance. *The Lancet*, 373(9657), 15–16. doi:10.1016/S0140-6736(08)61938-8
- Beratarrechea, A., Lee, A. G., Willner, J. M., Jahangir, E., Ciapponi, A., & Rubinstein, A. (2014). The impact of mobile health interventions on chronic disease outcomes in developing countries: a systematic review. *Telemedicine journal and e-health : the official journal of the American Telemedicine Association*, 20(1), 75–82. doi:10.1089/tmj.2012.0328
- Bowers, D; House, A; Owens, D. (2011). *Getting Started in Health Research*. Wiley.
- Car, J., Gurol-Urganci, T, D. J., Jamsek, V., & Atun, R. (2012). Mobile phone messaging reminders for attendance at healthcare appointments (Review). *The Cochrane Collaboration*, (7), 41. Retrieved from h
- Chen, Z., Fang, L., Chen, L., & Dai, H. (2008). Comparison of an SMS text messaging and phone reminder to improve attendance at a health promotion center : A randomized controlled trial, 9(1), 34–38. doi:10.1631/jzus.B071464
- Comolet, T., Rakotomalala, R.H. (1998). Factors determining compliance with tuberculosis treatment in an urban environment, Tamatave, Madagascar. *Int J Tuberc Lung Dis.*, 2(11)(891), 7.
- DeSouza, S. I., Rashmi, M. R., Vasanthi, A. P., Joseph, S. M., & Rodrigues, R. (2014). Mobile phones: the next step towards healthcare delivery in rural India? *PloS one*, 9(8), e104895. doi:10.1371/journal.pone.0104895
- Dowshen, N., Kuhns, L. M., Johnson, A., Holoyda, B. J., & Robert Garofalo. (2012). Improving Adherence to Antiretroviral Therapy for Youth Living with HIV AIDS A Pilot Study Using Personalized, Interactive, Daily Text Message

Reminders. *J Med Internet Res*. 2012 Mar-Apr; 14(2): e51. Retrieved from URL: <http://www.jmir.org/2012/2/e51/>

Farah, M.G., Tverdal, S.T. (2005). Treatment outcome of new culture positive pulmonary tuberculosis in Norway. *BMC Public Health*, 5(1), 14.

Fatehi, F., Malekzadeh, G., Akhavimirab, A., Rashidi, M., & Afkhami-ardekani, M. (2010). The Effect of Short Message Service on Knowledge of Patients with Diabetes in Yazd , Iran, (3), 27–31.

Fjeldsoe, B. S., Marshall, A. L., & Miller, Y. D. (2009). Behavior change interventions delivered by mobile telephone short-message service. *American journal of preventive medicine*, 36(2), 165–73.
doi:10.1016/j.amepre.2008.09.040

Free, C., Phillips, G., Galli, L., Watson, L., Felix, L., Edwards, P., Patel, V., et al. (2013). The effectiveness of mobile-health technology-based health behaviour change or disease management interventions for health care consumers: a systematic review. *PLoS medicine*, 10(1), e1001362.
doi:10.1371/journal.pmed.1001362

Gold, J., Lim, M. S. C., Hellard, M. E., Hocking, J. S., & Keogh, L. (2010). What 's in a message ? Delivering sexual health promotion to young people in Australia via text messaging. *BMC Public Health*, 10(1), 792. doi:10.1186/1471-2458-10-792

Green, L.W., Kreuter, M. (2005). The Precede-Proceed Model of Health Program Planning & Evaluation. Retrieved April 28, 2015, from <http://www.lgreen.net/precede.htm>

Gubernur DIY (2013). UMK Tahun 2014 di Daerah Istimewa Yogyakarta. Retrieved from <http://www.nakertrans.jogjaprov.go.id/contentdetil.php?kat=brta&id=MTI5&fle=Y29udGVudC5waHA=&lback=a2F0PWJydGEmbGJhY2s9JnBhZ2U9Mw==>

Heather Cole-Lewis, T. K. (2011). *NIH Epidemiol Rev*, 32(1), 56–69.
doi:10.1093/epirev/mxq004.Text

Herlina, S. (2013). *Short message service reminder sbg media promosi bumil di Kab. Banjar*. Gadjah Mada University, Yogyakarta.

Hoefman, A. B. (2010). Using SMS for HIV/AIDS education and to expand the use of HIV testing and counselling services at the AIDS Information Centre (AIC) Uganda, (November), 40–48. Retrieved from <http://kau.diva-portal.org/smash/get/diva2:357565/FULLTEXT01#page=43>

- Iribarren, S., Chirico, C., Echevarria, M., & Cardinali, D. (2012). mHIMMS 2012 Abstract TextTB: A parallel design randomized control pilot study to evaluate acceptance and feasibility of a patient-driven mobile phone based intervention to support adherence to TB treatment. *JMTM*, 1(4), 23–24. doi:10.7309/jmtm.46
- Jaiswal, A., Singh, V., Ogden, J.A. (2003). Adherence to tuberculosis treatment: lessons from the urban setting of Delhi, India. *Trop Med Int Health*, 8(7): 625-
- M.'Imunya, T, K. (2012). Patient education and counselling for promoting adherence to treatment for tuberculosis (Review). *The Cochrane Collaboration*, (5), 31. Retrieved from <http://www.thecochranelibrary.com>
- Kementerian Kesehatan RI (2011). *Pedoman Nasional Penanggulangan Tuberkulosis (TB)*. (T. Y. Aditama, Ed.). Jakarta, Indonesia. Retrieved from <http://www.hukor.depkes.go.id>
- Khan, A. (2012). mHealth to improve TB care. Retrieved from <http://irdresearch.org/wp-content/uploads/2012/05/mHealth-to-Improve-TB-Care.pdf>
- Kusfriyadi, M. (2010). *Pengaruh Pendidikan Gizi Ibu Hamil dan Pesan Gizi Melalui Short Message Service (SMS) Terhadap Pengetahuan, Perilaku, Kepatuhan Minum Tablet Besi dan Kadar Hemoglobin Ibu Hamil di Kota Palangkaraya*. Universitas Gadjah Mada. Retrieved from <http://infolib.med.ugm>
- Lester, R. T., Ritvo, P., Mills, E. J., Kariri, A., Karanja, S., Chung, M. H., Jack, W (2010). Effects of a mobile phone short message service on antiretroviral treatment adherence in Kenya (WelTel Kenya1): a randomised trial. *Lancet*, 376, 1838–45. doi:10.1016/S0140-6736(10)61997-6
- Lewis, C. P., & Newell, J. N. (2009). Improving tuberculosis care in low income countries - a qualitative study of patients' understanding of "patient support" in Nepal. *BMC public health*, 9, 190. doi:10.1186/1471-2458-9-190
- Liliweri, A. (2007). *Dasar-Dasar Komunikasi Kesehatan* (Pertama., p. 363). Yogyakarta: Pustaka Pelajar.
- Liu, Q., Abba, K., Mm, A., Sinclair, D. & Mad, L. (2014). Reminder systems to improve patient adherence to tuberculosis clinic appointments for diagnosis and treatment (Review), (11).
- Maulana, H. (2007). *Promosi Kesehatan*. (E. K. Yudha, Ed.) (Kesatu., p. 271). Jakarta.
- Mbuagbaw, L., Bonono-Momnougui, R. C., & Thabane, L. (2012). Considerations in using text messages to improve adherence to highly active antiretroviral therapy:

a qualitative study among clients in Yaoundé, Cameroon. *HIV/AIDS (Auckland, N.Z.)*, 4, 45–50. doi:10.2147/HIV.S29954

- Mbuagbaw, L., Mursleen, S., Lytvyn, L., Smieja, M., Dolovich, L., & Thabane, L. (2015). Mobile phone text messaging interventions for HIV and other chronic diseases: an overview of systematic reviews and framework for evidence transfer. *BMC health services research*, 15(1), 33. doi:10.1186/s12913-014-0654-6
- McKay, M., Davis, M., & Fanning, P. (1995). *Messages, The Communication Skills Book* (Second., p. 310). Oakland, CA 94609: New Harbinger Publications, Inc.
- Miller & Himelhoch, S. (2013). Acceptability of Mobile Phone Technology for Medication Adherence Interventions among HIV-Positive Patients at an Urban Clinic. *Hindawi Publishing Corporation AIDS Research and Treatment*, 2013. Retrieved from <http://www.hindawi.com/journals/art/2013/670525/>
- Minartiningtyas, B. (2012). SMS Gateway. Retrieved January 11, 2014, from <http://informatika.web.id/category/sms-gateway>
- Mohammed, S., Siddiqi, O., Ali, O., Habib, A., Haqqi, F., Kausar, A. (2012). User engagement with and attitudes towards an interactive SMS reminder system for patients with tuberculosis. *J Telemed Telecare*, 18 no. 7 4. doi:10.1258/jtt.2012.120311
- Mutire, B. N., Keraka, M. N., Kimuu, P. K., Kabiru, E. W., Ombeka, V. O. & Oguya, F. (2011). Factors associated with default from treatment among tuberculosis patients in Nairobi province, Kenya: a case control study. *BMC Public Health*, 11(1), 696. doi:10.1186/1471-2458-11-696
- Nglazi, M. D., Bekker, L.-G., Wood, R., Hussey, G. D. & Wiysonge, C. S. (2013). Mobile phone text messaging for promoting adherence to anti-tuberculosis treatment: a systematic review protocol. *Systematic reviews*, 2(1), 6. doi:10.1186/2046-4053-2-6
- Northouse, P. (1992). *Health Communication, Strategies For Health Professionals*. (William Brottmeier, Ed.) (Second Edi., p. 286). East Norwalk, Connecticut.
- Observatory, G. (2011). New horizons for health through mobile technologies, 3, 112. Retrieved from http://www.who.int/goe/publications/goe_mhealth_web.pdf
- Pablos-Méndez, Knirsch, Barr, F. (1997). Non adherence in Tuberculosis Treatment: Predictors and Consequences in New York City. *The American Journal of Medicine*, 102(2), 164–170. Retrieved from [http://www.amjmed.com/article/S0002-9343\(96\)00402-0/abstract?source=aemf](http://www.amjmed.com/article/S0002-9343(96)00402-0/abstract?source=aemf)

- Pakpahan, H. (2013). Pengertian SMS. <http://globalonlinebook.blogspot.com/>. Retrieved from <http://globalonlinebook.blogspot.com/2009/09/pengertian-sms.html>
- Pop-Elechesa, C., Habyarimanae, M., & Goldstein, M. P. (2011). Mobile Phone Technologies Improve Adherence to Antiretroviral Treatment in Resource-Limited Settings : A Randomized Controlled Trial of Text Message Reminders Mobile phone technologies improve adherence to antiretroviral treatment in a resource-limited se, (January). doi:10.1097/QAD.0b013e32834380c1
- Pratiknya, A. W. (1993). *Dasar-Dasar Metodologi Penelitian Kedokteran dan Kesehatan* (Ed.1., Cet., pp. 274–149). Jakarta Indonesia: PT. RajaGrafindo Persada.
- Ramadhika, A. (2012). SMS Gateway Menggunakan Gammu Dan MySQL. Retrieved January 11, 2014, from http://www.ubaya.ac.id/2013/content/articles_detail/33/SMS-Gateway-menggunakan-Gammu-dan-MySQL.html
- Ridha, A. (2012). *Motivasi Melalui Layanan Pesan Singkat Telepon Seluler Pada Perilaku Merokok*. Gadjah Mada University Yogyakarta, Yogyakarta.
- Rodrigues, R., Shet, A., Antony, J., Sidney, K., Arumugam, K., Krishnamurthy, S., D'Souza, G., (2012). Supporting adherence to antiretroviral therapy with mobile phone reminders: results from a cohort in South India. *PloS one*, 7(8), e40723. doi:10.1371/journal.pone.0040723
- Rogers, C. R. (1951). *Client-centered therapy*. Boston: Houghton Mifflin Company.
- Sastroasmoro, S. & Ismael, S. (1995). *Dasar-Dasar Metodologi Penelitian Klinis*. (S. Sastroasmoro, S & Ismael, Ed.) (Pertama., p. 283). Jakarta Indonesia: Binarupa Aksara, Jakarta.
- Scrhamm, W. (1969). *Mass Communications* (October 1s.). Illinois, USA: University of Illinois Press.
- Sembiring, T. (2014). Menkominfo: 270 Juta Pengguna Ponsel di Indonesia. Retrieved April 27, 2015, from <http://ugm.ac.id/id/berita/8776-menkominfo:270.juta.pengguna.ponsel.di.indonesia>
- Utomo, P. (2006). *Membangun Aplikasi SMS Berbasis Open Source*. Yogyakarta: Andi Offset.
- Vijay, S., Kumar, P., Chauhan, L. S., Vollepore, B. H., Kizhakkethil, U. P., & Rao, S. G. (2010). Risk factors associated with default among new smear positive TB

patients treated under DOTS in India. *PloS one*, 5(4), e10043.
doi:10.1371/journal.pone.0010043

WHO. (2009). *Treatment of tuberculosis: Guidelines for National Programs*.

WHO. (2013). Case TB Notification 2013. Retrieved April 28, 2015, from
http://apps.who.int/iris/bitstream/10665/137094/1/9789241564809_eng.pdf?ua=1

Wesolowski, A., Eagle, N., Noor, A. M., Snow, R. W., & Buckee, C. O. (2012). Heterogeneous Mobile Phone Ownership and Usage Patterns in Kenya, 7(4), 1–6. doi:10.1371/journal.pone.0035319

Widjanarko, B., Gompelman, M., Dijkers, M., & Van der Werf, M. J. (2009). Factors that influence treatment adherence of tuberculosis patients living in Java, Indonesia. *Dove Press Journal*, 3, 231–8. Retrieved from <http://www.dovepress.com>

World Health Organization. (2013). WHO Global Report. Geneva, Switzerland. Retrieved from <http://www.who.int/tb/publications/2009/airborne/background/info/en/>

Zignol, M., Gemert, W. V., Falzon, D., Sismanidis, C., Glaziou, P., Floyd, K., & Raviglione, M. (2012). Surveillance of anti-tuberculosis drug resistance in the world : an updated analysis , 2007 – 2010. *Bull World Health Organ* 2012;90:111–119D, (November 2011), 111–119. doi:10.2471/BLT.11.092585

Zurovac, D., Otieno, G., Kigen, S., Mbithi, A. M., Muturi, A., Snow, R. W., & Nyandigisi, A. (2013). Ownership and use of mobile phones among health workers, caregivers of sick children and adult patients in Kenya: cross-sectional national survey. *Globalization and health*, 9(1), 20. doi:10.1186/1744-8603-9-20

