



## DAFTAR PUSTAKA

- Abaza, I.M., M.A. Shehata, M.S. Shoieb, and I.I. Hasan. 2008. Evaluation of some natural feed additive in growing chicks diets. International Journal of Poultry Science. 7: 872-879.
- Abu-Dieyeh, B.H.M. and M.S. Abu-Darwish. 2008. Effect of feeding powder black cumin seeds (*Nigella sativa* L.) on growth performance of 4-8 weeks old broilers. Journal of Animal and Veterinary Advances. 7: 286-290.
- Ademola, S.G., G.O. Farinu, and G.M. Babatunde. 2009. Serum lipid and haematological parameters of broiler fed garlic, ginger and their mixtures. World Journal of Agricultural Sciences. 5: 99-104.
- Adjid, R.M.A., R. Damayanti, H. Hamid, T. Sjafriati, dan Darminto. 2002. Penyakit Marek pada ayam: I. Etiologi, patogenesis, dan pengendalian penyakit. Wartazoa. 12: 65-71.
- Aini, I. 1999. The Scope and Effect of Family Poultry Research and Development: Disease in rural family chickens in Southe-East Asia. International Network for Family Poultry Development (INFPD). INFPD E-Conference. FAO. 37-41.
- Aji, S.B., K. Ignatius, A.Y. Ado, J.B. Nuhu, A. Abdulkarim, U. Aliyu, M.B. Gambo, M.A. Ibrahim, H. Abubakar, M.B. Bukar, H.A.M. Imam and P.T. Numar. 2011. Effects of feeding onion (*Allium cepa*) and garlic (*Allium sativum*) some performance characteristics of broiler chickens. Research Journal of Poultry Science. 4: 22-27.
- Akhtar, M.S., Z. Nasir and A.R. Abid. 2003. Effect of feeding powdered *Nigella sativa* L. seed on poultry egg production and their suitability for human consumption. Veterinarski Arhiv. 73: 181-190.
- Al-Beitawi, N.A. and S.S. El-Ghousein. 2008. Effect of feeding different levels of *Nigella sativa* seeds (black cumin) on performance, blood constituents and carcass characteristic of broiler chicks. International Journal of Poultry Science. 7: 715-721.
- Al-Beitawi, N.A., S.S. El-Ghousein, and A.H. Nofal. 2009. Replacing bacitracin methylene disalicylate by crushed *Nigella sativa* seeds in broiler rations and its effects on growth, blood constituents and immunity. Livestock Science. 125: 304-307.
- Al-Munfareej, S.I. 2014. Immune-responiveness and performance of broiler chickens fed black cumin (*Nigella sativa* L.) powder. Journal of the Saudi Society of Agricultural Sciences. 13: 75-80.



- Al-Zahim, M., H.J. Newbury, and B.V. Ford-Lloyd. 1997. Classification of genetic variation in garlic (*Allium sativum L.*) revealed by RAPD. Hort Science. 32: 1102-1104.
- Alçılıçek, A., M. Bozkurt and M. Çabuk. 2003. The effect of an essential oil combination derived from selected herbs growing wild in Turkey on broiler performance. South African Journal of Animal Science. 33: 89-94.
- Anggorodi, R. 1985. Kemajuan Mutakhir dalam Ilmu Makanan Ternak Unggas. Penerbit Universitas Indonesia (UI-Press). Jakarta.
- Ao, X., J.S. Yoo, T.X. Zhou, J.P. Wang, Q.W. Meng, L. Yan, J.H. Cho, and I.H.Kim. 2011. Effects of fermented garlic powder supplementation on growth performance, blood profiles and breast meat quality in broilers. Livestock Science. 141: 85-89.
- Ashayerizadeh, O., B. Dastar, and M.S. Shargh. 2009. Use of garlic (*Allium sativum*), black cumin seeds (*Nigella sativa L.*) and wild mint (*Mentha longifolia*) in broiler chickens diets. Journal of Animal and Veterinary Advances. 8: 1860-1863.
- Astuti, M. 2008. Buku Ajar Rancangan Percobaan. Laboratorium Pemuliaan Ternak, Fakultas Peternakan, UGM. Yogyakarta.
- Aviagen. 2010. Lohmann Meat Broiler Stock Nutrition Specification. <http://en.aviagen.com>. Diakses pada 26 April 2014.
- Aviagen. 2012. Ross 308 Broiler: Performance Objectives. <http://en.aviagen.com>. Diakses pada 26 April 2014.
- Aydin, R., M.A. Bal, A.K. Ozugur, H.H.C. Toprak, A. Kamalak and M. Karaman. 2006. Effect of black seeds (*Nigella sativa*) supplementation on feed efficiency egg yield parameters and shell quality in chickens. Pakistan Journal of Biological Sciences. 9: 243-247.
- Bashandy, S.A.E. 2007. Effect of fixed oil of *Nigella sativa* on male fertility in normal and hyperlipidemic rats. International Journal of Pharmacology. 3 : 27-33.
- Begum, M., M.M. Hossain, and I.H. Kim. 2014. Effects of the plant extract YGF251 on growth performance, meat quality, relative organ weight, nutrient digestibility and blood profiles in broiler chickens: possible role of insulin-like growth factor 1. Veterinarski Medicina. 59: 415-423.
- Berg, T.P.V. 2010. Acute infectious bursal disease in poultry: a review. Avian Pathology. 29: 175-194.



- Botsoglou, N.A., E. Christaki, P.F. Paneri, I. Gianennas, G. Papagiorgiou, and A.B. Spaus. 2004. The effect of a mixture of herbal essential oils or  $\alpha$ -tocopheryl acetate on performance parameters and oxidation of body lipid in broilers. South Africa Journal of Animal Science. 34: 52-61.
- Clark, S., R. Daly, E. Jordan, E. Lee, A. Mathew, and P. Ebner. 2011. Extention Education Symposium: The future of biosecurity and antimicrobial use in livestock production in the united states and the role of extention. Journal of Animal Science. 90: 2861-2872.
- Damayanti, R. Dan A. Wiyono. 2003. Gambaran histopatologi kasus Marek pada ayam pedaging di Kabupaten Tasikmalaya dan Ciamis, Jawa Barat. JJIV. 8: 247-255.
- Demir, E., Ş. Sarica, M.A. Özcan, and M.S. Mez. 2003. The use of natural feed additives as alternatives for an antibiotic growth promoter in broiler diets. British Poultry Science. 44: 44-45.
- Dibner, J.J. and J.D. Richards. 2005. Antibiotic growth promoters in agriculture: history and mode of action. Poultry Science. 84: 634-643.
- Doley, S., J.J. Gupta, and P.B. Reddy. 2009. Effect of supplementation of ginger, garlic and turmeric in broiler chicken. Indian Veterinary Journal. 86: 644 (Abstr.)
- Dono, N.D. 2012. Nutritional strategies to improve enteric health and growth performance of poultry in the post of antibiotic era. Ph D thesis. Colledge of Medical, Veterinary and Life Sciences, University of Glasgow.
- Durrani, F.R., N. Chand, K. Zaka, A. Sultan, F.M. Khattak and Z. Durrani. 2007. Different levels of feed added black seed (*Nigella sativa L.*) on the performance of broiler chicks. Pakistan Journal of Biological Sciences. 10: 4164-4167.
- Erdoğan, Z., S. Erdoğan, Ö. Aslantaş and S. Çelik. 2010. Effects of dietary supplementation of synbiotics and phytobiotics on performance, caecal coliform population and some oxidant/antioxidant parameters of broilers. Journal of Animal Physiology and Animal Nutrition. 94: 40 (Abstr.).
- Fadlala, I.M.T., B.H. Mohammed, and A.O. Bakhet. 2010. Effect of feeding garlic on the performance and immunity of broilers. Asian Journal of Poultry Science. 4: 182-189.



- Giannenas, I., P. Florou-Paneri, M. Papazahariadou, E. Christaki, N.A. Botsoglou, and A.B. Spais. 2003. Effect of dietary supplementation with oregano essential oil on performance of broilers after experimental infection with *Eimeria tenella*. Archives of Animal Nutrition. 57: 99-106.
- Guler, T., B. Dalkılıç, O.N. Ertas and M. Çiftçi. 2006. The effect of dietary black cumin seeds (*Nigella sativa* L.) on the performance of broilers. Asian-Australian Journal of Animal Science. 19: 425-430.
- Halawani, E. 2009. Antibacterial activity of thymoquinone and thymohydroquinone of *Nigella sativa* L. and their interaction with some antibiotics. Advances in Biological Research. 3: 148-152.
- Hartadi, H., S. Reksohadiprodjo, S. Lebdosukojo, and A.D. Tillman. 1980. Tabel-Tabel dari Komposisi Bahan Makanan Ternak untuk Indonesia. Program EFD. Fakultas Peternakan, Universitas Gadjah Mada.
- Hashemi, S.R. and H. Davoodi. 2010. Phytopreventive as new class of feed additive in poultry industry. Journal of Animal and Veterinary Advances. 9: 2295-2304.
- Horton, G.M.J., M.J. Fennell, and B.M. Prasad. 1991. Effect of dietary garlic (*Allium sativum*) on performance, carcass composition and blood chemistry changes in broiler chickens. Canadian Journal of Animal Science. 71: 939-942.
- Ismail, Z.S.H. 2011. Effects of dietary black cumin growth seeds (*Nigella sativa* L.) or its extract on performance and total coliform bacteria count on broiler chicks. Research Article, Animal Production Department, Faculty of Agriculture, South Valley University. Egypt.
- Kamran, Z., M. Sarwar, M. Nisa, M.A. Nadeem, S. Mahmood, M.E. Babar, and S. Ahmed. 2008. Effect of low-protein diets having constant energy-to-protein ratio on performance and carcass characteristics of broiler chickens from one to thirty-five days of age. Poultry Science. 87: 468-474.
- Khalaji, S., M. Zaghari, K.H. Hatami, S. Hedari-Dastjerdi, L. Lotfi, and H. Nazarian. 2011. Black cumin seeds, *Artemisia* leaves (*Artemisia sieberi*), and *Camellia* L. plant extract as phytopreventive products in broiler diets and their effects on performance, blood constituents, immunity, and cecal microbial population. Poultry Science. 90: 2500-2510.



- Khawaja, T.; S.H Khan; N. Mukhtar and A. Parveen. 2012. Comparative study of growth performance, meat quality and haemotological parameters of Fayoumi, Rhode Island red and their reciprocal crossbred chickens. Italian Journal of Animal Science. 11: 211-216.
- Kırkpınar, F., H.B. Unlu, and G. Ozdemir. 2010. Effects of oregano and garlic essential oils on performance, carcass, organ and blood characteristics and intestinal microflora of broilers. Livestock Science. 137: 219-225.
- Kökdil, G. and H. Yilmaz. 2005. Analysis of the fixed oils of the genus *Nigella* L. (Ranunculaceae) in Turkey. 33: 1203-1209.
- Lee, S.H., H.S. Lillehoj, R.A. Dalloul, D.W. Park, Y.H. Hong, and J.J. Lint . 2007. Influence of *Pediococcus*-based probiotic on coccidiosis in broiler chickens. Poultry Science. 86: 63-66.
- Lević, J., S. Sredanovic, O. Đuragić, D. Jakić, Lj. Lević, and S. Pavkov. 2007. New feed additives based on phytogenics and acidifiers in animal nutrition. Biotechanology in Animal Husbandry. 23: 527-534.
- Mansoub, N.H. and M.P. Myandoab. 2011. Comaparative effect of using zizaphora (*Thymus vulgaris*), garlic and probiotic on performance and serum composition of broiler chickens. Scholars Research Libary. 2: 373-378.
- Matsuyama-Kato, A. , S. Murata, M. Isezaki, R. Kano, S. Takasaki, O. Ichii, S. Konnai, and K. Ohashi. 2012. Molecular characterization of immunoinhibitory factors PD-1/PD-L1 in chickens infected with Marek's disease virus. Virology Journal. 9: 1-12.
- Miraghaei, S.S., B. Heidary, H. Almasi, A. Shabani, M. Elahi, dan M.H.M. Nia. 2011. The effects of *Nigella sativa* powder (black seed) and *Echinacea purpurea* (L.) moench extract on performance, some blood biochemical and hematological parameters in broiler chickens. African Journal of Biotechnology. 10: 19249-19254.
- Mwale, M., J.F. Mupangwa, C. Mapiye, H. Saina, and J. Chipuramahwe. 2008. Growth performance of guinea fowl keets fed graded levels of baobab seed cake diets. International Journal of Poultry Science. 7: 429-432.
- Natsir, M.H., Hartutik, O. Sofjan and E. Widodo. 2013. Effect of either powder or encapsulated form of garlic and *Phylantus Niruri* L. mixture on broiler performances, intestinal characteristics and intestinal microflora. International Journal of Poultry Science.12 : 676-680.



- Nobo, G., J. Moreki, and N.J. Nsoso. 2012. Feed intake, body weight, average daily gain, feed conversion ratio and carcass characteristic of hemelted guinea fowl fed varying levels of phane meal (*Imbrasia belina*) as replacement of fishmeal under intensive system. International Journal of Poultry Science. 11: 378-384.
- Oguz, H. and V. Kurtoglu. 2000. Effect of clinoptilolite on performance of broiler chickens during experimental aflatoxicosis. British Poultry Science. 41: 512-517.
- Okwor, E.C. and D.C. Eze. 2011. Outbreak and persintance of Marek's disease in batches of birds reared in a poultry farm located in Nsukka, South East Nigeria. International Journal of Poultry Science. 10: 617-620.
- Onibi, G.E., O.E. Adebisi, A.N. Fajemisin, and A.V. Adetunji. 2009. Response of broiler chicken in terms of performance and meat quality to garlic (*Allium sativum*) supplementation. African Journal of Agricultural Research. 4: 511-517.
- Prasad, R., M.K. Rose, M. Virmani, S.L. Garg, and J.P. Puri. 2009. Lipid profile of chicken (*Gallus domesticus*) in response to dietary supplementation of garlic (*Allium sativum*). International Journal of Poultry Science. 8: 270-276.
- Raeesi, M., S.A.H. Alibabad, A. Roofchae, A.Z. Shahneh, and S. Pirali. 2010. Effect of periodically use of garlic (*Allium sativum*) powder on performance and carcass characteristics in broiler chicken. International Science Index. 4: 1421-1427.
- Rahimi, S., Z.T. Zadeh, M..K. Torshizi, R. Omidbaigi, and H. Rokni. 2011. Effect of the three herbal extracts on growth performance, immune system, blood factors, and intestinal selected bacterial population in broiler chickens. Journal Agriculture Science Technology. 13: 527-539.
- Roofchae, A., M. Irani, M.A. Ebrahimzadeh and M.R. Akbari. 2011. Effect of dietary oregano (*Origanum vulgare L.*) essential oil on growth performance, cecal microflora and serum antioxidant activity of broiler chickens. African Journal of Biotechnology. 10: 6177-6183.
- Salem, M.L. 2005. Immunomodulatory and therapeutic properties of the *Nigella sativa* L. seed. International Immunopharmacology. 5: 1749-1770.



- Shewita, R.S. and A.E. Taha. 2011. Effect of dietary supplementation of different levels of black seeds (*Nigella sativa* L.) on growth performance, immunological, hematological and carcass parameters of broiler chickens. World Academy of Science, Engineering and Technology. 5: 1071-1077.
- Sirvydis, H.V., R. Bobiniené, V. Priudokiené, and D. Vencius. 2003. Phyto-biotics add value to broiler feed. World Poultry. 19: 16-17.
- Tabbu, C. R. 2000. Penyakit Ayam dan Penanggulangannya: Penyakit Bakterial, Mikal, dan Viral volume 1. Penerbit Kanisius. Yogyakarta.
- Temburne, S.T., S. Feroz, B.H. More, and D.M. Sakarkar. 2014. A review on therapeutic potential of *Nigella sativa* (kalonji) seeds. Journal of Medicinal Plants Research. 8: 167-177.
- Wang, J.P., J.S. Yoo, H.D. Jang, J.H. Lee, J.H. Cho, and I.H. Kim. 2011. Effect of dietary fermented garlic by *Weisella koreensis* powder on growth performance, blood characteristic and immune response of growing pigs challenged with *Escherichia coli* lypopolysacharide. Journal of Animal Science. 89: 2123-2131.
- Windisch, W., K. Schedle, C. Plitzner, and A. Kroismayr. 2007. Use of phytogenic product as feed additives for swine and poultry. Journal of Animal Science. 86: 140-148.
- Yuwanta, T. 2000. Dasar Ternak Unggas. Fakultas Peternakan, Universitas Gadjah Mada. Yogyakarta.
- Ziarlarimi, A., M. Irani, S. Gharahveysi, and Z. Rahmani. 2011. Investigation of antibacterial effects of garlic (*Allium sativum*), mint (*Mentha* spp.) and Onion (*Allium cepa*) herbal extracts on *Escherichia coli* isolated from broiler chickens. African Journal of Biotechnology. 10: 10320-10322.
- Zhao, J.F., J.L. Chen, G.P. Zhao, M.Q. Zheng, R.R. Jiang, and J. Wen. 2009. Live performance, carcass composition, and blood metabolite responses to dietary nutrient density in two distinct broiler breeds of male chickens. Poultry Science. 88: 2575-2584.