UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Alfred. D.C., Mohsin. S.K., Fuqua. S.K., 2001. Histopathological dan Biological Evolution of Human Premalignant of Breast Disease. In: Endocrine Related Cancer Vol 8. P 47-61

American Cancer Society. 2016. Breast cancer facts dan figures 2015 - 2016. American Society Inc. Atlanta.

American Cancer Society. 2007. Global Cancer Facts dan Figures 2007. American Society Inc. Atlanta.

American Joint Committee on Cancer. 2009. AJCC CANCER STAGING ATLAS. Springer Science+Business Media, Inc.

Andre F, Cabioglu N, Assi H, Sabourin JC, Delaloge S, Sahin A, Broglio K, Spano JP, Combadiere C, Bucana C, Soria JC, Cristofanilli M. 2006. Expression of chemokine receptors predicts the site of metastatic relapse in patients with axillary node positive primary breast cancer. *Ann Oncol* 2006, 17(6):945–951.

Arens N, Gandhari M, Bleyl U, Hildenbrdan R. 2005. In vitro suppression of urokinase plasminogen activator in breast cancer cells–a comparison of two antisense strategies. *Int J Oncol,* 26(1):113–119.

Barbero S, Bonavia R, Bajetto A, Porcile C, Pirani P, Ravetti JL,Zona GL, Spaziante R, Florio T, Schettini G. 2003. Stromal cell-derived factor 1_ stimulates human glioblastoma cell growth through the activation of both extracellular signal-regulated kinases 1/2 dan Akt. In: Cancer Research vol 63. P 1969–1974

Barcellos-Hoff M.H, dan Ravani S.A. 2000. Irradiated mammary gldan stroma promotes the expression of tumorigenic potential by unirradiated epithelial cells. In: Cancer Research vol 60. P 1254.

Bartolomé RA, Gálvez BG, Longo N, Baleux F, Van Muijen GN, Sánchez-Mateos P, Arroyo AG, Teixidó J. 2004. Stromal cell-derived factor-1alpha promotes melanoma cell invasion across basement membranes involving stimulation of membrane-type 1 matrix metalloproteinase and Rho GTPase activities. Cancer Res;64:2534–43.

Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein
Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan
Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Behrens J. 1993. The role of cell adhesion molecules in cancer invasion dan metastasis. *Breast Cancer Res Treat*, 24(3):175–184.

Bhowmick,N,A, Neilson. E.G., and Moses.H.L, 2004 Stromal fibroblasts in cancer initiation and progression *Natur*e. 2004 Nov 18; 432(7015): 332–337. doi: 10.1038/ nature03096

Blot. E, Laberge-Le Couteulx S, Jamali. H, Cornic M, Guillemet. C, Duval. C, Hellot MF, Pille. JY, Picquenot JM, Veyret C. 2008. CXCR4 membrane expression in node-negative breast cancer. Breast J. 2008 May-Jun;14(3):268-74. doi: 10.1111/ j.15244741.2008.00573.x. Epub 2008 Mar 26.

Bodenstine. T &. Welch. D. 2008. Metastasis Suppressors dan the Tumor Microenvironment. In: *Cancer Microenvironment*.. Springer Science Business Media. P 1–11

Bockhorn M, Jain RK, Munn LL. Active versus passive mechanisms in metastasis: do cancer cells crawl into vessels, or are they pushed? *Lancet Oncol*. 2007 May;8(5):444-8.

Bramwell VH, Doig GS, Tuck AB, Wilson SM, Tonkin KS, Tomiak A, Perera F, Vandenberg TA, Chambers AF. 2006. Serial plasma osteopontin levels have prognostic value in metastatic breast cancer. Clinical Cancer Research, 12(11 Pt 1):3337–3343.

Burger.J dan Kipps. T. 2006 CXCR4: a Key Receptor in the Crosstalk Between Tumor Cells dan Their Microenvironment. In: *BLOOD*, Vol 107,no 5. The American Society of Hematology. P 1761-1767

Burstein.H.J, Jay. R.H., Morrow.M. 2008 Section 2: Malignant Tumors of the Breast. In Devita, Hellman & Rosenberg's Cancer: Principles & Practice of Oncology, 8th Edition. Editors: DeVita, Vincent T.; Lawrence, Theodore S.; Rosenberg, Steven A. Copyright 2008 Lippincott Williams & Wilkins

Bushillo, J.M. dan Benovic, J.L. 2008. Regulation of CXCR4 Signaling. *Biochim Biophys Acta.* 2008 April ; 1768(4): 952–963.

Bushillo, J.M., Armando, S., Sengupta,. R., Meucci, O. Bouvier, M., dan Benovic, J.L.. 2010. Site-specific Phosphorylation of CXCR4 Is Dynamically Regulated by Multiple Kinases and Results in Differential Modulation of CXCR4 Signaling. *Journal Biology Chem*. 2010 Mar 5; 285(10): 7805–7817. Published online 2010 Jan 4. doi: 10.1074/jbc.M109.091173

Bustreo.S, Osella-Abate.S, Cassoni,P, Donadio.M, Airoldi.M, Pedani.F, Papotti.M,

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Sapino.A. Castellano.I. 2016. Optimal Ki67 cut-off for luminal breast cancer prognostic evaluation: a large case series study with a long-term follow-up. *Breast Cancer Res Treat* (2016) 157:363–371 DOI 10.1007/s10549-016-3817-9

Carmeliet P, Jain RK. 2000. Angiogenesis in cancer and other diseases. Nature; 407:249–57.

Ceradini DJ, Kulkarni AR, Callaghan MJ, Tepper OM, Bastidas N, Kleinman ME, Capla JM, Galiano RD, Levine JP, Gurtner GC. 2004. Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. In: Nat Med vol 10.P 858–864

Chang HY, Chi. J.T., Dudoit. S., Bondre. C., Rijn. M.V., Botstein. D., dan Brown. P.O. 2002. Diversity, Topographic Differentiation, dan Positional Memory in Human Fibroblasts. In: Proc Natl Acad Sci U S A vol. 99 no. 20. P 12877-12882

Chang HY, Sneddon. J.B., Alizadeh. A.A., Sood. R., West. R.B., Montgomery. K., Chi.J.T, Rijn. M.V., Botstein. D., dan Brown. P.O. 2004. Gene Expression Signature of Fibroblast Serum Response Predicts Human Cancer Progression: Similarities Between Tumors dan Wounds. In: PLoS Biol Vol 2 issue 2. P 206-2014

Chang YS, di Tomaso E, McDonald DM, Jones R, Jain RK, Munn LL. 2000. Mosaic blood vessels in tumors: frequency of cancer cells in contact with flowing blood. Proc Natl Acad Sci USA ;97:14608–13.

Cheng. J.D. dan Weiner. L.M. 2003. The Biology Behind Tumors dan Their Microenvironments: Tilling the Soil. In: Clinical Cancer Research Vol. 9. P, 1590–1595

Cook AC, Tuck AB, McCarthy S, Turner JG, Irby RB, Bloom GC, Yeatman TJ, Chambers AF. 2005. Osteopontin induces multiple changes in gene expression that reflect the six “hallmarks of cancer” in a model of breast cancer progression. Mol Carcinog, 43(4):225–236.

Costa. A, Coradini. D, Carrasi. A, Erdas. R, Sardella. A, dan Daidone. M, 2001. Re: Levels of Hypoxia-Inducible Factor-1 During Breast Carcinogenesis. In: Journal of the National Cancer Institute, Vol. 93, no. 15, August 1. P 1175-1177

Cui Y, Niu A, Pestell R, Kumar R, Curran EM, Liu Y, Fuqua SA. 2006. Metastasis associated protein 2 is a repressor of estrogen receptor alpha whose overexpression leads to estrogen-independent growth of human breast cancer cells. Mol Endocrinol, 20(9):2020–2035.

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Darash-Yahana M, Pikarsky E, Abramovitch R, Zeira E, Pal B, Karplus R, Beider K, Avniel S, Kasem S, Galun E, Peled A. 2004. Role of high expression levels of CXCR4 in tumor growth, vascularization, dan metastasis. In: FASEB Journal vol 18.P 1240–1242,

de Silva Rudldan S, Martin L, Roshanlall C, Winstanley J, Leinster S, Platt- Higgins A, Carroll J, West C, Barraclough R, Rudldan P. 2006. Association of S100A4 dan osteopontin with specific prognostic factors dan survival of patients with minimally invasive breast cancer. Clin Cancer Res, 12(4):1192–1200.

Duffy MJ, Maguire TM, Hill A, McDermott E, O'Higgins N, 2000. Metalloproteinases: role in breast carcinogenesis, invasion dan metastasis. Breast Cancer Res, 2(4):252–257.

Duffy MJ, McCarthy K,1998. Matrix metalloproteinases in cancer: prognostic markers dan targets for therapy (review). Int J Oncol, 12(6):1343–1348

Dunnwald.L.K., Rossing.M.A. and Li.C.I. 2007 Hormone receptor status, tumor characteristics, and prognosis: a prospective cohort of breast cancer patients. Breast Cancer Research. 2007 9:R6 DOI: 10.1186/bcr1639

Egelbald .M., Littlepage. L.E., and Werb. Z..The Fibroblastic Coconspirator in Cancer Progression. Cold Spring Harb Symp Quant Biol. 2005; 70: 383–388. doi: 10.1101/sqb.2005.70.007

England National Statistics on request 2015. http://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/breast-cancer/incidence-invasive

Entschladen F, Drell TL 4th, Lang K, Joseph J, Zaenker KS. 2004. Tumour-cell migration, invasion, and metastasis: navigation by neurotransmitters. Lancet Oncol;5:254–58.

Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray, F. 2014. Cancer incidence & mortality worldwide: Sources, methods & major patterns in GLOBOCAN 2012. Available from: http://globocan.iarc.fr 15 September

Ferlay J, Autier P, Boniol M, Heanue.M , Colombet.M & Boyle.P.. 2007. Estimates of the Cancer Incidence dan Mortality in Europe in 2006. In: Ann Oncology vol 18 no 3: P 581.

Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein
Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan
Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Finlay TH, Tamir S, Kadner SS, Cruz MR, Yavelow J, Levitz M. 1993. Alpha 1-Antitrypsin- dan anchorage-independent growth of MCF-7 breast cancer cells. Endocrinology, 133(3):996–1002.

Fitzgibbons PL, Page DL, Weaver D, Thor AD, Allred DC, Clark GM, Ruby SG, O'Malley F, Simpson JF, Connolly JL, Hayes DF, Edge SB, Lichter A, Schnitt SJ. 2000. Prognostic factors in breast cancer. College of American Pathologists Consensus Statement 1999. *Arch Pathol Lab Med*. 2000 Jul;124(7):966-78.

Foekens JA, Ries C, Look MP, Gippner-Steppert C, Klijn JG, Jochum M, 2003. Elevated expression of polymorphonuclear leukocyte elastase in breast cancer tissue is associated with tamoxifen failure in patients with advanced disease. Br J Cancer, 88(7):1084–1090.

Foekens JA, Ries C, Look MP, Gippner-Steppert C, Klijn JG, Jochum M. 2003. The prognostic value of polymorphonuclear leukocyte elastase in patients with primary breast cancer. Cancer Res, 63(2):337–341.

Gandhari M, Arens N, Majety M, Dorn-Beineke A, Hildenbrdan R. 2006. Urokinasetype plasminogen activator induces proliferation in breast cancer cells. Int J Oncol, 28(6):1463–1470.

Genatlas. 2007. CXCL12. http://genatlas.medecine.univ-paris5.fr/

Genatlas. 2007. CXCR4. http://genatlas.medecine.univ-paris5.fr/

Giannelli G, Erriquez R, Fransvea E, Daniele A, Trerotoli P, Schittulli F, Grano M, Quaranta M, Antonaci S. Proteolytic imbalance is reversed after therapeutic surgery in breast cancer patients. *Int J Cancer* 2004, 109(5):782–785

Gilles C, Bassuk JA, Pulyaeva H, Sage EH, Foidart JM, Thompson EW. SPARC/osteonectin induces matrix metalloproteinase 2 activation in human breast cancer cell lines. Cancer Res 1998, 58(23):5529–5536.

Gnant.M, Harbeck. N, dan Thomssen. C. 2011. St Gallen 2011: Summary of the Consensus Discussion. Breast Care 2011;6:136-141 DOI: 10.1159/000328054

Graham JD, Balleine RL, Milliken JS, Bilous AM, Clarke CL. 1997. Expression of osteonectin mRNA in human breast tumours is inversely correlated with oestrogen receptor content. Eur J Cancer, 33(10):1654–1660.

Grigorian MS, Ambartsumian NS, Georgiev GP, Lukanidin EM. 1999. (Expression of mts1 gene in human breast cancer MCF-7 cells increases their malignancy). Mol Biol (Mosk), 33(4):651–656.

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Grimes.DA, Schulz. KF..2002. Cohort studies: Marching towards outcomes. *The Lancet*; Jan 26, 2002; 359, 9303; ProQuest Nursing & Allied Health Source pg. 341

Gutierrez LS, Eliza M, Niven-Fairchild T, Naftolin F, Mor G. 1999. The Fas/Fasligdan system: a mechanism for immune evasion in human breast carcinomas. Breast Cancer Res Treat, 54(3):245–253.

Hammond.ME, Hayes.DF, Dowsett,M, Allred.DC, Hagerty.KL, Badve.S, Fitzgibbons.PL, Francis.G, Goldstein. NS, Hayes. Hicks. MD, Lester.S, Love. R, Mangu.PB, McShane.L, Miller.K, Osborne.CK, Paik.S, Perlmutter. J, Rhodes.A, Sasano.H , Schwartz. J.N, Sweep.FC, Taube.S, Torlakovic.EE, Viale. PV, Visscher.D, Wheeler. T, Williams.RB, Wittliff.JL, dan Wolff, AC. 2010. American Society of Clinical Oncology/College of American Pathologists Guideline Recommendations for Immunohistochemical Testing of Estrogen and Progesterone Receptors in Breast Cancer. *Journal of Clinical Oncology ASCO Special Article*. Vol 28 Numbar 16 June1 2010

Hartmann TN, Grabovsky V, Pasvolsky R, Shulman Z, Buss EC, Spiegel A, Nagler A, Lapidot T, Thelen M, Alon R. 2008. A crosstalk between intracellular CXCR7 and CXCR4 involved in rapid CXCL12-triggered integrin activation but not in chemokine-triggered motility of human T lymphocytes and CD34+ cells. J Leukoc Biol. 2008;84:1130–1140

Hassan S, Ferrario C, Saragovi U, Quenneville L, Gaboury L, Baccarelli A, Salvucci O, Basik M. The influence of tumor-host interactions in the stromal cell-derived factor-1/CXCR4 ligand/receptor axis in determining metastatic risk in breast cancer. Am J Pathol. 2009 Jul;175(1):66-73. doi: 10.2353/ajpath.2009.080948. Epub 2009 Jun 4.

Hortobagyi G.N., Singletary S.E., Strom E.A. 2004. Locally Advanced Breast Cancer. In Harris J.R.,. Editors. *Disease of The Breast. 3rd Edition*. Lippincott William Wilkins. Philadelphia. 58 : 951 – 969.

Horuk R. 2001 Chemokine receptors. Cytokine Growth Factor Rev. 2001 Dec;12(4):313-35.

Inoue T., Plieth D, Venkov C.D., Xu C., dan Neilson E.G. 2005. Antibodies against macrophages that overlap in specificity with fibroblasts. In: *Kidney International* vol 67. P 2488.

Isaac P. Witz. 2009. The Tumor Microenvironment: The Making of a Paradigm. In: *Cancer Microenvironmen*t vol 2 Suppl 1. P 9–17

Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein
Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan
Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Jain RK. 2003. Molecular regulation of vessel maturation. *Nat Med*;9:685–93.

Jenkinson SR, Barraclough R, West CR, Rudldan PS. 2004. S100A4 regulates cell motility dan invasion in an in vitro model for breast cancer metastasis. *Br J Cancer*, 90(1):253–262

Kang. H., Watkins.G., Parr.C., Douglas-Jones. A., Mansel. R.E., and Jiang. W.G. 2005. Stromal cell derived factor-1: its influence on invasiveness and migration of breast cancer cells in vitro, and its association with prognosis and survival in human breast cancer. *Breast Cancer Research*. 20057:R402 DOI: 10.1186/bcr1022© licensee BioMed Central Ltd. 2005

Kang H, Watkins G, Douglas-Jones A, Mansel RE, Jiang WG. 2005. The elevated level of CXCR4 is correlated with nodal metastasis of human breast cancer. *Breast*. 2005 Oct;14(5):360-7.

Kang H, Escudero-Esparza A, Douglas-Jones A, Mansel RE, Jiang WG. 2009. Transcript analyses of stromal cell derived factors (SDFs): SDF-2, SDF-4 and SDF-5 reveal a different pattern of expression and prognostic association in human breast cancer. *Int J Oncol*. 2009 Jul;35(1):205-11.

Katayama A, Ogino T, Bandoh N, Nonaka S, Harabuchi Y. 2005. Expression of CXCR4 dan its down-regulation by IFN-_ in head dan neck squamous cell carcinoma. In: *Clinical Cancer Research* vol 11. P 2937–2946,

Katayama Y, Battista M, Kao WM, Hidalgo A, Peired AJ, Thomas SA, Frenette PS. 2006. Signals from the sympathetic nervous system regulate hematopoietic stem cell egress from bone marrow. In: *Cell* vol 124. P 407–421,

Katz MH, Hauck WW. 1993. Proportional hazards (Cox) regression. *Journal Gen Intern Med.* 1993 Dec;8(12):702-11.

Kim R, Emi M, Tanabe K, Arihiro K. 2006. Tumor-driven evolution of immunosuppressive networks during malignant progression. Cancer Res, 66(11):5527–5536.

Kim. J.O, Suh. K.S, Lee. D.O, Sul. H.J, Lee. J.U, dan Song. K.S. 2009. Expression of CXCR4 dan SDF-1 in Primary Breast Cancer dan Metastatic Lymph Nodes. In: *Journal of Breast Cancer* vol 12 no 4. P 249-256.

Kryczek I, Wei S, Keller E, Liu R, Zou W. Stromal-derived factor (SDF-1/CXCL12) and human tumour pathogenesis. *Am Journal Physiol*. 2007;292:C987–C995

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Kucia. M, Jankowski. K, Reca. R, Wysoczynski. M, Bandura. L, Allendorf. D , Zhang. J, Ratajczak. J dan Ratajczak. M. 2004. CXCR4–SDF-1 Signalling, Locomotion, Chemotaxis dan Adhesion. In: *Journal of Molecular Histology*. 35: P 233–245

Kumar R, Wang RA, Bagheri-Yarmdan R. 2003. Emerging roles of MTA family members in human cancers. *Semin Oncol*, 30(5 Suppl 16):30–37.

Larson CJ, Moreno JG, Pienta KJ, Gross.S, Repollet. M, O'hara. SM, Russel. T, Terstappen.LW. 2004. Apoptosis of circulating tumor cells in prostate cancer patients. *Cytometry*;62:46–53.

Lee WY, Su WC, Lin PW, Guo HR, Chang TW, Chen HH. 2004. Expression of S100A4 dan Met: potential predictors for metastasis dan survival in early-stage breast cancer. *Oncology*, 66(6):429–438.

Lee. B.C., Lee. T.H., Avraham. S., dan Avraham. H.K. 2004. Involvement of the Chemokine Receptor CXCR4 dan Its Ligdan Stromal Cell-Derived Factor 1α in Breast Cancer Cell Migration Through Human Brain Microvascular Endothelial Cells. In: *Molecular Cancer Research* vol 2 no6. P 327-338

Liang Z, Yoon Y, Votaw J, Goodman M, William L, Shim H. Silencing of CXCR4 blocks breast cancer metastasis. Cancer Res. 2005;65:967–971.

Liang Z, Wu T, Lou H, Yu X, Taichman RS, Lau SK, Nie S, Umbreit J, Shim H. 2004. Inhibition of breast cancer metastasis by selective synthetic polypeptide against CXCR4. *Cancer Res*. 2004; 64:4302–4308

Hernandez.L., Magalhaes.M.A., Coniglio.S.J., Condeelis. J.S. & Segal.J.E. 2011. Opposing roles of CXCR4 & CXCR7 in breast cancer metastasis. *Breast Cancer Research*. 201113:R128. BioMed Central Ltd

Littlepage.L, Egeblad.M, dan Werb.Z. 2008. The Tumor Microenvironment in Cancer Progression. In: *The Molecular Basic of Cancer* 3rd Edition. Elsevier Inc. Philadelphia. P 229-239.

Liu F, Lang R, Wei J, Fan Y, Cui L, Gu F, Guo X, Pringle GA, Zhang X, Fu L. 2009. Increased expression of SDF-1/CXCR4 is associated with lymph node metastasis of invasive micropapillary carcinoma of the breast. *Histopathology*. 2009 May;54(6):741-50. doi: 10.1111/j.1365-2559.2009.03289.x.

Ma Q, Jones D, Borghesani PR, Segal RA, Nagasawa T, Kishimoto T, Bronson RT, Springer TA. 1998. Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- dan SDF-1-deficient mice. In: *PNAS* vol 95 no 16. P 9448–9453.

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Mansel. R, Fodstad. O, dan Jiang. W. 2007. Metastasis of breast cancer: an introduction. In: *Metastasis Breast Cancer, Cancer Metastasis – Biology dan Treatment* Vol 11. Springer. Dordrecht. P 1-7

Manuaba. T. W. 2010. *Panduan Penatalaksanaan Kanker Solid Peraboi* 2010. Sagung Seto. Jakarta

Martin TA, Watkins G, Mansel RE, Jiang WG. 2004. Hepatocyte growth factor disrupts tight junctions in human breast cancer cells. *Cell Biol Int*, 28(5):361–371.

Matteucci E, Locati M, Desiderio MA. 2005. Hepatocyte growth factor enhances CXCR4 expression favoring breast cancer cell invasiveness. *Exp Cell Res*, 310(1):176–185.

Mukherjee dan Zhao. 2013. Review Article The Role of chemokine receptor CXCR4 in breast cancer metastasis. *American Journal Cancer Research*. 2013;3(1):46-57 www.ajcr.us /ISSN:2156-6976/ajcr0000162

Mine S, Fujisaki T, Kawahara C, Tabata T, Iida T, Yasuda M, Yoneda T, Tanaka Y. 2003. Hepatocyte growth factor enhances adhesion of breast cancer cells to endothelial cells in vitro through up-regulation of CD44. *Exp Cell Res*, 288(1):189–197.

Mirisola. V, Zuccarino, A, Bachmeier. BR, Sormani. MP, Falter. J, Nerlich. A, Pfeffer. U. 2009. CXCL12/SDF1 expression by breast cancers is an independent prognostic marker of disease-free dan overall survival. In: *European Journal of Cancer* 45 : 2579 –2587

Mohle R, Bautz F, Rafii S, Moore MA, Brugger W, Kanz L. 1998. The chemokine receptor CXCR-4 is expressed on CD34_ hematopoietic progenitors dan leukemic cells dan mediates transendothelial migration induced by stromal cell-derived factor-1. In: *Blood* vol 91. P 4523-4530.

Muller A, Homey B, Soto H,Ge. N, Catron. D., Buchanan. M.E., Clanahan. T.M, Murphy. E., Yuan. W, Wagner. S.N., Barrera. J.L., Mohar. A., Verástegui. E., dan Zlotnik. A. 2001. Involvement of Chemokine Receptors in Breast Cancer Metastasis. In: *Nature* vol 410. P 50-56

Nagasawa .T., Hirota. S., Tachibana. K., Takakura .N., Nishikawa. S., Kitamura. Y., Yoshida. N., Kikutani. H., dan Kishimoto. T. 1996. Defects of B-cell lymphopoiesis dan bone-marrow myelopoiesis in mice lacking the CXC chemokine PBSF/SDF-1. In: *Nature* vol 382. P 635-638

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Nicolson GL, Nawa A, Toh Y, Taniguchi S, Nishimori K, Moustafa A. 2003. Tumor metastasis-associated human MTA1 gene dan its MTA1 protein product: role in epithelial cancer cell invasion, proliferation dan nuclear regulation. *Clin Exp Metastasis*, 20(1):19–24

Olumi A.F., Grossfeld G.D., Hayward S.W., Carroll P.R., Tlsty T.D., dan Cunha G.R. 1999. Carcinoma-associated fibroblasts direct tumor progression of initiated human prostatic epithelium. American Association for Cancer Researc. In: *Cancer Research* vol 59. P 5002.

Orimo A, Weinberg RA. 2006 Stromal fibroblasts in cancer: a novel tumor-promoting cell type. *Cell Cycle*. 2006 Aug;5(15):1597-601. Epub 2006 Aug 1.

Orimo. A, Gupta. P.B, Sgroi D.C., Seisdedos. F.A., Delaunay. T., Naeem .R, Carey. V.J., Richardson. A.L., Weinberg. R.A. 2005. Stromal fibroblasts present in invasive human breast carcinomas promote tumor growth dan angiogenesis through elevated SDF-1/CXCL12.. In: *Cell* Vol 121. Elsevier inc. P 335

Paget, S. 1889. The distribution of secondary growths in cancer of the breast. In; *The Lancet* vol 133 issue 3421. P 571–573.

Parr.C, Watkins.G, Mansel.RE, dan Jiang,WG . 2004. The Hepatocyte Growth Factor Regulatory Factors in HumanBreast Cancer. Clin Cancer Res;10:202-211. Published online January 20, 2004.

Pecorino, L. 2005. Molecular Biology of Cancer Mechanisms, Targets dan Therapeutic. Oxford University Press Inc. New York. P 157-172.

Pei D, Majmudar G, Weiss SJ. 1994. Hydrolytic inactivation of a breast carcinoma cell-derived serpin by human stromelysin-3. J Biol Chem, 269(41): 25849–25855.

Pelekanos.RA, Ting.MJ, Sardesai.VS, Ryan.JM, Lim.YC, Chan.JKY, and Fisk.NM, 2014. Intracellular trafficking and endocytosis of CXCR4 in fetal mesenchymal stem/stromal cells. *BMC Cell Biology* 2014, 15:15 http://www.biomedcentral.com/1471-2121/15/15

Penault-Llorca.F., André.F., Sagan. C., Lacroix-Triki. M., Denoux.Y., Verriele.V., Jacquemier.J., Baranzelli.M.C. 2009. Ki67 Expression and Docetaxel Efficacy in Patients With Estrogen Receptor–Positive Breast Cancer. Journal of Clinical Oncology.. Volume 27 Number 17 June 10 2009

Perou CM, Sorlie T, Eisen MB, van de, Rijn M, Jeffrey SS, Rees CA, Pollack JR, Ross DT, Johnsen H, Akslen LA, Fluge O, Pergamenschikov A, Williams C, Zhu SX,

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Lonning PE, Borresen-Dale AL, Brown PO, Botstein D. (2000). Molecular portraits of human breast tumours. Nature 406:747-752

Perou CM, Børresen-Dale AL. 2011. Systems biology and genomics of breast cancer. Cold Spring Harb Perspect Biol. 2011 Feb 1;3(2). pii: a003293. doi: 10.1101/cshperspect.a003293.

Petit I, Szyper-Kravitz M, Nagler A, Lahav M, Peled A, Habler L, Ponomaryov T, Taichman RS, Arenzana-Seisdedos F, Fujii N, Sandbank J, Zipori D, Lapidot T. 2002.G-CSF induces stem cell mobilization by decreasing bone marrow SDF-1 dan up-regulating CXCR4. In: Nature Immunology vol 3. P 687–694,

Phillips R.J., Burdick M.D., Lutz M, Belperio J.A., Keane M.P., dan Strieter R.M. 2003. The stromal derived factor-1/CXCL12-CXC chemokine receptor 4 biological axis in non-small cell lung cancer metastases. In: American Journal Respiratory Critical Care Medicine vol 167 no12. P 1676-1686.

Phillips. R.J, Mestas .J, Gharaee-Kermani. M, Burdick. M.D, Sica. A, Belperio. J.A, Keane. M.P, dan Strieter. R.M. 2005. Epidermal Growth Factor dan Hypoxia-induced Expression of CXC Chemokine Receptor 4 on Non-small Cell Lung Cancer Cells Is Regulated by the Phosphatidylinositol 3-Kinase/PTEN/AKT/ Mammalian Target of Rapamycin Signaling Pathway dan Activation of Hypoxia Inducible Factor-1. In: The Journal of Biological Chemistry Vol. 280 No. 23. P 22473–22481.

Platt-Higgins AM, Renshaw CA, West CR, Winstanley JH, De Silva Rudldan S, Barraclough R, Rudldan PS. 2000. Comparison of the metastasis-inducing protein S100A4 (p9ka) with other prognostic markers in human breast cancer. Int J Cancer, 89(2):198–208.

Poznansky MC, Olszak IT, Foxall R, Evans RH, Luster AD, Scadden DT. 2000. Active movement of T cells away from a chemokine. In: Nature Medicine vol 6. P543–548,

Racila E, Euhus D, Weiss AJ, Rao.C, McConenell. J. Terstappen. LW, Uhr. JW. 1998. Detection and characterization of carcinoma cells in the blood. *Proc Natl Acad Sc*i USA;95:4589–94

QIAGEN. 2009. Gene Globe Pathways- CXCR4 Pathway. https://www.qiagen.com/de/shop/genes-and-pathways/pathway-details/?pwid

Rahman. W. 2005. The Role of SDF1 alpha dan CXCR4 in Metastatic Breast Cancer. U.S. Army Medical Research dan Material Command.

Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein
Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan
Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Ratajczak. M.Z., Surma. E.Z., Kucia. M., Reca1. R., Wojakowski. W., dan Ratajczak. J . 2006. The pleiotropic effects of the SDF-1–CXCR4 axis in organogenesis, regeneration dan tumorigenesis. In*: Leukemia* vol 20. Nature Publishing Group. P 1915–1924

Ree AH, Pacheco MM, Tvermyr M, Fodstad O, Brentani MM. 2000. Expression of a novel factor, com1, in early tumor progression of breast cancer. Clin Cancer Res, 6(5):1778–1783.

Rozhin J, Sameni M, Ziegler G, Sloane BF. 1994. Pericellular pH affects distribution dan secretion of cathepsin B in malignant cells. *Cancer Res*, 54(24): 6517–6525.

Rubinstein.LV, Gail.MH, dan Santner.TJ. Planning the duration of a comparative clinical trial with loss to follow-up and a period of continued observation. *J Chron Dis*. 1981;34:469–479.

Rudldan PS, Platt-Higgins A, Renshaw C, West CR, Winstanley JH, Robertson L, Barraclough R. 2000. Prognostic significance of the metastasis-inducing protein S100A4 (p9Ka) in human breast cancer. Cancer Res, 60(6):1595–1603

Saadi W, Wang SJ, Lin F, Jeon NL. 2006. A parallel-gradient microfluidic chamber for quantitative analisis of breast cancer cell chemotaxis. Biomed Microdevices;8:109–18.

Schier AF. 2003. Chemokine signaling: rules of attraction. In: *Current Biology* vol 13. Elsevier Science Ltd. P 192–194.

Schmitt M, Janicke F, Moniwa N, Chucholowski N, Pache L, Graeff H. Tumor-associated urokinase-type plasminogen activator: biological dan clinical significance. Biol Chem Hoppe Seyler 1992, 373(7):611–622

Schmitt M, Wilhelm O, Janicke F, Magdolen V, Reuning U, Ohi H, Moniwa N, Kobayashi H, Weidle U, Graeff H. 1995. Urokinase-type plasminogen activator (uPA) dan its receptor (CD87): a new target in tumor invasion dan metastasis. *J Obstet Gynaecol*, 21(2):151–165Schneider IC, Haugh JM. 2006. Mechanisms of gradient sensing and chemotaxis: conserved pathways, diverse regulation. Cell Cycle;5:1130–34.

Schneider IC, Haugh JM. 2006. Mechanisms of gradient sensing and chemotaxis: conserved pathways, diverse regulation. *Cell Cycle*;5:1130–34.

Semov A, Moreno. M.J., Onichtchenko. A, Abulrob. A, Ball. M, Ekiel. I, Pietrzynski. G, Stanimirovic. D, dan Alakhov. V. 2005. Metastasis-associated protein S100A4

Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein
Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan
Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

induces angiogenesis through interaction with Annexin II dan accelerated plasmin formation. In: The Journal of Biological Chemistry vol 280. P 20833–20841.

Sehgal A, Keener C, Boynton AL, Warrick J, dan Murphy GP. 1998. CXCR-4, a chemokine receptor, is overexpressed in dan required for proliferation of glioblastoma tumor cells. In: *Journal Surgical Oncolology vol 69 no 2*. P 99-104.

Sheen-Chen SM, Liu YW, Eng HL, Chou FF. 2005. Serum levels of hepatocyte growth factor in patients with breast cancer. Cancer Epidemiol Biomarkers Prev, 14(3):715–717.

Singhal H, Bautista DS, Tonkin KS, O'Malley FP, Tuck AB, Chambers AF, Harris JF. 1997. Elevated plasma osteopontin in metastatic breast cancer associated with increased tumor burden dan decreased survival. Clin Cancer Res,3(4):605–611.

Sorlie T1, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, Johnsen H, Pesich R, Geisler S, Demeter J, Perou CM, Lønning PE, Brown PO, Børresen-Dale AL, Botstein D. 2003. Repeated observation of breast tumor subtypes in independent gene expression data sets. Proc Natl Acad Sci U S A. 2003 Jul 8;100(14):8418-23. Epub 2003 Jun 26.

Su. J.L., Yen. C.J., Chen. P.S., Chuang. S.E., Hong. C.C., Kuo. I.H., Chen. H.Y., Hung. M.C., Kuo. M.L. 2007. The role of the VEGF-C/ VEGFR-3 axis in Cancer Progression. In: *British Journal of Cancer* Vol 96. PP 541-545.

Su YC, Wu MT, Huang CJ, Hou MF, Yang SF, Chai CY. Expression of CXCR4 is associated with axillary lymph node status in patients with early breast cancer. *Breas*t. 2006;15:533–539

Sun YX, Schneider A, Jung Y, Wang J, Dai J, Cook K, Osman NI, Koh-Paige AJ, Shim H, Pienta KJ, Keller ET, McCauley LK, dan Taichman RS. 2005. Skeletal localization dan neutralization of the SDF- 1(CXCL12)/CXCR4 axis blocks prostate cancer metastasis dan growth in osseous sites in vivo. In: *Journal Bone Miner Research* vol 20. 318–329,

Sun. X., Wei. L., Chen. Q., dan Terek.R.M. 2010. CXCR4/SDF1 mediate hypoxia induced chondrosarcoma cell invasion through ERK signaling dan increased MMP1 expression. In: Molecular Cancer vol 9. P 17

Tachibana K, Hirota S, Iizasa H, Yoshida H, Kawabata K, Kataoka Y, Kitamura Y, Matsushima K, Yoshida N, Nishikawa S, Kishimoto T, Nagasawa T. 1998. The chemokine receptor CXCR4 is essential for vascularization of the gastrointestinal tract. In: Nature. Vol 393 no 6685. P 524-525.

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Talvensaari-Mattila A, Paakko P, Hoyhtya M, Blanco-Sequeiros G, Turpeenniemi-Hujanen T. 1998. Matrix metalloproteinase-2 immunoreactive protein: a marker of aggressiveness in breast carcinoma. Cancer, 83(6):1153– 1162.

Tashiro K, Tada H, Heilker R, Shirozu M, Nakano T, Honjo T. 1993. Signal sequence trap: a cloning strategy for secreted proteins dan type I membrane proteins. In: Science vol 261. P 600–603.

Thomssen C, Schmitt M, Goretzki L, Oppelt P, Pache L, Dettmar P, Janicke F, Graeff H. 1995. Prognostic value of the cysteine proteases cathepsins B dan cathepsin L in human breast cancer. *Clin Cancer Res*, 1(7):741–746.

Tjindarbumi, D. dan Mangukusumo, R. 2002. Cancer in Indonesia Present dan Future. In: Japan Journal Clinical Oncology vol 32 (Supplement). P 17-21

Tong RT, Boucher Y, Kozin SV, Winkler F, Hicklin DJ, Jain RK. 2004. Vascular normalization by vascular endothelial growth factor receptor 2 blockade induces a pressure gradient across the vasculature and improves drug penetration in tumors. Cancer Res; 64:3731–36.

Tuck AB, Chambers AF. 2001. The role of osteopontin in breast cancer: clinical dan experimental studies. J Mammary Gldan Biol Neoplasia, 6(4):419–429.

Tuck AB, Elliott BE, Hota C, Tremblay E, Chambers AF. 2000. Osteopontin induced, integrin-dependent migration of human mammary epithelial cells involves activation of the hepatocyte growth factor receptor (Met). J Cell Biochem, 78(3):465–475.

Tuck AB, Hota C, Wilson SM, Chambers AF. 2003. Osteopontin-induced migration of human mammary epithelial cells involves activation of EGF receptor dan multiple signal transduction pathways. Oncogene, 22(8):1198–1205.

Tuck AB, O'Malley FP, Singhal H, Harris JF, Tonkin KS, Kerkvliet N, Saad Z, Doig GS, Chambers AF. 1998. Osteopontin expression in a group of lymph node negative breast cancer patients. Int J Cancer, 79(5):502–508.

Vaupel. P., dan Mayer. A. 2007. Hypoxia in Cancer: Significance dan Impact on Clinical Outcome. In: Cancer Metastasis Reviews Vol 26, No 2. P 225–239.

Vogel .V.G. 2004. Epidemiology of Breast Cancer. In: The Breast Comprehensive Management of Benign dan Malignant Disorder 3rd Edition. Saunders Company. St Louis. P 341-354

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Wang M, Liu YE, Greene J, Sheng S, Fuchs A, Rosen EM, Shi YE. 1997. Inhibition of tumor growth dan metastasis of human breast cancer cells transfected with tissue inhibitor of metalloproteinase 4. Oncogene, 14(23):2767–2774.

Wang SJ, Saadi W, Lin F, Minh-Canh Nguyen C, Li Jeon N. 2004. Differential effects of EGF gradient profiles on MDA-MB-231 breast cancer cell chemotaxis. Exp Cell Res;300:180–89.

Watkins G, Douglas-Jones A, Bryce R, Mansel RE, Jiang WG. 2005. Increased levels of SPARC (osteonectin) in human breast cancer tissues dan its association with clinical outcomes. Prostaglandins Leukot Essent Fatty Acids, 72(4): 267–272.

World Health Organization (WHO). 2014. Cancer Country Profile – World Health Orgazitation. http://www.who.int/cancer/country-profiles/idn_en.pdf

Wolff. AC , Hammond. ME , Hicks. DG , Dowsett.M , McShane.LM , Allison. KH, Allred.DC , Bartlett.JM , Bilous. M, Fitzgibbons.P , Hanna. W, Jenkins. RB, Mangu.PB , Paik.S, Perez.EA, Press.MF, Spears.PA, Vance.GH, Viale.G, dan Hayes.DF. 2013. Recommendations for human epidermal growth factor receptor 2 testing in breast cancer: American Society of Clinical Oncology/College of American Pathologists clinical practice guideline updateJournal. Journal of Clinical Oncology. 31(31):3997-4013

Woo SU, Bae JW, Kim CH, Lee JB, Koo BW. A significant correlation between nuclear CXCR4 expression and axillary lymph node metastasis in hormonal receptor negative breast cancer. Ann Surg Oncol. 2008 Jan;15(1):281-5. Epub 2007 Sep 1.

Yarso. K.Y dan Sudarsa. I.W. Tripel Negative as basal type of breas cancer in Balinese Women. Dipresentasikan Pada Mukatamar Mabi XV 2010 surabaya

Yarso. K.Y, Sudarsa. I.W, dan Manuaba. 2012. Intitial respon chemotherapy in triple negative Breast Cancer. Bali Medical Journal. Vol 1. No 1 2012.

Yarso. K.Y., Suyatmi, Syah.FK. Manuaba. IBTW, Hakimi.M, Astuti.I, dan Aryandono. T. 2016. Association of SDF-1 with Metastasis in Breast Cancer Patient at Sanglah Hospital Bali. *Bali Medical Journal*.. Vol 6 no 1 2017.

Yamashita J, Akizuki M, Jotsuka T, Harao M, Nakano S. 2006. Neutrophil elastase predicts trastuzumab responsiveness in metastatic breast cancer. Breast J, 12(3):288.

UNIVERSITAS GADJAH MADA
Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Yamashita J, Ogawa M, Shirakusa T. 1995. Free-form neutrophil elastase is an independent marker predicting recurrence in primary breast cancer. J Leukoc Biol, 57(3):375–378.

Yang SY, Lee J, Park CG, Kim S, Hong S, Chung HC, Min SK, Han JW, Lee HW, Lee HY. 2002. Expression of autotaxin (NPP-2) is closely linked to invasiveness of breast cancer cells. Clin Exp Metastasis, 19(7):603–608

Yasuoka H, Tsujimoto M, Yoshidome K, Nakahara M, Kodama R, Sanke T, Nakamura Y. Cytoplasmic CXCR4 expression in breast cancer: induction by nitric oxide and correlation with lymph node metastasis and poor prognosis. BMC Cancer. 2008 Nov 23;8:340. doi: 10.1186/1471-2407-8-340.

Yoneda T, Sasaki A, Dunstan C, Williams PJ, Bauss F, De Clerck YA, Mundy GR. 1997. Inhibition of osteolytic bone metastasis of breast cancer by combined treatment with the bisphosphonate ibandronate dan tissue inhibitor of the matrix metalloproteinase-2. J Clinical Invest, 99(10):2509–2517.

Young.K.C., Huang.C.T., Lin.S.L., dan Hsiao.W.C., 2006. Two-Phased Alteration dan Clinical Relevance of MTA1dan MTA1s Gene Expression: A Quantitative PCR Analisis in Breast Cancer Patients. *Journal of Cancer Molecules* 2(2): 79-84

Yui S, Tomita K, Kudo T, Ando S, Yamazaki M. 2005. Induction of multicellular 3-D spheroids of MCF-7 breast carcinoma cells by neutrophil-derived cathepsin G dan elastase. Cancer Science, 96(9):560–570.

Zager. J, Solorzano. C, Thomas. E, Feig, B, dan Babiera. G. 2006. Invasive Breast Cancer. In: MD Anderson Surgical Oncology Handbook, The, 4th Edition. Lippincott Williams dan Wilkins. P

Zeelenberg IS, Ruuls-Van Stalle L, Roos E. 2003. The chemokine receptor CXCR4 is required for outgrowth of colon carcinoma micrometastases. In: Cancer Research vol 63. P 3833–3839

Zhang H, Stephens LC, Kumar R. 2006. Metastasis tumor antigen family proteins during breast cancer progression dan metastasis in a reliable mouse model for human breast cancer. Clin Cancer Res, 12(5):1479–1486

Zhang.Z., Ni.C., Chen.W., Wu.P., Wang.Z., Yin.J., Huang.J. and Qiu.F. 2014. Expression of CXCR4 and breast cancer prognosis: a systematic review and meta-analysis†Contributed equally. BMC Cancer201414:49 DOI: 10.1186/1471-2407-14-49

Pengaruh Ekspresi mRNA Kemokin Ligan SDF-1 pada Jaringan Kanker Payudara dan Ekspresi Protein
Kemokin Reseptor CXCR4 pada Inti Sel Kanker Payudara Terhadap Resiko Metastasis Jauh dan Ketahanan
Hidup
Kristanto Yuli Yarso, Prof. Dr. dr. Teguh Aryandono Sp.BOnk(K).; Prof. dr. H. M Hakimi. PhD. SpOG (K),; DR. Med.d
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Zlotnik A. 2006. Involvement of chemokine receptors in organ-specific metastasis. Contrib Microbiol, 13:191–199.

Zou L, Barnett B, Safah H, Larussa VF, Evdemon-Hogan M, Mottram P, Wei S, David O, Curiel TJ, Zou W. 2004. Bone marrow is a reservoir for CD4_CD25_ regulatory T cells that traffic through CXCL12/CXCR4 signals. In: Cancer Research vol 64. P 8451–8455

Zou YR, Kottmann AH, Kuroda M, Taniuchi I, Littman DR. 1998. Function of the chemokine receptor CXCR4 in haematopoiesis dan in cerebellar development. In: Nature vol 393. P 595–599