

DAFTAR PUSTAKA

- Abdallah, S., Alaa, E., & Naseem, M. (2010). Obesity and eating habits among college students in Saudi Arabia: A cross sectional study. *Nutr. J.*, 9, 3-4.
- Almatsier, Sunita. (2003). *Prinsip Dasar Ilmu Gizi*. Jakarta : PT Gramedia Pustaka Utama, 77-78.
- Arnau, A., Espauellad, J., Serrarolse, M., Canudase, J., Formigaf, F., & Ferrerb, M. (2016). Risk factors for functional decline in a population aged 75 years and older without total dependence: A one-year follow-up. *Gerontology and Geriatrics*, 65, 239–247.
- Badan Pusat Statistik. (2015). *Statistik Penduduk Lanjut Usia 2014 Hasil Survey Sosial Ekonomi Nasional*. Jakarta: Badan Pusat Statistika, 11-15.
- Beriault, K., Carpentier, A.C., Gagnon, C., Ménard, J., Baillargeon, J.P., Ardilouze, J.L., & Langlois, M.F. (2009). Reproducibility of the 6-minute walk test in obese adults. *Int J Sports Med*, 30(10), 725-727.
- Bonnefoy, M., Cornu, C., Normand, S., Boutitie, F., Bugnard, F., & Rahmani, A. (2003). The effect of exercise and protein energy supplements on body composition and muscle function in frail elderly individual: a long term controlled randomized study. *BJN*, 89, 731-738.
- Borges, L.S., Fernandes, M.H., Schettino, L., Coqueiro, R.D.S., & Pereira, R. (2015). Handgrip explosive force is correlated with mobility in the elderly women. *Acta of Bioengineering and Biomechanics*, 17(3), 134-141.
- Brach, J.S., & VanSwearingen, J.M. (2002). Physical impairment and disability : Relationship to performance of activities of daily living in communitydwelling older men. *Journal of Physical Therapy*, 82(8), 753.
- Brown, J.E. (2011). *Nutrition trough the life cycle (Fourth Edition)*. USA: Wardsworth Cengage Learning, 125-128.
- Camry Bodyfit EF973 *manual book*, 6.
- Casanova, C., Celli, B.R., Barria, P., Casas, A., Cote, C., de Torre, J.P., Jardim, J., Lopez, M.V., Marin, J.M., Montes de Oca, M. Pinto-Plata, V., & Aguirre-Jaime, A. (2001). The 6-min walk distance in healthy subjects: reference standards from seven countries. *Eur Respir J*, 37, 150–156.
- Cereda, M., Pedrolli, C., Zagami, A., Vanotti, A., Piffer, S., Faliva, M., Rondanelli, M., & Caccialanza, R. (2013). Nutritional risk, functional status and mortality in newly institutionalised elderly. *British Journal of Nutrition*, 110, 1903–1909.

- Chatchalit, R. and Rattana, L. (2004). Gender differences of regional abdominal fat distribution and their relationships with insulin sensitivity in healthy and glucose-intolerant. *Thai. J. Clin. Endocrinol.Metabol.*, 89, 6266-6270.
- Chevalier, S., Saoud, F., Gray-Donald, K., &Morais, J.A. (2008). The physical functional capacity of frail elderly persons undergoing ambulatory rehabilitation is related to their nutritional status. *J Nutr Health Aging.*, 12(10), 721-726.
- Chodzko-Zajko, W.J., Proctor, D.N., Singh, M.A.F., Minson, C.T., Nigg, C.R., & Salem, G.J. (2009). Exercise and physical activity for older adults. *Med Sci Sports Exerc*, 41(7), 1510-1530
- Craig, C.L., Marshall, M. & Sjostrom, (2003). International physical activity questionnaire: 12-country reliability and validity. *Medicine and science in sports and exercise*, 35(8), 1381–95.
- Crim, M.T., Yoon, S.S., Ortiz, E., Wall, H.K., Schober, S., Gillesie, C., Sorlie, P., Keenan, N., Labarte, D., Hong, Y. (2012). National urveillance definitions for hypertension prevalence and control among adult. *Circ Cardiovasc Qual Outcome*, 5(3), 343–351.
- Darmojo, B. (2015). *Buku ajar geriatri (Edisi 5)*. Jakarta: Badan Penerbit FKUI, 8 – 17.
- De Faria Santarém, G.C., de Cleve, R., Santo, M.A., Bernhard, A.B., Gadducci, A.V., Greve, J.M.D., & Silva, P.R.S. (2015). Correlation between body composition and walking capacity in severe obesity. *PLoS ONE*, 10(6), 1-10.
- De Souza, S.A., Faintuch, J., Fabris, S.M., Nampo, F.K., Luz, C., Fabio, T.L., Sitta, I.S., & de Batista Fonseca, I.C. (2009). Six-minute walk test: functional capacity of severely obese before and after bariatric surgery. *Surg Obes Relat Dis*, 5(5), 540-543.
- Dias, F.M., Costa, S.O., Freitas, J.P., Pinto, A.C.R., Vigário, P.S., & Mainenti, M.R.M. (2014). Functional capacity of oldest old living in a long-stay institution in Rio De Janeiro Brazil. *J Phys Ther Sci*, 26, 1097–1105.
- Diehr, P., Thielke, S., O'Meara, E., Fitzpatrick, A.L., & Newman, A. (2012). Comparing years of healthy life, Measured in 16 ways, for normal weight and overweight older adults. *Journal of Obesity*, 1-13.
- Dinas Kesehatan Kota Yogyakarta. (2015). *Profil Kesehatan Tahun 2015 Kota Yogyakarta*. Yogyakarta : Dinas Kesehatan Kota Yogyakarta, 34-35; 80.
- Dong, H.J., Marcusson, J., Wressle, E., & Unosson, M. (2015). Obese very old women have low relative handgrip strength, poor physical function, and difficulties in daily living. *J Nutr Health Aging*, 19(1), 20-25.

Enright, P.L., McBurnie, M.A., Bittner, V., Tracy, R.P., McNamara, R., Arnold, A., & Newman, A.B. (2003). The 6 min walk test: A quick measure of functional status in elderly adults. *Chest Clinical Journal*, 123, 387-397.

Esmayel, E.M.M., Eldarawy, M.M., Hassan, M.M., Hassanin, H.M., Ashour, W.M.R., & Mahmoud, W. (2013). Nutritional and functional assessment of hospitalized elderly : Impact of sociodemographic variables. *Journal of Aging Research*, 1 – 7.

Eynon, N., Yamin, C., Ben-Sira, D., & Sagiv, M. (2009). Optimal health and function among the elderly: lessening severity of ADL disability. *Eur Rev Aging Phys Act*, 6, 55–61.

Falsarella, G.R., Coimbra, I.B., Barcelos, C., Iartelli, I., Montedori, K.T., Santos, M., Neri, A.L., & Coimbra, A.M.V. (2014). Influence of muscle mass and bone mass on the mobility of elderly women: an observational study. *BMC Geriatrics*, 14(13), 1-7.

Fatmah. (2010). *Gizi usia lanjut*. Jakarta: Penerbit Erlangga, 11-12, 36, 46-47.

Ferruci, L., Gularnik, J.M., Buchner, D. (1997). The relationship between measures of muscular strength and physical performance of the lower extremities : The woman's health ang aging study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 52, 12-15.

Gallagher, D., Visser, M., De Meersman, R.E., Sepulveda, D., Baumgartner, R.N., Pierson, R.N., Harris, T., Heymsfield, S.B. (1996). Appendicular skeletal muscle mass: effects of age, gender, and ethnicity. *J Appl Physiol*, 83, 229–239.

Gibbons, W.J. (2001). Reference values for a multiple repetition 6-minute walk test in healthy adults older than 20 years. *Journal of Cardiopulmonary Rehabilitation*, 21(2), 87-93.

Gibson, R.S. (2005). *Principles of Nutritional Assessment*. New York : Oxford University Press. 290-291.

Gomes-Neto, M., Araujo, A.D., Junqueira, I. D.A., Oliveira, D., Brasileiro, A., & Arcanjo, F.L. (2016). Comparative study of functional capacity and quality of life among obese and non-obese elderly people with knee osteoarthritis. *Revista Brasileira de Reumatologia*, 56(2), 126–130.

Goya, W., Geral, S., & Richard, M. (2005). Measures of adiposity in the identification of metabolic abnormalities in elderly men. *Am. J. Clin. Nutr.*, 81, 1313-1321.

Guralnik, J.M., Ferruci, L., & Pipier, C.F. (2000). Lower extremity function and subsequent disability : consistency across studies, predictive models,

and value gait speed alone compared with the short physical performance battery. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 55, 221-231.

Hardywinoto, Setiabudhi T. (2005). *Panduan Gerontologi Tinjauan Dari Berbagai Aspek*. Jakarta : PT Ikrar Mandiri Abadi, 90-92.

Helena, S., Elena, Z., & Paolo, T. (2001). Relation between body composition, fat distribution, and lung function in elderly men. *Am. J. Clin. Nutr.*, 73, 827-831.

Henningsson, E. and Ekelund, U. (2007). Is the association between physical activity and body mass index obesity dependent?. *Int J Obes*, 31: 663-668.

Hulens, M., Vansant, G., Lysens, R., Claessens, A.L., Muls, E., & Brumagne, S. (2001). Study of differences in peripheral muscle strength of lean versus obese women: an allometric approach. *Int J Obes Relat Metab Disord*, 25, 676-68.

IPAQ group, (2002). *International Physical Activity Questionnaires*. <http://www.ipaq.ki.se/questionnaires/IPAQ>, 15-21.

Jankowski, C.M., Gozansky, W.S., Van Pelt, R.E., Schenkman, M.L., Wolfe, P., Schwartz, R.S., & Kohrt, W.M. (2008). Relative contributions of adiposity and muscularity to physical function in community-dwelling older adults. *Obesity*, 16, 1039-1044.

Jamar hydraulic hand dynamometer. (2003). *Manual book*, 3-4.

Jette, A., & Haley, S. (2002). Late life function and disability instrument : development and evaluation of the disability component. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 57, 209 – 216.

Jorgic, B., Pantelic, S., Milanovic, Z., & Kostic, R. (2011). The effect of physical exercise on the body composition of the elderly: a systematic review. *J Phys Educ Sport*, 9(4), 439 – 453.

Kamarul, T., Ahmad T.S., & Loh, W.Y.C. (2006). Hand grip strenght in the adult Malaysian population. *Journal of Orthopaedic Surgery*, 14 (2), 172 – 177.

Kamper, A.M., Stott, D.J., Hyland, M., Murray, H.M., & Ford, I. (2005). Predictors of functional decline in elderly people with vascular risk factors or disease. *Age and Ageing*, 34, 450-455.

Kementrian Kesehatan RI. (2013). *Riset Kesehatan Dasar Tahun 2013*. Jakarta : Badan Penelitian dan Pengembangan Kesehatan Kementrian Kesehatan RI, 88-90.

Kementrian Kesehatan RI. (2014). *Hipertensi*. Jakarta : Pusat Data dan Informasi
Kementrian Kesehatan RI, 2-3.

King-sheen *Electronic Hand Dynamometer Instructions for Use*, 8.

Kozicka, I., Kostka, T. (2016). Handgrip strength, quadriceps muscle power, and optimal shortening velocity roles in maintaining functional abilities in older adults living in a longterm care home: a 1-year follow-up study. *Clinical Interventions in Aging*, 11, 739–747.

Lauretani, F., Russo, C.R., & Bandinelli, S. (2003). Age-associated changes in sceletal muscle and their effect on mobility : an operational diagnosis of sarcopenia. *Journal of Applied Physiologi*, 95, 1851-1860.

Lee, R.D., and Nieman, D.C. (2013). *Nutritional Assessment (Sixth Edition)*. New York : Mc Graw Hill, 185 – 186.

Lemeshow, S., David, W., Janelle, K., Stephen, K. (1997). *Adequacy of Sample Size in Health Studies*. Penerjemah : Pramono, Dibyo. *Besar Sample dalam Penelitian Kesehatan*. Yogyakarta : Gama Press, 21-22.

Maria, A. (2004). Exercise and aging. *Clin. Geriatr.Med.*, 20, 201-221.

Milanović, Z., Pantelić, S., Trajković, N., & Sporiš, G. (2011). Basic anthropometric and body composition characteristics in a elderly population: a systematic review. *Facta Universitatis Series: Physical Education and Sport*, 9(2), 173 – 182.

Mohanty, S.P., Babu, S.S., & Nair, N.S. (2001). The use of arm span as a predictor of height: a study of south Indian women. *Journal of Orthopaedic Surgery*, 9, 19-23.

Nadeak, D.C. (2004). *Diagnosis malnutrisi pada usia lanjut*. Majalah Gizi Medik Indonesia, 3, 14-16.

Ng Tze Pin.(2009). Chronic disease, functional status and quality of life among the elderly in Singapore. *Symposium and Healthcare Challenges for a Ageing Population : Managing Healthcare and End of Life Decisions in Singapore*, 2-4.

Norman, K., Stobaus, N., & Gonzales M.C. (2011). Handgrip strength: outcome predictor and marker of nutritional status. *Clinical Nutrition*, 30, 135-142.

Payette, H., Hanusaik, N., Boutier, V., Morais, J.A., Gray-Donald, K. (1998). Muscle strength and functional mobility in relation to lean body mass in free-living frail elderly women. *Eur J Clin Nutr*, 52, 45–53.

Peolsson, A., Hedlund, H., & Oerberg, B. (2001). Intra and inter tester reliability and reference values of hand strength. *J Rehab Med*, 33, 36-41.

- Perls, T., & Fretts, R. (1998). *Why women live longer than men*. <http://www.sciam.com/1998/0698womens/0698perls.html>, 1-8.
- Pietrerse, S., Manandhar, M., & Ismail, S. (2002). The association between nutritional status and handgrip strength in older Rwanda refugees. *European Journal of Clinical Nutrition*, 56, 933-939.
- Poh, H., Eastwood, P.R, Cecins, N.M., Ho, K.T., & Jenkins, S.C. (2006). Six-minute walk distance in healthy Singaporean adults cannot be predicted using reference equations derived from Caucasian populations. *Respirology*, 11(2), 211-216.
- Ramos, R.A., & Ferreira, A.S. (2014). Functional capacity in adults with hypertension as assessed by the six-minute walk distance test: systematic review. *Fisioter Pesq.*, 21(3), 257-263.
- Rantanen, T., Guralnik, J.M., Foley, D., Masaki, K., Leveille, S., & Curb, J.D. (1999). Midlife handgrip strength as a predictor of old age disability. *JAMA*, 281, 558-560.
- Sedenkova, B., Stejska, P., Simicek, J., Elfmark, M., Businova, T., & Ranikova, B. (2012). The influence of a six month aerobic programme on middle aged woman's aerobic capacity and body composition. *Acta Univ Palacki Olomuc Gymn*, 42(3), 55-66.
- Setia, A. (2008). *Faktor-faktor yang berhubungan dengan tingkat kebugaran lanjut usia penghuni Panti Budi Agung Kupang*. Yogyakarta: UGM, 35-36.
- Setiati, S. (2013). Geriatric medicine, sarkopenia, frailty dan kualitas hidup pasien usia lanjut: tantangan masa depan pendidikan, penelitian dan pelayanan kedokteran di Indonesia. *eJKI*, 1(3), 234-242.
- Shechtman, O., Mann, W.C., Justiss, M.D., & Tomita, M. (2004). Grip strength in the frail elderly. *Am J Phys Med Rehabil*, 83, 819-826.
- Silva, R., Giribela, C., Wolosker, N., & Consolim-Colombo, M. (2012). Functional limitation and intermittent claudication: Impact of blood pressure measurements. *Arq Bras Cardiol*, 98(2), 161-166.
- Steffen, T.M., Hacker, T.A., & Mollinger, L. (2002). Age and gender related test performance in community dwelling elderly people : Six Minutes Walk Test, Berg Balance, Timed Up and Go Test , and Gait Speeds. *Journal Physical Therapy*, 82, 128-137.
- Sugiarto, Andi. (2005). *Penilaian keseimbangan dengan aktivitas kehidupan sehari-hari pada lansia di panti werdha Pelkris Elim Semarang dengan menggunakan Berg Balance Scale dan Indeks Barthel*. Semarang : UNDIP, 7 – 8.

Supariasa, I.D.N., Bachyar Bakri, Ibnu Fajar. (2002). *Penilaian Status Gizi*. Jakarta: Penerbit Buku Kedokteran EGC, 18; 94-96; 114.

Torres, G.V., & Reis, L.A. (2010). Assessment of functional capacity in elderly residents of an outlying area in the hinterland of Bahia/Northeast Brazil. *Arq Neuropsiquiatr*, 68(1), 39-43.

Tower, J., and Arbeitman, M., (2009). The genetics of gender and life span. *Journal of Biology*, 8(38), 1 -3.

Trihandini, Indang. (2007). Peran medical check up terhadap aktivitas fisik dasar lansia : Studi panel kelompok lanjut usia 1993-2000. *Jurnal Makara Kesehatan*, 2(2), 90-91.

Utomo, B dan Surini, S. (2003). *Fisioterapi Pada Lansia*. Jakarta : Penerbit Buku Kedokteran EGC, 11-12.

Vahlberg, B., Zetterberg, L., Lindmark, B., Hellström, K., & Cederholm, T. (2016). Functional performance, nutritional status, and body composition in ambulant community-dwelling individuals 1–3 years after suffering from a cerebral infarction or intracerebral bleeding. *BMC Geriatrics*, 16(48), 1-9.

Wanderley, F.A.C. (2011). Six-Minute Walk Distance (6MWD) is associated with body fat, systolic blood pressure, and rate-pressure product in community dwelling elderly subjects. *Archives of Gerontology and Geriatrics*, 52, 206–210.

WHO. (2000). *The Asia-Pacific perspective : Redefining obesity and its treatment*. Health Communications Australia, 17-18.

Wiraharja, R.S. (2014). *Sindroma Geriatri pada Lansia di Komunitas*. Jakarta: Universitas Atma Jaya, 25-41.

Zoico, E., Di Francesco, V., Guralnik, J.M., Mazzali, G., Bortolani, A., Guarient, S., Sergi, G., Bosello, O., & Zamboni, M. (2004). Physical disability and muscular strength in relation to obesity and different body composition indexes in a sample of healthy elderly women. *International Journal of Obesity*, 28, 234–241.