

## DAFTAR PUSTAKA

- [1] R. V. Krishnamurthi *et al.*, “Stroke prevalence, mortality and disability-adjusted life years in adults aged 20-64 years in 1990-2013: Data from the global burden of disease 2013 study,” *Neuroepidemiology*, vol. 45, no. 3, hal. 190–202, 2015.
- [2] B. H. Dobkin, “Rehabilitation after stroke,” *N. Engl. J. Med.*, vol. 352, no. 16, hal. 1677–1684, 2005.
- [3] Centers for Disease Control and Prevention (CDC), “Outpatient rehabilitation among stroke survivors--21 States and the District of Columbia, 2005,” *MMWR Morb. Mortal. Wkly. Rep.*, vol. 56, no. 20, hal. 504–507, Mei 2007.
- [4] D. J. Gladstone, C. J. Danells, dan S. E. Black, “The Fugl-Meyer Assessment of Motor Recovery after Stroke: A Critical Review of Its Measurement Properties,” *Neurorehabil. Neural Repair*, vol. 16, no. 3, hal. 232–240, Sep 2002.
- [5] A. R. Fugl-Meyer, L. Jääskö, I. Leyman, S. Olsson, dan S. Steglind, “The post-stroke hemiplegic patient. 1. a method for evaluation of physical performance.,” *Scand. J. Rehabil. Med.*, vol. 7, no. 1, hal. 13–31, 1974.
- [6] C. Collin dan D. Wade, “Assessing motor impairment after stroke: a pilot reliability study.,” *J. Neurol. Neurosurg. Psychiatry*, vol. 53, no. 7, hal. 576–579, Jul 1990.
- [7] Medical Research Council, *Aids to the examination of the peripheral nervous system*. HM Stationery Office, 1976.
- [8] S. L. Wolf, P. A. Catlin, M. Ellis, A. L. Archer, B. Morgan, dan A. Piacentino, “Assessing Wolf motor function test as outcome measure for research in patients after stroke,” *Stroke*, vol. 32, no. 7, hal. 1635–1639, 2001.
- [9] M. Wintz, “Variations in current manual muscle testing.,” *Phys. Ther. Rev.*, vol. 39, no. 7, hal. 466, 1959.
- [10] P. W. Duncan, M. Propst, dan S. G. Nelson, “Reliability of the Fugl-Meyer assessment of sensorimotor recovery following cerebrovascular accident,” *Phys. Ther.*, vol. 63, no. 10, hal. 1606–1610, 1983.
- [11] A. R. Fugl-Meyer dan L. Jääskö, “Post-stroke hemiplegia and ADL-performance.,” *Scand. J. Rehabil. Med. Suppl.*, vol. 7, hal. 140, 1980.
- [12] S. L. Wood-Dauphinee, J. I. Williams, dan S. H. Shapiro, “Examining outcome measures in a clinical study of stroke.,” *Stroke*, vol. 21, no. 5, hal. 731–739, 1990.
- [13] E. Taub *et al.*, “Technique to improve chronic motor deficit after stroke.,” *Arch. Phys. Med. Rehabil.*, vol. 74, no. 4, hal. 347–354, 1993.
- [14] F. Cary, O. Postolache, dan P. S. Girão, “Kinect based system and Artificial Neural Networks classifiers for physiotherapy assessment,” in *2014 IEEE International Symposium on Medical Measurements and Applications (MeMeA)*, 2014, hal. 1–6.

- [15] S. D. Din, S. Patel, C. Cobelli, dan P. Bonato, "Estimating fugl-meyer clinical scores in stroke survivors using wearable sensors," in *2011 Annual International Conference of the IEEE Engineering in Medicine and Biology Society*, 2011, hal. 5839–5842.
- [16] W.-S. Kim, S. Cho, D. Baek, H. Bang, dan N.-J. Paik, "Upper Extremity Functional Evaluation by Fugl-Meyer Assessment Scoring Using Depth-Sensing Camera in Hemiplegic Stroke Patients," *PLOS ONE*, vol. 11, no. 7, hal. e0158640, Jul 2016.
- [17] S. H. Lee, M. Song, dan J. Kim, "Towards clinically relevant automatic assessment of upper-limb motor function impairment," in *2016 IEEE-EMBS International Conference on Biomedical and Health Informatics (BHI)*, 2016, hal. 148–151.
- [18] E. V. Olesh, S. Yakovenko, dan V. Gritsenko, "Automated Assessment of Upper Extremity Movement Impairment due to Stroke," *PLOS ONE*, vol. 9, no. 8, hal. e104487, Agu 2014.
- [19] P. Otten, S. H. Son, dan J. Kim, "Automating Stroke Patient Evaluation Using Sensor Data and SVM," in *2014 IEEE 7th International Conference on Service-Oriented Computing and Applications*, 2014, hal. 223–229.
- [20] O. Patsadu, C. Nukoolkit, dan B. Watanapa, "Human gesture recognition using Kinect camera," in *2012 International Joint Conference on Computer Science and Software Engineering (JCSSE)*, 2012, hal. 28–32.
- [21] A. Scano, M. Caimmi, A. Chiavenna, M. Malosio, dan L. M. Tosatti, "Kinect One-based biomechanical assessment of upper-limb performance compared to clinical scales in post-stroke patients," in *2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*, 2015, hal. 5720–5723.
- [22] A. Scano, M. Caimmi, M. Malosio, dan L. M. Tosatti, "Using Kinect for upper-limb functional evaluation in home rehabilitation: A comparison with a 3D stereoscopic passive marker system," in *5th IEEE RAS/EMBS International Conference on Biomedical Robotics and Biomechatronics*, 2014, hal. 561–566.
- [23] J. Wang, L. Yu, J. Wang, L. Guo, X. Gu, dan Q. Fang, "Automated Fugl-Meyer Assessment using SVR model," in *2014 IEEE International Symposium on Bioelectronics and Bioinformatics (ISBB)*, 2014, hal. 1–4.
- [24] L. Yu, D. Xiong, L. Guo, dan J. Wang, "A remote quantitative Fugl-Meyer assessment framework for stroke patients based on wearable sensor networks," *Comput. Methods Programs Biomed.*, vol. 128, hal. 100–110.
- [25] J. A. Chambers *et al.*, "Adherence to medication in stroke survivors: a qualitative comparison of low and high adherers," *Br. J. Health Psychol.*, vol. 16, no. 3, hal. 592–609, 2011.
- [26] P. Otten, J. Kim, dan S. H. Son, "A Framework to Automate Assessment of Upper-Limb Motor Function Impairment: A Feasibility Study," *Sensors*, vol. 15, no. 8, hal. 20097–20114, Agu 2015.
- [27] Y.-W. Hsieh, I.-P. Hsueh, Y.-T. Chou, C.-F. Sheu, C.-L. Hsieh, dan G. Kwakkel, "Development and Validation of a Short Form of the Fugl-Meyer

- Motor Scale in Patients With Stroke,” *Stroke*, vol. 38, no. 11, hal. 3052–3054, Nov 2007.
- [28] P. Otten, S. H. Son, dan J. Kim, “Automating Stroke Patient Evaluation Using Sensor Data and SVM,” in *2014 IEEE 7th International Conference on Service-Oriented Computing and Applications*, 2014, hal. 223–229.
- [29] D. S. Nichols-Larsen, P. Clark, A. Zeringue, A. Greenspan, dan S. Blanton, “Factors influencing stroke survivors’ quality of life during subacute recovery,” *Stroke*, vol. 36, no. 7, hal. 1480–1484, 2005.
- [30] A. Kedavra, “Official Kinect SDK vs. Open-source alternatives - Stack Overflow,” 09-Okt-2011. [Daring]. Tersedia pada: <http://stackoverflow.com/questions/7706448/official-kinect-sdk-vs-open-source-alternatives>. [Diakses: 23-Apr-2016].
- [31] M. K4W Team, “Kinect API Overview.” [Daring]. Tersedia pada: <https://msdn.microsoft.com/en-us/library/dn782033.aspx>. [Diakses: 24-Apr-2016].
- [32] R. Julianjatsono, R. Ferdiana, dan R. Hartanto, “PENGEMBANGAN DAN PENGUJIAN APLIKASI PERMAINAN OLAHRAGA RITMIS (EXERGAME) MENGGUNAKAN SENSOR MICROSOFT KINECT,” *JNTETI*, 2016.
- [33] H.-L. Teulings, J. L. Contreras-Vidal, G. E. Stelmach, dan C. H. Adler, “Parkinsonism reduces coordination of fingers, wrist, and arm in fine motor control,” *Exp. Neurol.*, vol. 146, no. 1, hal. 159–170, 1997.
- [34] B. Jeff, *Azure Machine Learning Microsoft Azure Essentials*. Redmond, Washington 98052-6399: Microsoft Press.
- [35] J. O. Rawlings, S. G. Pantula, dan D. A. Dickey, *Applied regression analysis: a research tool*. Springer Science & Business Media, 2001.
- [36] C. M. Bishop dan M. E. Tipping, “Bayesian regression and classification.”
- [37] J. N. Morgan dan J. A. Sonquist, “Problems in the analysis of survey data, and a proposal,” *J. Am. Stat. Assoc.*, vol. 58, no. 302, hal. 415–434, 1963.
- [38] W. Loh, “Classification and regression trees,” *Wiley Interdiscip. Rev. Data Min. Knowl. Discov.*, vol. 1, no. 1, hal. 14–23, 2011.
- [39] C. J. Burges, “From ranknet to lambdarank to lambdamart: An overview.”
- [40] T. M. Mitchell, “Machine learning. WCB,” 1997.