

Daftar Pustaka

- Baroni, M.P., Sanchis, G.J., de Assis, S.J., dos Santos, R.G., Pereira, S.A., Sousa, K.G., Lopes, J.M. (2015). Factors associated with scoliosis in schoolchildren: a cross-sectional population-based study. *Journal Epidemiology*. 25(3):212-20.
- Barrios, C., Cortes, S., Perez-Encinas, C., Escriva, M.D., Benet, I., Burgos, J., Hevia, E., Piza, G., Domenech, P. (2011). Anthropometry and body composition profile of girls with nonsurgically treated adolescent idiopathic scoliosis. *Spine*, 36 (18): 1470-1477.
- Bjure, J., Grimby, G., Nachemson, A. (1968). Correction of body height in predicting spirometric values in scoliotic patients. *Scandinavian Journal of Clinical and Laboratory Investigation*. 21(2):191-2.
- Brazolino, M.A.N., Maia, T.C., Junior, C.J., Cardoso, I.M., Junior, J.L.B., Batista P.R. (2015). Low body weight in patients with adolescent idiopathic scoliosis. *Salus Journal Health and Science*, 1(1): 82-86.
- Budi, D.P.S. (2011). *School Screening For Scoliosis In Surabaya*. Artikel Penelitian. Surabaya : Universitas Airlangga.
- Burwell, R.G., Aujla, R.K., Grevitt, M.P., Dangerfield, P.H., Moulton, A., Randell, T.L., Anderson, S.I. (2009). Pathogenesis of adolescent idiopathic scoliosis in girls - a double neuro-osseous theory involving disharmony between two nervous systems, somatic and autonomic expressed in the spine and trunk: possible dependency on sympathetic nervous system and hormones with implications for medical therapy. *Scoliosis*, 31;4:24. doi: 10.1186/1748-7161-4-24.
- Brox, J.I. (2014). Idiopathic Scoliosis. *European Medical Journal Rheumatology*, 1: 48-55.
- Cheung, C.S.K., Lee, W.T.K., Tse, Y.K. (2003). Generalized osteopenia in adolescent idiopathic scoliosis—association with abnormal pubertal growth, bone turnover, and calcium intake? *Spine (Phila Pa 1976)* 2006;31:330–8.
- Cobb, N.J. (2001). *Adolescence: Continuity, Change, And Diversity* (4th ed.). California: Mayfield Publishing Company.
- Demerath, E.W., Schubert, C.M., Maynard, L.M., Sun, S.S., Chumlea, W.C., Pickoff, A., Czerwinski, S.A., Towne, B., Siervogel, R.M. (2006). Do changes in body mass index percentile reflect changes in body composition in children? Data from The Fels longitudinal study. *Pediatrics*, 117: 487–495.

- Departemen Kesehatan Republik Indonesia. (2006). *Pengukuran Antropometri*. Jakarta : Salemba Medika.
- Departemen Kesehatan Republik Indonesia. (2010). *Kesehatan Remaja: Problem dan Solusinya*. Jakarta: Salemba Medika.
- Ellis, K.J. (2001). Selected body composition methods can be used in field studies. *American Society for Nutritional Sciences*, 131: 1589S-1595S.
- Gibson, R.S. (2005). *Principle of Nutritional and Assessment*. Newyork : Oxford University Press.
- Heymsfield, S.B., Lohman, T.G., Wang, Z., Going, S.B. (2005). *Human Body Composition*. Champaign, IL: Human Kinetics.
- Heyward, V. (2001). ASEP methods recommendation: body composition assessment. *An International Electronic Journal*, 4 (4).
- Hume, K. (2008). Scoliosis: To screen or not to screen. *British Journal of School Nursing*, 3 (5). EBSCO database.
- Hurlock, E. B. (1994). *Psikologi Perkembangan: Suatu Pendekatan Sepanjang Rentang Kehidupan*. Jakarta : Erlangga..
- Ippolito, E., Versari, P., and Lezzerini, S. (2006). The role of rehabilitation in juvenile low back disorders. *Pediatric Rehabilitation*, 9 (3): 174-184.
- Irianto, Koes. (2007). *Gizi dan Pola Hidup Sehat*. Bandung : Yrama Widya.
- Jebb, S., McCarthy, D., Fry, T., Prentice, A.M. (2004). New body fat reference curves for children. *Obesity Reviews* (NAASO Suppl) A156
- Konieczny, M.R., Senyurt, H., Krauspe, R. (2013). Epidemiology of adolescent idiopathic scoliosis. *Journal of Children's Orthopaedics*, 7 (1): 3–9.
- Koya, C. and Rawlinson. (2009). Pain management: an adolescent scoliosis patient. *Journal of Perioperative Practice*, 19 (7): 205-212. ProQuest Nursing & Allied Health Source.
- Kyle, U.G., Bosaeus, I., De Lorenzo, Antonio. (2004). Bioelectrical impedance analysis Part I: Review of principles and methods. *Clinical Nutrition*, 23: 1226–1243.

- Marfell-Jones, M., Olds, T., Stewart, A., and Carter, J.L. (2006). *International Standards for Anthropometric Assesment*. Potchefstroom : International Society for the Advancement of Kinanthropometry.
- Martelli, R.C. and Traebert, J. (2006). Descriptive study of backbone postural changes in 10 to 16 year-old schoolchildren. *Revista Brasileira de Epidemiologia*. vol.9 no.1.
- Matusik, E., Durmala, J. Matusik P., Pitrowski, J. (2012). Evaluation of nutritional status of children and adolescent with idiopathic scoliosis. *Ortopedia*, 14: 351-362.
- Matusik, E., Durmala, J. Matusik P. (2016). Association of body composition with curve severity in children and adolescents with idiopathic scoliosis (IS). *Nutrients*, 8: 71.
- McCance, Kathryn, L., and Sue, E. H. (2006). *Pathophysiology: The Biologic Basis for Disease in Adults and Children*. 5th ed. St. Louis, MO : Elsevier Mosby.
- Moore, K.L., Dalley, A. F., and Agur, A.M.R. (2010). *Clinically Oriented Anatomy*. Philadelphia : Wolter Kluwer Health.
- Mukaromah, S. (2011). *Pengalaman Psikososial Remaja Penyandang Skoliosis Di Wilayah Karesidenan Surakarta, Jawa Tengah: Studi Fenomenologi*. Tesis. Depok : Universitas Indonesia.
- National Scoliosis Foundation and DePuy Spine. (2009). *Scoliosis Media and Community Guide* [booklet]. Stoughton.
- Nicolopoulos, K.S., Burwell, R.G., Webb, J.K. (1985). Stature and its components in adolescent idiopathic scoliosis. Cephalo-caudal disproportion in the trunk of girls . *Journal of Bone and Joint Surgery Bristish*, 67 (4):594 – 601 .
- Normelli, H., Sevastik, J., Ljung, G. (1985). Anthropometric data relating to normal and scoliotic Scandinavian girls . *Spine (Phila Pa 1976)*, 10 : 123 – 6.
- Qiu, Y., Sun, X., Qiu, X., Li, W., Zhu, Z., Zhu, F., Wang, B., Yu, Y., Qian, B: (2007). Decreased circulating leptin level and its association with body and bone mass in girls with adolescent idiopathic scoliosis. *Spine*, 32(24):2703-2710.
- Ramirez, M., Martinez-Llorens, J., Sanchez, JF., Bago, J., Molina, A., Gea, J., Caceres, E. (2013). Body composition in adolescent idiopathic scoliosis. *Europe Spine Journal*, 22: 324-329.

- Santoso, S. (2004). *Kesehatan Dan Gizi*. Jakarta : Rineka Cipta.
- Sari, S. (2013). *Swiss Ball Exercise Dan Koreksi Postur Tidak Terbukti Lebih Baik Dalam Memperkecil Derajat Skoliosis Idiopatik Daripada Klapp Exercise Dan Koreksi Postur Pada Anak Umur 11 – 13 Tahun*. Tesis. Denpasar : Universitas Udayana.
- Schlenker, E. (2000). *Nutrition and The Aging Adult*. in: B. Worthington-Roberts, S. Williams (Eds.) *Nutrition Throughout The Life Cycle*. 4th edition. McGraw-Hill, 350–383.
- Simanjuntak, C.A. (2008). Sudah Perlukah Dilakukan Skrining Skoliosis pada Anak Sekolah Lanjutan Tingkat Pertama. *DEXA Media*, 1 (1).
- Siu, K.C.C., Tak, K.L.W., Kit, T.Y, Ping, T.S., Man, L.K., Guo, X., Qin, L., Chun, Y.C.J. (2003). Abnormal peri-pubertal anthropometric measurements and growth pattern in adolescent idiopathic scoliosis: a study of 598 patients. *Spine*, 28(18): 2152-2157.
- Slaughter, M.H., Lohman, T.G., Boileau, R.A., Horswill, C.A., Stillman, R.J., Van Loan, M.D., Bembien, D.A. (1988). Skinfold equations for estimation of body fatness in children and youth. *Human Biology*, 60: 709-723.
- Smith, B.G., Hakim-Zargar, M., Thomson, J.D. (2009). Low body mass index: a risk factor for superior mesenteric artery syndrome in adolescents undergoing spinal fusion for scoliosis . *Journal of Spinal Disorder and Technique*, 22 : 144 – 8.
- Snell, Richard S. (2006). *Anatomi Klinik* Ed. 6. Jakarta : EGC.
- Soucacos, P.N., Zacharis, K., Soultanis, K., Gelalis, J., Xenakis, T., Beris, A.E. (2000). Risk factors for idiopathic scoliosis: review of a 6-year prospective study. *Orthopedics*, 23(8): 833-838.
- Stanhope, M., and Lancaster, J. (2004). *Community & Public Health Nursing* (6th ed.). St.Louis: Mosby, Inc.
- Sugiyono. (2001). *Metode Penelitian Administrasi*. Bandung: Alfabeta.
- Suh, S.W., Modi, H.N., Yang, J.H., Hong, J.Y. (2011). Idiopathic scoliosis in Korean schoolchildren: a prospective screening study of over 1 million children. *Europe Spine Journal*, 20 (7): 1087-1094.
- Supariasa, I.D.N, Bakri, B., Fajar, I. (2002). *Penilaian Status Gizi*. Jakarta : EGC.

- Tachdjian's. (2008). *Scoliosis. In: Pediatric Orthopaedics* (4th edition). Saunders Elsevier, Philadelphia.
- Tam EMS. Yeung KH. Tang S. Lam TP. Ng BKW. Lee SKM. Qiu Y and Cheng JCY. (2016). Lower muscle mass and body fat in adolescent idiopathic scoliosis are associated with abnormal leptin bioavailability. *Spine*, 41(11): 940-946.
- Taspinar, F., Saracoglu, I., Afsar, E., Okur, E.O., Seyyar, G.K. (2017). Assessing the Relationship between Body Composition and Spinal Curvatures in Young Adults. *Archive of Sports Medicine and Physiotherapy*, 2(1): 010-015.
- Trobisch, P., Sues, O., Schwab, F. (2010). Idiopathic scoliosis. *Deutsches Arzteblatt International*. 107(49): 875-884.
- UWSP University Health Service. (2000). *Body Composition*. Diakses dari <http://www.wellnessproposals.com/health-promotion/presentations/bodycomposition-body-fatness-presentation.pdf>
- Wei-jun, W., Xu, S., Zhi-wei, W. (2012). Abnormal anthropometric measurement and growth pattern in male adolescent idiopathic scoliosis. *Europe Spine Journal*, 21: 77-83.
- Weili, Y., He, B., Yao, H., Dai, J., Cui, J., Ge, D., Zhang, Y., Li, L., Guo, Y., Xiao, K., Fu, X., Ma, D. (2007). Waist-to-height ratio is an accurate and easier index for evaluating obesity in children and adolescent. *Obesity* 15: 748-752.
- WHO. (2007). *WHO Reference 2007 for Child and Adolescent*. WHO, Geneva.
- WHO. (2008). *Waist Circumference and Waist-Hip Ratio Report of WHO Expert Consultation*. World Health, 8-11.
- Wolpert, DK. (2006). *Scoliosis Surgery: The Definitive Patient's Reference*. Swordfish Communications. ISBN 978-0-9741955-2-0.
- Wong, D.L. (2008). *Buku Ajar Keperawatan Pediatrik*. Vol.1 & 2. alih bahasa, Agus Sutarna, Neti Juniarti, H.Y.Kuncara; editor edisi bahasa Indonesia, Egi Komara Yudha. Ed.6. Jakarta: EGC.
- Wong, H.K., Hui, J.H., Rajan, U., Chia, H.P. (2005). Idiopathic scoliosis in Singapore schoolchildren: A prevalence study 15 years into the screening program. *Spine*, 30 (10): 1188-1196.

- Yim, A.P., Yeung, H.Y., Hung, V.W. (2012). Abnormal skeletal growth patterns in adolescent idiopathic scoliosis: a longitudinal study until skeletal maturity. *Spine*, 37:E1148–54.
- Ylikoski, M. (2003). Height of girls with adolescent idiopathic scoliosis . *Eur Spine*, 12 : 288 – 91.
- Zaimul Haq, Ahmad. (2011). *Waspadai Tulang Belakang Miring*. 20 Januari 2011. <http://issuu.com/surya-epaper/docs/surya>.
- Zaina, F., Donzelli, S., Lusini, M., Vismara, L., Capodaglio, P., Neri, L., Negrini, S. (2013). Adolescent idiopathic scoliosis and eating disorders: is there a relation? Results of a cross-sectional study. *Research in Developmental Disabilities*, 34(4):1119-24.
- Zheng, R.D., Chen, Z.R., Chen J.N. Lu, Y.H., Chen, J. (2012). Role of body mass index, waist-to-height, waist-to-hip ratio in prediction of nonalcoholic fatty liver disease. *Gastroenterology Research and Practice*, 362147.