

DAFTAR PUSTAKA

- Acevedo, P. et al., 2005. Environmental constraints in the colonization sequence of roe deer (*Capreolus capreolus* Linnaeus, 1758) across the Iberian Mountains, Spain. *Journal of Biogeography*, 32(9), pp.1671–1680.
- Ackers, S.H. et al., 2015. The evolution of mapping habitat for northern spotted owls (*Strix occidentalis caurina*): A comparison of photo-interpreted, Landsat-based, and lidar-based habitat maps. *Remote Sensing of Environment*, 156, pp.361–373.
- Agoramoorthy, G., Alagappasamy, C. & Hsu, M.J., 2004. Can proboscis monkeys be successfully maintained in captivity? A case of swings and roundabouts. *Zoo Biology*, 23(6), pp.533–544.
- Alikodra, H.S., 2002. *Pengelolaan Satwaliar* Jilid I., Bogor: Yayasan Penerbit Fakultas Kehutanan Institut Pertanian Bogor.
- Anwar, J. Damanik, S. J. Hisyam, N. Whitten, A.J., 1984. *Ekologi ekosistem Sumatra*, Yogyakarta: Gadjah Mada University Press.
- Aplin, P., 2004. Remote sensing as a means of determining ecological conditions. *Bioscience*, 17(7), pp.444–449.
- Asner, G.P., Scurlock, J.M.O. & Hicke, J. a, 2003. Global synthesis of leaf area index observations : *Global Ecology & Biogeography*, 12(2003), pp.191–205.
- Atmoko, T. A. Ma'ruf, I. Syahbani. Rengku, M.T., 2007. *Kondisi habitat dan penyebaran bekantan (*Nasalis larvatus Wurm*) di Delta Mahakam, Kalimantan Timur* D. S. K. Sidiyasa, M. Omon, ed., Bogor: Pusat Litbang Hutan dan Konservasi Alam.
- Bailey, J.A., 1984. *Principles of Wildlife Management* 1st ed., Wiley; 1 edition (March 2, 1984). Available at: <https://www.amazon.com/Principles-Wildlife-Management-James-Bailey/dp/0471016497>.
- Baldwin, R.A., 2009. Use of maximum entropy modeling in wildlife research. *Entropy*, 11(4), pp.854–866.
- Barclay, H.J., 1998. Conversion of total leaf area to projected leaf area in lodgepole pine and Douglas-fir. *Tree Physiology*, 18, pp.185–193. Available at: <http://www.heronpublishing.com>.
- Barnes, T.G., 2000. Landscape ecology and ecosystems management. *Cooperative Extension Service University of Kentucky, Lexington*, p.8.
- Beadle, C.L., 1993. Growth analysis. In *Photosynthesis and Production in a Changing Environment*. Dordrecht: Springer Netherlands, pp. 36–46. Available at: http://link.springer.com/10.1007/978-94-011-1566-7_3 [Accessed November 15, 2016].
- Bennett, E.L. & Sebastian, A.C., 1988. Social organization and ecology of proboscis monkeys (*Nasalis larvatus*) in mixed coastal forest in Sarawak. *International Journal of Primatology*, 9(3), pp.233–255. Available at: <http://link.springer.com/10.1007/BF02737402> [Accessed October 1, 2016].
- Bismark, M. Iskandar, S., 2002. Kajian total populasi dan struktur sosial bekantan (*Nasalis larvatus*) di Taman Nasional Kutai, Kalimantan Timur. *Buletin Penelitian Hutan*, 631, pp.17–29.
- Bismark, M. Soerianegara, I. Sastradipradja, D. Suratmo, F. G. Alikodra, H. S.

- Pawitan, H., 1994. *The potency of mangrove forest habitat to the proboscis monkey's food source at Kutai National Park*, Bali.
- Bismark, M., 1995. Analisis Populasi Bekantan (*Nasalis Larvatus* Wurmbe). *Rimba Indonesia*, p.vol. XXX.
- Bismark, M., 2009. *Biologi Konservasi Bekantan (Nasalis larvatus)*,
- Bismark, M., 2004. Daya Dukung Habitat dan Adaptasi Bekantan (*Nasalis Larvatus* Wurmbe). *Penelitian Hutan dan Konservasi Alam*, 1(3), pp.309–320.
- Bismark, M., 2005. Model pengukuran biomasa populasi primata. *Jurnal Penelitian Hutan dan Konservasi Alam II*, 5, pp.491–496.
- Bismark, M., 1987. ngkah-laku makan bekantan (*Nasalis larvatus*) di hutan bakau, Taman Nasional Kutai, Kalimantan Timur. *Buletin Penelitian Hutan*, 492, pp.1–10.
- Bismark, M., 1997. *Pengelolaan habitat dan populasi bekantan (Nasalis larvatus) di Cagar Alam Pulau Kaget*, Kalimantan Selatan.
- Bismark, M., 1980. *Populasi dan tingkahlaku bekantan (Nasalis larvatus) di Suaka Margasatwa Tanjung Puting*, Kalimantan Tengah,
- Bloom, S., 1999. *In praise of primates*, Köln, Germany: Könemann. Available at: <http://kindlebooku.com/read-online-in-praise-of-primates.html>.
- Chivers, D.J., 1974. The siamang in Malaya: a field study of a primate in tropical rain forest. *Primata*, 4, pp.1–335.
- Clement, M.J. & Castleberry, S.B., 2013. Estimating density of a forest-dwelling bat: A predictive model for Rafinesque's big-eared bat. *Population Ecology*, 55(1), pp.205–215.
- Clutton-Brock, T.H., 1977. Some aspects of intraspecific variation in feeding and ranging behaviour in primates. *Primate Ecology*, pp.539–556.
- Curran, A.L.M. et al., 2004. Lowland Forest Loss in Protected Areas of Indonesian Borneo Published by : American Association for the Advancement of Science Stable URL : <http://www.jstor.org/stable/3836134>. , 303(5660), pp.1000–1003.
- Danoedoro, P., 2012. *Pengantar Penginderaan Jauh Digital*, Yogyakarta: Andi Yogyakarta.
- Darsidi, A., 1984. Pengelolaan hutan mangrove di Indonesia. *Prosiding Seminar II Ekosistem Mangrove*, pp.19–28.
- Digital Globe, *WorldView-2*,
- Elith, J., Leathhwick, J.R., 2009. Evolution and ecology of species range limits. *Annual Review of Ecology, Evolution, and Systematics*, 40(2009), pp.415–436. Available at: <http://www.annualreviews.org/doi/pdf/10.1146/annurev.ecolsys.110308.120317>.
- Elith, J. et al., 2006. Novel Methods Improve Prediction of Species ' Distributions from Occurrence Data. *Ecography*, 29(129–151). Available at: <http://www.jstor.org/stable/3683475>.
- Estes, L.D. et al., 2008. Habitat selection by a rare forest antelope: A multi-scale approach combining field data and imagery from three sensors. *Remote Sensing of Environment*, 112(5), pp.2033–2050.
- Fleagle, J.G., 1980. Locomotion and Posture. In *Malayan Forest Primates*. Boston,

- MA: Springer US, pp. 191–208. Available at: http://link.springer.com/10.1007/978-1-4757-0878-3_7 [Accessed November 14, 2016].
- Forman, R.T.T., 1995. Some general principles of landscape and regional ecology. *Landscape Ecol*, 10, p.133.
- Fourcade, Y. et al., 2014. Mapping species distributions with MAXENT using a geographically biased sample of presence data: A performance assessment of methods for correcting sampling bias. *PLoS ONE*, 9(5), pp.1–13.
- Franklin, J., 2009. Mapping species distributions. Spatial inference and prediction. *Ecology, biodiversity and conservation*, 53(9), p.340.
- Franklin, J., 2009. *Mapping species distributions: spatial inference and prediction*, New York, United States of America: Cambridge University Press.
- Freeland, A.W.J., 1976. Pathogens and the Evolution of Primate Sociality Published by : The Association for Tropical Biology and Conservation Stable URL : <http://www.jstor.org/stable/2387816> . , 8(1), pp.12–24.
- Gobron, N. Michel, M. V., 2009. Leaf Area Index. *Assessment of the status of the development of the standards for the Terrestrial Essential Climate Variables*, 10, p.22.
- Goodman, M. et al., 1998. Toward a phylogenetic classification of primates based on DNA evidence complemented by fossil evidence. *Molecular Phylogenetics and Evolution*, 9(3), pp.585–598.
- Greaves, G.J., Mathieu, R. & Seddon, P.J., 2006. Predictive modelling and ground validation of the spatial distribution of the New Zealand long-tailed bat (*Chalinolobus tuberculatus*). *Biological Conservation*, 132(2), pp.211–221.
- Gron, K., 2009. Primate Factsheets: Proboscis monkey (*Nasalis larvatus*) Taxonomy, Morphology, & Ecology. , p.1. Available at: http://pin.primate.wisc.edu/factsheets/entry/proboscis_monkey [Accessed September 10, 2015].
- Gusmira, D., 2013. *Kajian Dampak Keberadaan Jalan terhadap Lingkungan Kawasan Taman Nasional Tesso Nilo, Provinsi Riau*. UGM.
- Hall, L. S. Krausman, P. R. Morrison, M.L., 1997. The Habitat Concept and A Plea for Standard Terminology. *Wildlife Society Bulletin*, 25, p.173–182.
- Helms, J.A., 1998. *The Dictionary of Forestry*, The Society of American Forestry & CABI Publishing Bethesda.
- Hernández-Clemente, R. et al., 2014. A Novel Methodology to Estimate Single-Tree Biophysical Parameters from 3D Digital Imagery Compared to Aerial Laser Scanner Data. *Remote Sensing*, 6(11), pp.11627–11648. Available at: <http://www.mdpi.com/2072-4292/6/11/11627/> [Accessed October 13, 2016].
- Huck, M. et al., 2008. Predicting European badger *Meles meles* sett distribution in urban environments Predicting European badger *Meles meles* sett distribution in urban environments. *Wildlife Biology*, 14(2), pp.188–198. Available at: //000258027500003.
- Huete, A.R., 1988. A soil-adjusted vegetation index (SAVI). *Remote Sensing of Environment*, 25, pp.295–309.
- Hurd, J.D. et al., 2006. Tidal wetland classification from landsat imagery using an integrated pixel-based and object-based classification approach. *American*

- Society for Photogrammetry and Remote Sensing - Annual Conference of the American Society for Photogrammetry and Remote Sensing 2006: Prospecting for Geospatial Information Integration*, 2, pp.590–600. Available at: <http://www.scopus.com/inward/record.url?eid=2-s2.0-84869014796&partnerID=tZOtx3y1>.
- Hutchins, M., 2003. *Grzimek 's Animal Life Encyclopedia*,
- Iskandar, E., 2006. *Habitat dan populasi owa Jawa (*Hylobates moloch*) di TN Gunung Halimun Salak Jawa Barat*. Institut Pertanian Bogor.
- Jaberg, C. & Guisan, A., 2001. Modelling the distribution of bats in relation to landscape structure in a temperate mountain environment. *Journal of Applied Ecology*, 38(6), pp.1169–1181.
- Jain, A.K., 1989. *Fundamentals of digital image processing*, Prentice Hall.
- Jolly, A., 1972. *The evolution of primate behavior*, New York: Mac-Millan Publishing Co., Inc.
- Jones, D.B. et al., 2004. Asian primate classification. *Int. J. Primatol*, 25(1), pp.97–164.
- Juntti, T.M. & Rumble, M. a, 2006. Arc Habitat Suitability Index Computer Software. *Natural Resources Research*, (September), p.31.
- Kamal, M., Phinn, S. & Johansen, K., 2016. Assessment of multi-resolution image data for mangrove leaf area index mapping. *Remote Sensing of Environment*, 176, pp.242–254.
- Kemp, K.K., 1993. *Environmental modeling with GIS: A strategy for dealing with spatial continuity*,
- Kern, J.A., 1964. Observation on the habit of the proboscis monkey, *Nasalis larvatus* (Wurmb), made in the Brunei Bay area. *Borneo Zoologica*, 49, pp.183–192.
- Kerr, J.T. & Ostrovsky, M., 2003. From space to species: ecological applications for remote sensing. *Trends in Ecology & Evolution*, 18(6), pp.299–305.
- Kocev, D. et al., 2009. Using single- and multi-target regression trees and ensembles to model a compound index of vegetation condition. *Ecological Modelling*, 220(8), pp.1159–1168.
- Kuemmerle, T. et al., 2010. European Bison habitat in the Carpathian Mountains. *Biological Conservation*, 143(4), pp.908–916. Available at: <http://dx.doi.org/10.1016/j.biocon.2009.12.038>.
- Kupfer, J.A. et al., 2004. Identifying the Biodiversity Research Needs Related to Forest Fragmentation. *Geospatial Wetlands Impacts & Mitigation Forecasting Models*, (November 2016), p.218.
- Lambin, E.F., Geist, H. & Rindfuss, R.R., 2006. Introduction: Local Processes with Global Impacts. In *Land-Use and Land-Cover Change*. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 1–8. Available at: http://link.springer.com/10.1007/3-540-32202-7_1 [Accessed September 24, 2016].
- Lear, R. & Turner, T., 1977. *Mangrove of Australia*, University of Queensland Press.
- Ma'ruf, A., 2004. *Studi perilaku bekantan (*Nasalis larvatus*) di daerah Balikpapan dan sekitarnya*, Kalimantan Timur.

- MacKinnon, J.R. & MacKinnon, K.S., 1980. Niche Differentiation in a Primate Community. In *Malayan Forest Primates*. Boston, MA: Springer US, pp. 167–190. Available at: http://link.springer.com/10.1007/978-1-4757-0878-3_6 [Accessed November 14, 2016].
- Manel, S., Williams, H.C. & Ormerod, S.J., 2001. Evaluating presence absence models in ecology; the need to count for prevalence. *Journal of Applied ecology*, 38, pp.921–931.
- Marfai, M.A., 2011. *Pengantar Pemodelan Geografi* 1st ed., Yogyakarta: Badan Penerbit Fakultas Geografi (BPGF) Fakultas Geografi Universitas Gadjah Mada.
- McGarigal, K. & Marks, B.J., 1994. FRAGSTATS: spatial pattern analysis program for quantifying landscapes Structure. *General Technical Report PNW-GTR-351. U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. Portland, OR, 97331(503)*, p.134.
- McNeely, J.A. et al., 1990. *Conserving the World'S Biological Diversity World Bank Conserving the World'S*,
- Meijaard, E., Nijman, V. & Supriatna, J., 2008. *Nasalis larvatus*. The IUCN Red List of Threatened Species 2008: , 8235(Boonratana 1993), pp.9–11. Available at: <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T14352A4434312.en>.
- Meijaard, E. Nijman, V., 2000. Distribution and conservation of the proboscis monkey (*Nasalis larvatus*) in Kalimantan, Indonesia. *Biological Conservation*, 92, pp.15–24.
- Meijaard, E. et al., 2006. *Hutan Pasca Pemanenan*,
- Meixler, M.S., Bain, M.B. & Walter, M.T., 2009. Predicting barrier passage and habitat suitability for migratory fish species. , 220, pp.2782–2791.
- Mitsch, W. J. Gosselink, J.G., 1984. *Wetlands* V. N. C. Reinhold, ed., New York.
- Mochizuki, S. Murakami, T., 2011. Change in habitat selection by Japanese macaques (*Macaca fuscata*) and habitat fragmentation analysis using temporal remotely sensed data in Niigata Prefecture, Japan. *International Journal of Applied Earth Observation and Geoinformation*, 13(4), pp.562–571. Available at: <http://dx.doi.org/10.1016/j.jag.2011.02.006>.
- Monteith, J.L., Unsworth, M.H. & Arnold, E., 1990. *Principles of environmental physics* Second Edi. E. Alnold, ed., London.
- Morrison, M. L. Marcot, B. G. Mannan, R.W., 2006. *Wildlife-Habitat Relationships* Third Edit., Washington: Island Press.
- Morrison, M. L. Marcot, B. G. Mannan, R.W., 2007. Wildlife-habitat Relationships: Concepts and Applications. *The Condor*, 109(4), p.980. Available at: [http://www.bioone.org/perlserv/?request=get-abstract&doi=10.1650%2F0010-5422\(2007\)109%5B980%3AWRCAA%5D2.0.CO%3B2](http://www.bioone.org/perlserv/?request=get-abstract&doi=10.1650%2F0010-5422(2007)109%5B980%3AWRCAA%5D2.0.CO%3B2) [Accessed November 14, 2016].
- Murai, T., 2006. Mating behaviors of the proboscis monkey (*Nasalis larvatus*). *American Journal of Primatology*, 68(8), pp.832–837. Available at: <http://doi.wiley.com/10.1002/ajp.20266> [Accessed November 13, 2016].
- Nagendra, H., 2001. Using remote sensing to assess biodiversity. *Remote Sensing*, 22(12), pp.2377–2400.

- Napier, J.R. & Napier, P.H. (Prue H., 1985. *The natural history of the primates*, British Museum (Natural History).
- Newton-Cross, G., White, P.C.L. & Harris, S., 2007. Modelling the distribution of badgers *Meles meles*: comparing predictions from field-based and remotely derived habitat data. *Mammal Review*, 37(1), pp.54–70. Available at: <http://doi.wiley.com/10.1111/j.1365-2907.2007.00103.x> [Accessed December 9, 2016].
- Northrup, J.M., Stenhouse, G.B. & Boyce, M.S., 2012. Agricultural lands as ecological traps for grizzly bears. *Animal Conservation*, 15(4), pp.369–377.
- Nowak, R.M., 1999. *Primates of The World*, Loondon. Available at: <https://books.google.co.id/books?id=unODoWa7CM4C&pg=PP7&lpg=PP7&dq=Primates+of+The+World.+The+Johns+Hopkins+University+Press.+Ba+ltimore+and+London+nowak&source=bl&ots=lv7e8XUYj2&sig=cgtG5DVAR-Kx8tHUQXCwbHbICQc&hl=id&sa=X&ved=0ahUKEwjTp9qkp6XQAhUIs48KHxbD>.
- Osborne, P.E., Alonso, J.C. & Bryant, R.G., 2001. Modelling landscape-scale habitat use using GIS and remote sensing: a case study with great bustards. *Journal of Applied Ecology*, 38(2), pp.458–471. Available at: <http://doi.wiley.com/10.1046/j.1365-2664.2001.00604.x>.
- Ottaviani, D. et al., 2009. Modelling semi-aquatic vertebrates' distribution at the drainage basin scale: The case of the otter *Lutra lutra* in Italy. *Ecological Modelling*, 220(2), pp.111–121.
- Panigada, S. et al., 2008. Modelling habitat preferences for fin whales and striped dolphins in the Pelagos Sanctuary (Western Mediterranean Sea) with physiographic and remote sensing variables. *Remote Sensing of Environment*, 112(8), pp.3400–3412.
- Pearson, R.G., 2008. Species' distribution modeling for conservation educators and practitioners. *Lessons in Conservation*, (3), pp.54–89.
- Phillips, S. J., Anderson, R. P., Schapire, R.E., 2006. Maximum Entropy Modeling of Species Geographic Distributions. *International Journal of Global Environmental Issues*, 6(2–3), pp.231–252.
- Phillips, S., Dudík, M. & Schapire, R., 2004. A maximum entropy approach to species distribution modeling. *Proceedings of the twenty-first ...*, pp.655–662. Available at: <http://dl.acm.org/citation.cfm?id=1015412>.
- Phillips, S.J. & Dudík, M., 2008. Modeling of species distribution with Maxent: new extensions and a comprehensive evaluation. *Ecography*, 31(December 2007), pp.161–175.
- Poplar-Jeffers, I.O. et al., 2009. Culvert Replacement and Stream Habitat Restoration: Implications from Brook Trout Management in an Appalachian Watershed, U.S.A. *Restoration Ecology*, 17(3), pp.404–413. Available at: <http://doi.wiley.com/10.1111/j.1526-100X.2008.00396.x> [Accessed December 9, 2016].
- Primack, R. B. Supriatna, J. Indrawan, M. Kramadibrata, P., 2007. *Biologi Konservasi*, Jakarta: Yayasan Obor Indonesia.
- Raemaekers, J.J. & Chivers, D.J., 1980. *Socio-Ecology of Malayan Forest Primates*.

- In *Malayan Forest Primates*. Boston, MA: Springer US, pp. 279–316. Available at: http://link.springer.com/10.1007/978-1-4757-0878-3_10 [Accessed November 14, 2016].
- Rajanathan, R.B.E.L., 1990. Notes on the sosial behaviour of wild proboscis monkeys (*Nasalis larvatus*). *Malayan Nature Journal*, 44, pp.35–44.
- Rich, P.M., 1990. Characterizing plant canopies with hemispherical photographs. *Remote Sensing Reviews*, 5(1), pp.13–29.
- Ross, J., 1981. *The Radiation Regime and Architecture of Plant Stands*, Boston, London: Junk Publishers.
- Rushton, S. P.; Ormerod, S. J. & Kerby, G., 2004. New paradigms for modelling species distributions? *J. Appl Ecol*, 41, pp.193–200.
- Salter, R. E. Mackenzie, N. A. Nightingale, N. Aken, K.M. Chai, P., 1985. Habitat use, ranging behaviour and food habits of proboscis monkey *Nasalis larvatus* (Van Wurmb) in Sarawak. *Primates*, 26(4), pp.436–451.
- Sanders, L. ed., 2007. *Model in Spatial Analysis*, ISTE Ltd London: Wiley. Available at: http://yp3mz3xt4m.search.serialssolutions.com/?ctx_ver=Z39.88-2004&ctx_enc=info:ofi/enc:UTF-8&rft_id=info:sid/summon.serialssolutions.com&rft_val_fmt=info:ofi/fmt:kev:mtx:book&rft.genre=book&rft.title=Models+in+Spatial+Analysis&rft.au=.
- Setiono, S.B., Said, S., E., 2008. Jumlah Individu dan Kelompok Bekantan (*Nasalis Larvatus*, Wurmb) di Taman Nasional Danau Sentarum Kabupaten Kapuas Hulu. , 1, pp.269–277.
- Shaw, J., 1985. *Introduction to Wildlife Management*, New York: McGraw-Hill Book Company.
- Soehartono, T., M., 2003. *Pelaksanaan konvensi CITES di Indonesia | OPAC Integrasi | Online Public Access Catalog | Universitas Gadjah Mada*, JICA. Available at: http://opac.lib.ugm.ac.id/index.php?mod=book_detail&sub=BookDetail&act=view&typ=html&obyek_id=1&buku_id=188724.
- Soendjoto, A. M. Alikodra, H. S. Bismark, M. Setijanto, H., 2006. Aktivitas harian bekantan (*Nasalis larvatus Wurmb.*) di hutan karet Kabupaten Tabalong, Kalimantan Selatan. *Biota*, XI (2), pp.101–109.
- Soendjoto, M. A. Akhdiyati, M. Haitami. Kusumajaya, I., 2001. Bekantan di hutan galam: Quo vadis? *Warta Konservasi Lahan Basah*, pp.18–19.
- Soendjoto, M.A., 2005. *Adaptasi Bekantan (Nasalis larvatus Wurmb) Terhadap Hutan Karet: Studi Kasus Di Kabupaten Tabalong, Kalimantan Selatan*. Institut Pertanian Bogor. Available at: http://www.iwf.or.id/opini_adaptasi_bekantan.pdf.
- Soerianegara, I. D. Sastradipradja, H. S. Alikodra. Bismark, M., 1994. Studi habitat sumber pakan dan perilaku bekantan (*Nasalis larvatus*) sebagai parameter ekologi dalam mengkaji sistem pengelolaan habitat hutan mangrove di Taman Nasional Kutai. *Pusat Penelitian Lingkungan Hidup Institut Pertanian Bogor*.
- Srivathsan, A., 2011. Proboscis Monkeys (*Nasalis Larvatus* (Wurmb , 1787)) Have Unusually High-Pitched Vocalizations. , 59(2), pp.319–323.
- Sterman, J.D., 2002. *System Dynamics: Systems Thinking and Modeling for a*

- Complex World. *In Proceedings of the ESD Internal Symposium.*, (May), pp.1–29.
- Stoms, D.M. & Estes, J.E., 1993. A remote sensing research agenda for mapping and monitoring biodiversity. *International Journal of Remote Sensing*, 14(10), pp.1839–1860.
- Sugiyono, 2007. *Metode Penelitian pendidikan pendekatan kuantitatif, kualitatif, dan R&D*, Bandung: Alfabeta.
- Sukardjo, S., 1987. Tanah dan status hara di hutan mangrove Tritis, Indramayu, Jawa Barat. *Rimba Indonesia XXI*, p.(2-4): 12-23.
- Sulistyo, B., Gunawan, T. & Danoedoro, P., 2013. Modeling of Percentage of Canopy in Merawu Catchment Derived. *Forum Geografi*, 27(1), pp.23–32.
- Supriatna, J. Wahyono, E.H., 2000. *Panduan lapangan primata Indonesia*, Yayasan Obor Indonesia.
- Teder, T. et al., 2007. Monitoring of biological diversity: A common-ground approach. *Conservation Biology*, 21(2), pp.313–317.
- Turner, W. et al., 2003. Remote sensing for biodiversity science and conservation. *Trends in Ecology and Evolution*, 18(6), pp.306–314.
- Verburg, P.H. et al., 2006. Modeling Land-Use and Land-Cover Change.
- Vina, A. et al., 2008. Evaluating MODIS data for mapping wildlife habitat distribution. *Remote Sensing of Environment*, 112(5), pp.2160–2169.
- Wainer, G.A., 2009. *Discrete-Event Modeling and Simulation: A Practitioner's Approach*, New York: CRC Pres. Available at: https://books.google.co.id/books?hl=en&lr=&id=E0T5dmTrG7EC&oi=fnd&pg=PP1&dq=Discrete-Event+Modelling+and+Simulation:+A+Practitioner%E2%80%99s+Approach,+CRC&ots=ceBhVpD6Hn&sig=M6wgl1yo7k-fg7OkfA9bKTA2HHR4&redir_esc=y#v=onepage&q&f=false.
- Wainwright, J. & Mulligan, M., 2004. *Environmental modelling finding simplicity in complexity*, New York: John Wiley & Sons.
- Walker, B.K., Jordan, L.K.B. & Spieler, R.E., 2009. Relationship of Reef Fish Assemblages and Topographic Complexity on Southeastern Florida Coral Reef Habitats. *Journal of Coastal Research*, 10053(10053), pp.39–48.
- Walters, D.M., Roy, A.H. & Leigh, D.S., 2009. Environmental indicators of macroinvertebrate and fish assemblage integrity in urbanizing watersheds. *Ecological Indicators*, 9(6), pp.1222–1233.
- Wang, T. et al., 2016. Textural-spectral feature-based species classification of mangroves in Mai Po nature reserve from worldview-3 imagery. *Remote Sensing*, 8(1), pp.1–15.
- Whitten, A.J., 1982. Diet and feeding behaviour of kloss gibbon on Siberut Island, Indonesia. *Animal Behavior*, 30, pp.182–198.
- Whitten, A.J., 1982. Diet and Feeding Behaviour of Kloss Gibbons on Siberut Island, Indonesia. *Folia Primatologica*, 37(3–4), pp.177–208. Available at: <http://www.karger.com/doi/10.1159/000156032> [Accessed November 14, 2016].
- Wiens, J.A., 1995. Habitat fragmentation: island. *Ibis*, (137 (I)), pp.97–104.
- Wilcove, D.S., McLellan, C.H. & Dobson, A.P., 1986. Habitat Fragmentation in

- the temperate zone. *Conservation Biology: The Science of Scarcity and Diversity*, pp.237–256.
- Wilson, C. C. Wilson, W.L., 1975. The influence of selective logging on primates and some other animals in East Kalimantan. *Folia Primatol*, 23, pp.245–274.
- Wulder, M.A. et al., 2004. High Spatial Resolution Remotely Sensed Data for Ecosystem Characterization. *BioScience*, 54(6), pp.511–521. Available at: [http://bioscience.oxfordjournals.org/cgi/doi/10.1641/0006-3568\(2004\)054\[0511:HSRRSD\]2.0.CO;2](http://bioscience.oxfordjournals.org/cgi/doi/10.1641/0006-3568(2004)054[0511:HSRRSD]2.0.CO;2).
- Xiaoxia, S. Jixian, Z. Zhengjun, L., 2004. An Object-Oriented Classification Method On High Resolution Satellite Data. *ACRS*, pp.347–350.
- Yasuma, S., 1989. *The present situation of proboscis monkey *Nasalis larvatus*, concern with it's distributions in and around the Bukit Soeharto protection forest*,
- Yeager, C.P., 1989. Feeding ecology of the proboscis monkey (*Nasalis larvatus*). *International Journal of Primatology*, 10(6), pp.497–530. Available at: <http://link.springer.com/10.1007/BF02739363> [Accessed October 1, 2016].
- Yeager, C.P., 1990. Proboscis monkey (*Nasalis larvatus*) social organization group structure. *Am. J. of Primatology*, 20, pp.73–86.