



DAFTAR PUSTAKA

- Adeniji, O.T.P, E.Odo. P.E and B. Ibrahim..2011. Genetic relationships and selection indices for cassava root yield in Adamawa state. Nigeria. *African Journal of Agricultural Research* 6 (13): 2931-2934.
- Ademosun, O. C., M. O. Jimoh, and O. J. Olukunle. 2012. Effect of physical and mechanical properties of cassava tubers on the performance of an automated peeling machine. *International Journal of Development and Sustainability*. 1. 3: 810-822.
- Adetan, D. A., L. O. Adekoya, L. O and O. B. Aluko. 2003, Characterisation of some properties of cassava root tubers, *Journal of Food Engineering*, Vol. 59, pp. 349-353.
- Agahie, A. E., K.P. Baiyeri and R. O. Ogbuji. 2011. Correlation analysis of tuber yield and yield related characters in two cassava (*Manihot esculenta* Crantz) morphological-types grown under nine weed management systems in the Guinea savanna zone of Nigeria. *Journal of Applied Bioscience* 48:3316-3321.6pp. www.m.elewa.org/JABS/2011/48/6.pdf
- Ahmadi, A. M, Joudi. and M, Janmohamadi. 2009. Late defoliation and wheat yield: little evidence of post anthesis source limitation. *Filed Crop Res.* 113: 90-93.
- Aji, T.G. dan S, Susanto. 2009. Pengaruh jumlah cabang terhadap pertumbuhan vegetatif dan generatif rosella. Makalah Seminar Departemen Agronomi dan Horikultura, Institut Pertanian Bogor.
- Al Afas. N, N. Marron, and R. Ceulemans. 2006. Clonal variation in stomatal characteristics related to biomass production of 12 poplar (*Populus*) clones in a short rotation coppice culture. *Journal Environmental and Experimental Botany*, vol. 58, pp. 279-286.
- Alizadeh, O. K, Farsinejad. S, Korani and A, Azarpanah. 2013. A study on source-sink relationship, photosynthetic ratio of different organs on yield and yield components in bread wheat (*Triticum aestivum* L). *International Journal of Agriculture and Crop Sciences*. 5 (1):69-79.
- Alves A.G.C. 2001. Cassava botany and physiology. In: Hillocks RJ, Thresh JM, Bellotti AC, eds. Cassava biology, production and utilization. Wallingford: CABI, 67–90
- Alves. A.A.A, 2002. Cassava botany and physiology, in: R.J. Hillock, J.M. Thresh, A.C. Bellotti (Eds), Cassava: Biology, Production and Utilization, CABI International Oxford, 2002, pp. 67-89.
- Amarullah, 2013. Teknologi Budidaya Singkong Gajah (*Manihot esculenta* Crantz). Prosiding *Seminar Nasional UNS "Akselerasi Pembangunan Pertanian Berkelanjutan menuju Kemandirian Pangan dan Energi"* di UNS Solo, Tanggal 17 April 2013
- Amarullah, 2015. Teknologi Budidaya Singkong Gajah (*Manihot esculenta* Crantz)..*Scientific Journal of Agricultural Science*. AgroUPY. 6 (2): 45-54.
- Ambarwati, E dan H. Murti. 2001. Analisis korelasi dan koefisien lintas sifat-sifat agronomi terhadap komposisi kimia umbi iles-iles (*Amorphophallus variabilis*). *Ilmu Pertanian* 8 (2): 55-61



- Angela, C. W. A, Rogers. M, Rees and C, P. Osborne. 2016 How can we make plants grow faster? A source–sink: perspective on growth rate. *Journal of Experimental Botany*, 67 (1):31–45. doi:10.1093/jxb/erv447
- Aranjuelo, I. Á, Sanz-Sáez. I, Jauregui. J, J. Irigoyen. J, L. Araus. M, Sánchez-Díaz and G, Erice. 2013. Harvest index, a parameter conditioning responsiveness of wheat plants to elevated CO₂. *Journal of Experimental Botany*. 64(7):1879-1892. doi:10.1093/jxb/ert081
- Arifin. 2003. Dasar-dasar Klimatologi. Fakultas Pertanian Universitas Brawijaya .Malang
- Armas . R, E. Ortega, and R. Rodes. 1988. Fisiologis vegetal. La Habana (ed) Pueblo y Education.
- Aslam. M. S.B. Lows, and L.A. Hunt. 2011. Effect of leaf age on photosynthesis and transpiration on cassava (*Manihot esculenta* Crantz). *Canadian Journal of Botany* 55(17):2288-2295. doi: 10.1139/b77-260
- Asli, D.E, A, Eghdami and A, Houshmandfar. 2011. Evaluation of Sink and Source Relationship in Different Rice (*Oryza Sativa* L.) Cultivars. *Advances in Environmental Biology*, 5(5): 912-919.
- Aspiazú, I. T. Sedyama. Jr. Ribeiro. A.A; Silva. G. Concenco. E.A. Ferreira. L. Galon, L. A.F. Silva. E.T. Borges and. W.F. Araujo. 2006. Photosynthetic activity of cassava plants under weed competition. *Encyclopedia Britannica*. Inc.
- Ashraf, M. M. Akbar. M, Salim. 1994. Genetic improvement in physiological traits of rice yield. Slafer CA (ed). Genetic improvement of yield crops. *Marcel Dekker Incorporates* New York, pp: 413-455.
- Atkins C.A and P, M.C Smith.. 2007. Translocations in Legumes; Assimilates, Nutrients and Signaling Molecules. *Plant Physiology* 144:550-561.
- Bahri, S. 2010. Klorofil. Diktat Kuliah. Kapita selekta Kimia organik. Universitas Lampung. 48p.
- Balitkabi. 2012. Deskripsi varietas unggul kacang-kacangan dan umbi-umbian. Balitkabi Malang. 179p.
- Bart. R.I. 1964. Carbohydrate utilization as a factor in plant growth. *Aust. J. Biol. Sci.* 17: 867-877.
- Basuchaudhuri, P. 2016. Source-sink relationship in soybean. Centre for Info Bio Technology (CIBTech). *Indian Journal of Plant Science*. 19-25. <http://www.cibtech.org/jps.htm>
- Bioversity International, CIAT. 2009. Key access and utilization descriptors for cassava genetic resources. Bioversity International, Rome, Italy, International Center for Tropical Agriculture (CIAT), Cali, Colombia.
- Bio Research Development. 2012. Peluang pasar ubikayu. **Info dari Big Cassava**. <http://kebun-singkong.blogspot.co.id/archive.html> (diakses 14 Februari 2014)
- BPS, 2013. Statistik indonesia. Badan Pusat Statistik. Jakarta-Indonesia; hal 216-218
- Boerboom, B. W. J. 1978. A model of dry matter distribution in cassava (*Manihot esculenta* Crantz). *Neth.J.agric.Sei.* 26 (3): 267-277.



- Connor, D.J and J.H Cock, 1981. The Response of cassava to water shortage II. Canopy dynamics, *Field Crops Res*, 4: 285-296
- Cardoso, P. Mirione, E. Ernesto. M. Massaza, F. Cliff, J. Haque, M.R and Bradbury, J.H. 2005. Processing of cassava roots to remove cyanogens. *Journal of food composition and analysis*, vol. 18. P: 451-460.
- Casteleanos, R.L and, F.F Betancourt, 1985. Relationship between origins of cassava cuttings and yields in Camaguey-Tunas Penplane, *Viandas Tropicales* 8: 57-67.
- Ceballos, H., E. Okogbenin, J. C. Perez, L. A. Becerra, and D. Debouck. 2010. Cassava. In: *Bradshaw Journal*, ed. Root and tuber crops. New York: Springer, 53–96
- Chávez, A. L., T. Sánchez., G. Jaramillo., J. M. Bedoya., J. Echeverry., E. A. Bolaños., H. Ceballos.and C. A. Iglesias. 2005. Variation of quality traits in cassava roots evaluated in landraces and improved clones. *Euphytica* 143:125-133.
- CIAT. 1991. Cassava Program Annual Report for 1987-1991. Cali, Colombia
- Cock, J.H and S.Ross. 1975. Ecophysiology of Cassava' dalam Symp on Ecophysiology of Tropical Crops hal 1-14 Communications Division of CEPLAC. Km 22. Rodovia. Ilheus-Itabuna. Bahia. Brazil.
- Cock, J.H. 1976. Characteristics the high yielding Cassava varieties. *Eksperimental Agriculture*. 23: 135-143.
- Cock, J.H., D. Franklin, G. Sandoval and P. Juri, 1979. The ideal cassava plant for maximum yield. *Crop Science* 19: 271-279.
- Cock, J. H. 2001. Stability of performance of cassava genotypes in cassava breeding: *Proceedings workshop held in Phillipines*, 4-7: 100-106.
- Connor, D.J and J. Palta, 1981. The Response of cassava to water shortage III. Stomatal Control of plant water status, *Field Crops Res*, 4: 297-311.
- Dahniya, M. T., Oputa, C. O. and Hahn, S. K., 1982. Investigating source-sink relations in cassava by reciprocal grafts. *Expl. Agric*, 18: 399-402.
- Daie, J. 1985. Carbohydrate partitioning and metabo-lism in crops. *Hort. Rev.* 7, 69-108
- De Bruijn, G.H. and T.S. Dharmaputra, 1974. The Mukibat system, a high-yielding method of cassava production in Indonesia. *Neth.J.agric.Sci.* 22.- * 89-100
- DJTPKP. 2013. Pedoman teknis pengelolaan dan Produksi ubikayu, DJTP Kementerian Pertanian. 144p.
- Doku, E.V., 1965. Breeding for yield in cassava. I. Indices of yield. *Ghana J.of Science* 5(1): 42-59
- Dueck, T.A., Janse, J, Schapendonk, AHCM, Kempkes, FLK, Eveleens-Clark, BA, Scheffers, CP, Pot, S, Trouwborst, G, Nederhoff, EM, and Marcelis, LFM. 2010b. Lichtbenutting van onder LED en SON-T belichting. Wageningen: Wageningen UR Glastuinbouw/Plant Dynamics BV, Rapporten GTB 1040



- Dufour, D.L. 1988. Cyanide content of cassava (*Manihot esculenta* Crantz) cultivar used by tukanoan indians in northwest Amazonia. In *Economic Botany*, vol. 42. P; 255-266.
- Duncan, W.G., R.S. Loomis, W.A. Williams and R. Hanau, 1967. A model for simulating photosynthesis in plant communities. *Hilgardia* 38: 181-205.
- Edet, M.A. H.T, Eniola. S.T.O. Lagoke and H. Tarawali. 2015. Relationship of cassava growth parameters with yielded and yiled related compenents and harvest time in Ibadan Southwestern. *Journal of Natural Sciences Research*. 5 (9): 87-92
- Enyi B.A.C 1972. The Effect of Spacing on Growth, Development and Yield of Single and Multishoot Plants of Cassava. In: Root tuber Yield and Tributes. *East Africa Agric. and Forestry Journal* 1: 27-34.
- Enyi, B.A.C. 1973.'Growth rate of three cassava (*Manihot esculenta* Crantz) varietties under varying population densities', *J. Agric. Sci. UK*, 81: 15-28
- Evans, L.T J, Biogham. P, Jackson. And J, Sutherland. 1972. Effect of awns and draought on the supply of photosynthate and its distribution within wheat ears. *Annals Applied Biol.* 70: 67-76.
- FAO. 2011. FAO's initiative on soaring food prices Guide for policy and programmatic actions at country level to address high food prices.
- Foyer, C. H, N. Galtier and P. Quick. 1994. Modifications in carbon assimilation, carbon partitioning and total biomass as a result of over-expression of sucrose phosphate synthase in transgenic tomato plants. *Plant Physiol.* 105: 123-135
- Fukuda, W.M.G., C.L. Guevara, R. Kawuki, and M.E. Ferguson, 2010. Selected morphological and agronomic description for the characterization of cassava. International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria. 19pp.
- Garcia.M dan M. Rodriguez. 1983. Comparative study on cassava cutting from different parts of the plant, *Viandas Tropicale* 6: 39-49
- Gardner, F.P., R.B. Pearce, and R.L. Mitchell. 2008. Physiology of Crop Plants (Fisiologi Tanaman Budidaya. Alih Bahasa: Herawati Susilo). Universitas Indonesia Press, Jakarta
- Gasper, J,-Ann, and B,B. Schweig, 1985. Wages and Public Higher Education Defined Benefit Pension Plans, 11(2): 19-38.
- Goldsworthy.P.R dan N.N. Fisher, 1994. Fisiologi Tanaman Budidaya Tropika. Gadjah Mada University Press. Yogyakarta. Indonesia. 874p.
- Gunadi N dan Subhan. 2011. Pengaruh jumlah cabang per tanaman terhadap pertumbuhan dan hasil tiga varietas paprika. *Jurnal Hortikultura* 21: 124-134.
- Hahn, S.K., 1977. A quantitative approach to source potentials and sink capacities among reciprocal grafts of sweet potato varieties. *Crop Science* 17: 559-562.
- Hahn, S.K., E.R. Terry, K. Leuschner, I.O. Akobundu, C. Okali and R. Lai, 1979. Cassava improvement in Africa. *Field Crops Research* 2: 193-226.



- Hairiah, K., M. Van Noordwijk and G. Cadisch. 2000. Crop yield, C and N balance of three types of cropping systems on an Ultisol in North Lampung. Netherlands *Journal of agricultural science*. 46: 3-17.
- Haryanti. S dan T. Meirina. 2009. Optimalisasi Pembukaan Porus Stomata Daun Kedelai (*Glycine max* L Merrill) Pagi dan Sore Hari. *BIOMA*. 11 (1):18-23 .
- Hartt. C.F., H.P. Korstchak and G.O. Burr. 1964. Effect of defoliation, deradiation and darkening the blade upon translocation of C⁺⁺, in sugarcane. *Plant Physiol*. 39:15-22.
- Hayford, M. A. 2009. Growth, Yield and Quality of Cassava as Influenced by Terramend 21, Poultry Manure and Inorganic Fertilizer Master of Agronomy Thesis, School of Graduate Studies, Kwame Nkrumah University of Science And Technology (KNUST), Kumasi
- Henry, G and C. Hershey. 2002. Cassava in South America and Caribbean, In: R.J. Hillocks, J.M. Thresh, A.C. Bellotti (Eds.), *Cassava: Biology, Production and Utilization*, CABI Publishing Oxon, UK and New York, USA, 2002, pp. 17-40.
- Herzog, H. 1982. Relation of source and sink during the grain-filling period in wheat and some aspects of the regulation. *Physiol Plant* 56:155-160.
- Heuberger, C. 2005. Cyanide content of cassava and fermented product with focus on attieke and attieke Gaarba. Dissertation of technology Zurich. Swiss federal institute of technology
- Howeler.R.H and Cadavid. L.P. 1983. Accumulation and distribution of dry matter and nutrients during a 12 month growth cycle of cassava, *Field Crops Research*. 7: 123-139.
- Hopkinson. 1964. *Plant in action*. A resource for teachers and students of plant science Second edition. Australian Society of Plant Scientists, New Zealand Society of Plant Biologists, and New Zealand Institute of Agricultural and Horticultural Science 2010
- Hopkins WG. 2004. *Introduction to Plant Physiology*. New York: John Wiley & Sons. Inc.
- Howeler.R.H and Cadavid. L.P. 1983. Accumulation and distribution of dry matter and nutrients during a 12 month growth cycle of cassava, *Field Crops Research*. 7: 123-139.
- Hozyo and Kato. 1976. The interrelationship between source and sink of the grafts of wild type and improved cultivar of *Ipomea*. *Proc. Crop Sci. Japan*. 45: 117-123
- Humphries, F. C. 1963. Dependence of net assimilation rate on root-growth isolated leaves. *Ann. Bot. N.S.* 37(5): 175-183
- Humphries. F.C. and G.N. Thorne. 1964. The effect of root formation on photosynthesis detached leaves. *Ann Bot*. 28:391-400.
- Hutami, F.D. dan Harijono. 2014. Pengaruh penggantian larutan dan konsentrasi NaHCO₃ terhadap penurunan kadar sianida pada pengolahan tepung ubikayu. *Jurnal pangan dan agroindustri*. Vol. 2 No.4p. Oktober 2014.
- Iglesias, D.J. I, Lliso. F,R. Tadeo and M, Talon. 2002. Regulation of photosynthesis through source-sink imbalance in citrus is mediated by carbohydrate content in leaves. *Physiologia plantarum*. 116: 563-572.



- Ijoyah M.O., R.I.B. and C, A. Iheadindueme. 2012. Productivity of Cassava-Sweet Potato Intercropping System as Influenced by Varying Lengths of Cassava Cutting at Makurdi, Nigeria. *Journal of Biology, Agriculture and Healthcare*. 2(5): 87-92.
- Ikeh.A.O, Udaeyo.N.U, Udoh.E.I, Iboko.K.O and Udounang. P.I. 2012. Growth and Yield of Cassava (*Manihot esculenta* Crantz) as Influenced by The Number of Shoots Retained Per Stand on as Ultisol. *Journal Nature and Science*. 2012; 10 (8).
- Iland, P., Dry, P., Proffitt, T. and Tyerman, S., 2011. The Grapevine from the Science to the Practice of Growing Vines for Wine. Patrick Iland Wine Promotions, Adelaide, South Australia
- Indira, P and T. Ramanujam. 1979. Effect of photoperiod on tuberization in cassava, *J.Root Crops* 5 (1/2): 39-42.
- Indira, P. and S.K. Sinha, 1970. Studies on the initiation and development of tubers in *Manihot esculenta* Crantz. *Indian Journal of Plant Physiology* 13: 24-39.
- Indira, P. and T. Kurian. 1977. A comparative study of the anatomical changes associated with tuberization in the roots of cassava and sweet potato. In: Leakey, C.L.A. (Ed.), *Proceedings of the third symposium of the International Society for Tropical Root Crops*, UTA, Ibadan, Nigeria, 2-9 December 1973, pp. 78-80.
- Irikura, Y., J. H. Cock and K. Kawano. 1979. The physiological basis of genotype temperature interactions in cassava, *Field Crops Res*, 2: 227-239.
- Jarvis, P.G and J.I. Morison. 1982. The control of transpiration and photosynthesis by the stomata. In: Jarvis PG, Mansfield TA (Eds.). *Society of experimental Biology Seminar 8*. Cambridge University Press. Pp: 247-279.
- Kakani. V.G, K.R. Reddy, D. Zhao, and A.R. Mohammed. 2003, "Effects of ultraviolet-B radiation on cotton (*Gossypium hirsutum* L) morphology and anatomy", *Annals of Botany*, 91 817-826.
- Karnjanakorn. K, 1968. Planting positions and weed control in cassava plantations at Si Racha, Ph.D. Dissertation, Kasetsart University, Thailand.
- Kawano K., W.M.G. Fukuda and U. Cenpukdee. 1987. Genetic and environmental effects on dry matter content of cassava root. *Crop Science* 27: 69-74.
- Kawano, K. 1990. Harvest index and evolution of major food crops cultivars in the tropics. *Euphytica* 46: 195-202.
- Keating, B.A. 1982. 'Environmental effects on growth and development of cassava (*Manihot esculenta* Crantz). Assimilate distribution and storage organ yield. *Field Crops.Res.* 5: 283-292.
- Keating, B.A and J.B. Evenson.1979. Effect of soil temperature on sprouting and sprout elongation of stem cuttings of cassava. *Field crops Res.* 2: 241-252.



- Khairullah. I. 2012. Gatra fisiologis dan agronomis pengerauh pengendalian keracunan besi padi sawah di lahan rawa pasang surut sulfat asam. (Disertasi). Universitas Gadjah Mada Yogyakarta.
- Khan, A.A., and G.R. Sagar. 1966. Distribution of C¹⁴ labelled products of photosynthesis during the commercial life of the tomato crop. *Ann. Bot.* 30:727-743.
- Khan, A.A., and G.R. Sagar. 1969. Alteration rice pattern of distribution of photosynthetic products in the tomato by manipulation of the plant. *Ann Bot.* 33:753-762.
- King.R.W, I.F. Warldlaw and I.T. Evans.1969. Effect of assimilate utilization on photosynthesis in wheat. *Planta.* 77: 261-276.
- Kirschbaum, M.U.F and R.W.Pearcy.1988. Gas exchange analysis of the relative importance of stomatal and biochemical factors in photosynthetic induction in *Alocasia macrorrhiza*. *Plant Physiol.* 86(3): 782-785.
- Kleinhenz, V. K, Katroschan. F, Schitt and H, Stitzel. 2006. Biomass accumulation and partitioning of tomato under protected cultivation in the humid tropics. *European Journal of Horticultural Science.* 71(4): 173-182.
- Kozlowski, T.T. and S.G. Pallardy. 1997. Physiology of woody trees. 2nd ed. Academic Press, San Diego, CA.
- Lafitte. H.R., and R.I. Travis. 1984. Photosynthesis and assimilate partitioning and closely related lines of rice exhibiting different sink-source relationship. *Crop. Sci.* 24: 447-452
- Lahai, M.T. I, J. Ekanayake and J, P.C. Koroma. 2013. Influence of canopy structure on yield of cassava cultivars at various toposequences of an inland valley agro-ecosystem. *Journal of Agricultural Biotechnology and Sustainable Development.* 5 (3): 36-47. Doi: 10-5897/JABSD10.006.
- Lakso, A. N and J. A. Flore. 2003. Carbohydrate partitioning and plant growth. Pp. 21-30. In: Baugher, T.A. and S. Singh (eds.). Concise encyclopedia of temperate tree fruit. Food Product Press, New York, N.Y.
- Lambers, H., F.S. Chapin and T.L. Pons. 2008. Plant Physiological Ecology. Springer. New York.
- Lebot, V. 2009.. Tropical root and tuber crops: cassava, sweet potato, yam and aroids. CABI publication. Amazon.com.413.
- Legesel. H, L. Gobeze, A. Shegro dan N. Geleta, 2011. Impact of Planting Position and Planting Material on Root Yield of Cassava (*Manihot esculenta* Crantz). *Journal of Agricultural Science and Technology,* 5(4):1939-1250, USA .
- Lemoine, R., Sylvain L.C., Rossitza A., Fabienne, D., Thierry, A., Nathalie, P., Jean-Louis, B., Maryse, L., Pierre, C.T., Laurence, M., Mireille, F., Christine, G., Pauline, L., Jonathan, P., and Mickael, D. 2013. Source-to-sink transport of sugar and regulation by environmental factors. *Front. Plant Sci.* 4: 272
- Lenis, J.L. F, Calie. G, Jaramillo. J,C. Perez. H, Caballos and J,H. Cock. 2006. Leaf retention and cassava productivity. *Filed Crops Res.* 95: 126-134.



- Li, M.H. G, Hoch and C, Korner. 2002. Source-sink removal affects mobile carbohydrates in Pinus cembreat the Swiss treeline. *Trees*. 16: 331-337.
- Li, R., P. Guo, M. Baum, S. Grando, and S. Ceccarelli. 2006. Evaluation of chlorophyll content and fluorescence parameters as indicators of drought tolerance in barley. *Agricultural Sciences in China*. 5(10): 751-757
- Li.T, E. Heuvelink and L.F.M. Marcelis. 2015. Quantifying the source-sink balance and carbohydrate content in three tomato cultivars. *Front Plant Sci*. 6:461. Doi; 10:3889/Fpls. 2015. 00416.
- Ly, J. 1998. Cassava roots (*Manihot esculenta* Crantz) for pigs; A short review on its nutrient content. *Revista Computadorizada de Produccion Porcina*, 5: 1-13
- Mahon, J.D, S.B. Lowe and I.A. Hunt, 1976. 'Variation in rate of photosynthetic CO₂ uptake in cassava and selected species of Manihot', *Photosynthetic*, 11: 131-138.
- Mahon, J.D., S.B. Lowe, L.A. Hunt and M. Thiagarajah, 1977a. Environmental effects on photosynthesis and transpiration in attached leaves of cassava (*Manihot esculenta* Crantz). *Photosynthetic* 11(2): 121-130.
- Marschner, H. 1995. Mineral Nutrition of Higher Plants. Academic Press.131-183p
- Mason, T.G., 1972. A note on growth and the transport of organic substances in bitter cassava (*Manihot utHissima*), *Scientific proceedings Royal Dublin society* 17: 105-112
- Mason,T.G and F. Phillip. 1957. The migratiuon of solute. *Bot. Rev.* 3:47-78.
- Matsuda R, K.Suzuki, A.Nakano, T. Higashide and M.Takaichi. 2011. Responses of leaf photosynthesis and plant growth to altered source–sink balance in a Japanese and a Dutch tomato cultivar. *Scientia Horticulturae*,127: 520-527. DOI: 10.1016/j.scienta.2010.12.008
- Max,J, F.J. L, Schmidt. U, N. Mutwiwa and K, Kahlen. 2016. Effects of shoot pruning and inflorescence thinning on plant growth, yield and fruit quality of greenhouse tomatoes in a tropical climate. *Journal of Agriculture and Rural Development in the Tropics and Subtropics*. 117(1): 45-56.
- Mengel, K. 1996. Potassium movement within plant and its importance in assimilate transport. pp: 408–409. In R. D. Munson (ed). Potassium In Agriculture. *American Soils Society*. 1207 p.
- Mogllner, I., J. D. P. Arias, A.D. Gotuzzo and J. A. Acosta. 1967. Influencia de la parte aerea de *Manihot flabellifolia* en la formacion de raices reservantes de *Manihot esculenta* utilizado como pie. *Bonplandia*. 2(10):137-142.
- Montagnac, J.A. Davis, C.R. and S.A.Tanumihardjo. 2008. Processing techniques to reduce toxicity and antinutrients of cassava for use as a staple food. In *Comprehensive reviews in Food science and food safety*, vol 8, p: 17-27.
- Morsis H., M.E. Medri, C.J. Marur, P.H. Caramori, A.M.D.A Riberio and J.C. Gomes. 2004. Modifications on leaf anatomy of coffea arabica caused by shade pigeonpea. (*Cajanus cajan*). *Brazillian Archives of biology and technology*, 47(6): 863-871.



- Moss, D. N and R. B. Musgrave. 1971. Photosynthesis and Crop Production. *Advances in Agronomy*. 23: 317-336.
- Murugeswari, R., Balakrishnan, V. and Vijayakumar, R. 2006. Studies to assess the suitable conservation method for tapioca leaves for effective utilization by ruminants Livestock Research for Rural Development 18(32).
- Naves-Barbiero, C.C. 2000. Fluxo de seve e conductancia estomatica de duas especies lenhosas sempre-verdes no campo sujoe cerrado. R. Bras. *Physiol. Veg.* 12(2):119-134
- Nazar, R. S, Umar. N, A. Khan and O, Sareer. 2015. Salicylic acid supplementation improves photosynthesis and growth in mustard through changes in proline accumulation and ethylene formation under drought stress. *South Africa Journal of Botany* 98: 84-94
- Nio, S. A dan Y. Banyo. 2011. Konsentrasi klorofil daun sebagai indikator kekurangan air pada tanaman. *Jurnal ilmiah Sains*. 11(2): 165-173.
- Nishio, J.N; Sun, J; and T.C.Vogelmann. 1994. Photoinhibition and the light environment within leaves, In: Baker, N.R; Bowyer, J.R. (Eds) Photoinhibition of Photosynthesis. Oxford: BIOS Scientific Publisher. P.1-24.
- Noerjiwati, K. 2012. Keragaan klon-klon ubikayu dengan potensi hasil umbi dan pati tinggi sebagai bahan baku industri. Prosiding; Peningkatan Daya Saing dan Implementasi Pengembangan Komoditas Kacang dan Umbi Mendukung Pencapaian Empat Sukses Pembangunan Pertanian. Balitkabi. Malang. Indonesia
- Nosberger, J and F.C, Humpries. 1965. The influence of removing tuber and dry matter production and net assimilation rate at potato plants. *Annual Botany*. 29: 579-588.
- Nugroho, W. H., H. Y. Sugito, B. Guritno dan W. H. Utomo. 1985. Teknologi Budidaya Ubkayu secara monokultur dan Tumpangsari. Dalam. Prosiding pengembangan Ubikayu di Jawa Timur. Pusat Penelitian Tanaman Ubi-ubian Univ. Brawijaya IDRC. IDC ford Foundation.
- Ntawuruhunga, P., and Dixon, A.G.O. 2010. Quantitative variation and interrelationship between factors influencing cassava yield. *Journal of Applied Bioscience*, 26, 1594-1602.
- Okogbenin. T.L.Setter. M, Ferguson. R, Mutegi. A,C. Alves. H, Caballos. and M, Fregene. 2010. Phenotyping cassava for adaptration to drought. Generation Challenge Programme. Mexico City. Pp. 381-400
- Oliveira E.C., and Miglioranza E. 2014. Stomatal Density in Six Genotypes of Cassava. *International Journal of Engineering Science and Innovative Technology* (IJESIT) 3(3): 305-308
- Oliveira, C.M. and C.A. Priestley. 1988. Carbohydrate re-serves in deciduous fruit trees. *Hort. Rev.* 10, 403-430
- Onwueme I.C, 2002. Cassava in Asia and the Pacific, in: R.J. Hillock, J.M. Thresh, A.C. Bellotti (Eds.), Cassava: Biology, Production and Utilization, CABI Publishing Oxon, UK and New York, USA, 2002, pp. 55-65.



- Osorio S, Ruan YL, and A.R. Fernie. 2014. An update on source-to-sink carbon partitioning in tomato. *Frontiers in Plant Science*, 5: 516. 06 October 2014 | <http://dx.doi.org/10.3389/fpls.2014.00516>
- Pangestuningsih dan Syamsudin. 1980. Studi kerapatan tanaman ubikayu (*Manihot esculenta* Crantz) dengan sistem FAN design. *Jurnal Agronomi Indonesia*. IPB. Bogor..458: 1-9.
- Patrick JW, and K. Colyvas. 2014. Crop yield components -photoassimilate supply- or utilisation limited-organ development? *Functional Plant Biology*, 41: 893-913
- Peroni, N. ; Kageyama, P.Y.and A.Begossi,.2007. Molecular differentiation, diversity, and folk classification of "sweet" and "bitter" cassava (*Manihot esculenta*) in Caicara and Caboclo management systems (Brazil). *Genetic Resources and Crop Evolution*, 54 (6): 1333-1349
- Prasetiaswati, N., A. Munip, B. S. Radjit, N. Saleh dan Y. Widodo. 2008. Kelayakan usahatani ubikayu sistem mukibat. *Pros. Sem. Nas. Pengembangan kacang-kacangan dan umbi-umbian*. Surakarta, 7 Agustus 2008. P: 223-233.
- Prihatman, K. 2000. Ketela Pohon/Singkong (*Manihot utilissima* Pohl). Available at: <http://www.ristek.go.id> (diakses tangga 6 Februari 2015)
- Rahmatian, A. M, Delshad and R, Salchi. 2014. Effect of grafting on growth, yield and fruit quality of single and double stemmed tomato plants grown hydroponically. *Horticulture Environment and Biothechnology*. 55(2): 115-119.
- Ramanujam, T and P, Indira. 1983. Canopy structure on growth and development of cassava (*Manihot esculenta* Crantz). *Turrialba*. 33: 321-326.
- Ramanujam, T. and S. P. Ghosh 1990. Investigations of source-sink relations in cassava using reciprocal grafting. *Expl. Agric*, 26: 189-195.
- Régnier, C., 2011. Valorisation des ressources alimentaires tropicales (feuilles et tubercules) chez le porc. Thèse (INRA Antilles-Guyane, Unité de Recherches Zootechniques – URZ)
- Retherford, R. D and M. K. Choe. 1993. *Statistical Model for Causal analysis*. New York; John Wiley & Sons. Inc.
- Ristono dan Amarullah. 2011. *Singkong Gajah Berjuang*. Petrogas Press. Cetakan II. Balikpapan. 202p.
- Rogers, D. J and S. G. Appan. 1973. *Manihot and manihotoides (Euphorbiaceae) a computer assisted study*. Organization for flora neotropica. Monograph no. 13. New York. 278 pp.
- Roja, A. 2009. *Ubikayu; varietas dan teknologi budidaya*. Pelatihan Spesifik Lokalita Kabupaten 50 Kota Sumatera Barat, Payakumbuh, 7-18 Oktober 2009
- Ross, C , J.H. Cock and G. Sandoval, 1976. Leaf fall in cassava. *Expl.Agric*. 12: 395-400.
- Rubatzky V.E, and M.Yamaguchi. 1995. *Sayuran Dunia 1*. Ed ke-2. Herison Catur, penerjemah. Bandung: ITB. Terjemahan dari: *World Vegetables: Principles, Production, and Nutritive Values*. p: 291-296



- Saleh, N., S. A. Rahayuningsih dan M. Adie. 2010. Peningkatan Produksi dan Kualitas Umbi-umbian. Balai Penelitian Kacang-kacangan dan Umbi-umbian (Balitkabi). Malang. Indonesia p:1-21
- Saleh, N., S.A. Rahayuningsih dan M Adie. 2012. Peningkatan produksi dan kulaitas umbi-umbian. Balai Penelitian Kacang-kacangan dan Umbi-umbian. BALITKABI. Malang p:1-21
- Salisbury, F.B and C.W Ross. 1995. Plant Physiology. 3 ed. Wadsworth Publishing Co. Belmont California
- Salunkhe, D.K. and S.S. Kadam. 1998. Handbook of vegetable science and technology : production, composition, storage, and processing. New York : Marcel Dekker, c1998. 721 p
- Samonte, S.O. P.B., Wilson and A.M. Mc. Clung. 1998. Path analysis of yield and yiel and yield-related traits of fitten diverse rice genotype. *Crop Sci.* 38: 1130-1136.
- SAS. 2002. *Statistical Analysis System for Windows* 9.0. SAS Institute Inc. SAS Campus Drive. Cary. North Carolina. USA
- Scofield G.N, Ruuska S.A., Aoki N., Lewis D.C., Tabe LM, and C.L.D.Jenkins.. 2009. Starch storage in the stems of wheat plants: localization and temporal changes. *Annals of Botany*, 103: 859-868. doi: [10.1093/aob/mcp010](https://doi.org/10.1093/aob/mcp010)
- Serna, L and C. Fenoli, 2000. Stomatal development in Arabidopsis; how to make a functional pattern. Trends in *Plant Science*, vol. 5, pp. 458-460.
- Shanumugha A and C. Srinivasan. 1973. Influence of Number of Shoot Per Plant on the Growth and Yield of Cassava. *Farm Journal of India*, 1973 14(7):17-19.
- Sharkawy, M. A. 2004. Cassava Biology and Physiology. *Plant Moluculer Biology*. 56: 481-501
- Sharkey, T.D and Raschke, K. 1981. Effect of light quality on stomatal opening in leaves of *Xanthium strumarium* L. *Plant Physiol.*, 68 (5):1170-1174.
- Sholihin. 2009. Peluang klon-klon ubikayu dalam menunjang industri bioetanol. Risalah Seminar Penelitian dan pengembangan tanaman pangan 2007-2008. Balai Penelitian dan Pengembangan Pertanian 324-341.
- Singh, R. K., and B. D. Chaudhary, 1979. Biometrical Methodes in Quantitative Genetic Analysis. Kalyani Pub. Ludhiana, New Delhi. 303 p.
- Sinha, S.K. and T.V.R. Nair, 1971. Leaf area during growth and yielding capacity of cassava. *Indian J.Genetics and Plant Breeding* 31(1): 16-20
- Sinthuprama. S and C. Tiraporn, Improving the productivity of cassava in Thailand, in: Cassava in Asia, Its Potential and Research Development Needs, Proceedings of Regional Workshop, Bangkok, Thailand, June 5-8, 1984, pp. 277-287.
- Smith, O.B. 1988. A review of ruminant responses to cassava-based diets Cassava as livestock feed in Africa. *Proceedings* of the IITA/ILCA/University of Ibadan. Workshop on the Potential Utilization of Cassava as Lives tock Feed in Africa. International Institute of Tropical Agriculture. Ibadan, Nigeria.



- Sugito. 1990. Effect of removing leaves on cassava yield. Nat. Seminar on cassava pre and post harvest Tech. Res. And Development, held in Lampung. Indonesia. Feb. 15, 1990. Pp:189-2008.
- Sukartono, Suwardji, Mulyati, Baharuddin dan T. Wulan. 2014. Modifikasi aplikasi biomassa pada pertanaman ubikayu di tanah lempung berpasir (sandy loam) lahan kering Lombok Utara. Pusat penelitian dan pengembangan lahan kering Univ. Mataram.
- Sundari, T. 2010. Pengenalan varietas unggul dan teknik budidaya ubikayu. Balai Penelitian Kacang-kacangan dan Umbi-umbian. Malang. Indonesia.
- Suryana, A. 2007. Kebijakan penelitian dan pengembangan ubikayu untuk agroindustri dan ketahanan pangan. Hlm. 1-9. *dalam* Hernowo, D., Subandi dan N. Saleh. Prospek, Strategi, dan Teknologi Pengembangan Ubikayu untuk agroindustri dan ketahanan pangan. Pusat Penelitian dan Pengembangan Tanaman Pangan. Bogor.98p.
- Swanson. C.A. 1959. Translocation of organic solutes. Pages 481-545 *in* Plant physiology, vol. II. F.C. Stewart, cd. Academic Press. New York
- Taiz, L and E. Zeiger. 2006. *Plant physiology* 4th ed. Sinuer Associates, Sunderland, MA
- Taiz, L and E. Zeiger. 2006. *Plant physiology* 4th ed. Sinuer Associates, Sunderland, MA.
- Tan, S.L. and J.H. Cock, 1979. Branching habit as a yield determinant in cassava. *Field Crops Research* 2: 281-289
- Teerawanichpan, P., Lertpanyasampatha, M., Netrphan, S., Varavinit, S., Boonseng, O., and J. Narangajavana. 2008. Influence of cassava storage root development and environmental conditions on starch granule an size distribution. *Starch Starke* 60, 696-705.
- Thonre. G.N and A.F. Evans. 1964. Influence of tops and root on net assimilation rate of sugar beet and spinach beet and grafts between them. *Ann Bot.* 28: 499-508.
- Tongglum, A., C. Tirapon and S. Sinthuprama. 1987. Cassava cultural practices research in Thailand. *In*: R.H. Howeler and K. Kawano (Eds). Cassava Breeding and Agronomy Research in Asia. *Proc. 2nd Regional Workshop*, held in Rayong, Thailand. Oct 26-28, 1987. pp. 131-144.
- Tongglum, A., V. Vichukit, S. Jantawat, C. Sittibusaya, C. Tirapon, S. Sinthuprama, and R. H. Howeler. 1990. Recent progress in cassava agronomy research in Thailand. *In*: R.H. Howeler and K. Kawano (Eds). Cassava Breeding and Agronomy Research in Asia. *Proc. 2nd Regional Workshop*, held in Malang, Indonesia. Oct 22-27, 1990. pp. 199-223.
- Velez-Ramirez AI, van Ieperen W, Vreugdenhil D, van Poppel PM, Heuvelink E, and F.F.Millenaar. 2014. A single locus confers tolerance to continuous light and allows substantial yield increase in tomato. *Nature communications*, 5: DOI:10.1038/ncomms5549
- Veltkamp. H.J. 1985. Photosynthesis, transpiration, water use efficiency and leaf and mesophyll resistance of cassava as influenced by light intensity. *Agricultural university Wagenigen Papers* 85: 27-36



- Venkateswarlu. B. 1976. Source-sink interrelationships in lowland rice. *Plant Soil* 44(3):575-586
- Venkateswarlu. B. and. R.M.Visperas. 1987. Source-sink Relationship in crop plants. IRRI Research Paper Series. No. 125
- Viana, A.E.S., T. Sedyama, S.C. Lopes, C.S. sediyama, and V.SW. Rocha. 2000. Effect of length in stem cutting and its planting position on cassava yield. *Acta Scientiarum* 22(4): 1011-1015
- Villamayor, F. G., G. A. Dingal, F. A. Evangelo, J. C. Loder, A. C. Medellin, G. E. Sajise, and G. B. Burgos. 1992. Recent progress in cassava agronomy research in the Philippines, in: R.H. Howeler (Ed.), *Cassava breeding, agronomy and utilization research in Asia. Proc. 3rd Regional Workshop, held in Malang, Indonesia, 1992*, pp. 245-259
- Wahyuningsih, S dan T, Sundari. 2006. Evaluasi klon-klon harapan ubikayu untuk karakter hasil umbi dan pati. Balai Penelitian Kacang-kacangan dan Umbi-umbian. Malang. Indonesia.
- Wargiono, J., A. Hasanuddin, dan Suyamto. 2006. Teknologi Produksi Ubikayu Mendukung Industri Bioethanol. Puslitbangtan Bogor; 42 hlm
- Wartopo, A., Y. Efendi, dan Sukadi. 2009. Pengaturan jumlah cabang utama dan penjarangan buah terhadap hasil dan mutu benih tomat varietas kaliurang. *Jurnal Ilmu Pertanian* 5: 150-163
- Widodo, Y. 2009. Sinkronisasi aspek ekonomi dan ekologi dalam pengembangan ubikayu guna memenuhi kebutuhan pangan dan industri. Risalah Seminar Penelitian dan pengembangan tanaman pangan 2007-2008. Balai Penelitian dan Pengembangan Pertanian 342-355.
- Williams, C.N. and S.M. Ghazali, 1969. Growth and productivity of tapioca (*Manihot utilissima*). I. Leaf characteristics and yield. *Expl.Agric.* 5: 183-194.
- Williams,C.N. 1972. Growth and productivity of tapioca (*Manihot utilissima*) III; Crop ratio, spacing and yield. *Exp. Agric.* 10: 193-198.
- Williams, C.N.,1974. Growth and productivity of tapioca (*Manihot utilissima*). IV. Development and yield of tubers. *Expl. Agric*, 10: 9-16
- Woodward F.I. 1998. Do plants really need stomata. *Journal of Experimental Botany.* 49: 471-480.
- Wubs, A.M., Y. T. Ma., E. Heuvelink and L.F.M, Marcelis 2009. Genetic differences in fruit-set patterns are determined by differences in fruit sink strength and a source: sink threshold for fruit set. *Annals of Botany*,104: 957-964. DOI: 10.1093/aob/mcp181
- Yoo, C. Y., H.E., P. M. Hasegawa and M. V. Mickelbart. 2009. Regulation of Transpiration to Improve Crop Water Use. *Plant Science*, 26 (1): 410-425