

## TABLE OF CONTENTS

	Page.
<b>COVER PAGE</b> .....	i
<b>APPROVAL PAGE</b> .....	ii
<b>STATEMENT</b> .....	iii
<b>ACKNOWLEDGEENT</b> .....	iv
<b>TABLE OF CONTENTS</b> .....	v
<b>LIST OF TABLE</b> .....	vii
<b>LIST OF FIGURE</b> .....	viii
<b>INTISARI</b> .....	x
<b>ABSTRACT</b> .....	xi
<b>I. INTRODUCTON</b> .....	1
1.1. Background .....	1
1.2. Issues .....	2
1.3. Research Purposes.....	2
1.4. Advantages .....	2
1.5. Authenticity .....	3
<b>II. LITERATURE REVIEW AND THEORITICAL BASIS</b> .....	4
2.1. Literature Review .....	4
2.1.1. General N <sub>2</sub> O Emission .....	4
2.1.2. N <sub>2</sub> O Processes and Production in Paddy Soil .....	5
2.1.3. Cover Crops as Green Manure.....	8
2.1.4. Midseason Drainage: Water Management on Paddy Field .....	10
2.2. Theoritical Basis .....	12
<b>III. MATERIALS AND METHODS</b> .....	14
3.1. Site Description .....	14
3.2. Experimental Design .....	14
3.3. Research Procedures.....	14
3.3.1. Gas Investigation Method.....	15
3.3.2. Soil Properties Analysis Method .....	16
3.3.3. Plant Measurement .....	16
3.4. Statistical Analysis.....	17

<b>IV. RESUL AND DISCUSSION .....</b>	<b>18</b>
4.1. Result .....	18
4.1.1. Soil Characteristic in Paddy Field in Ehime, Japan.....	18
4.1.2. Soil Analysis in Paddy Field in Ehime, Japan .....	19
4.1.2.1. Soil Temperature in Paddy Field in Ehime, Japan .....	19
4.1.2.2. Soil Eh in Paddy Field in Ehime, Japan .....	20
4.1.2.3. Soil pH in Paddy Field in Ehime, Japan .....	22
4.1.2.4. Soil Water Content in Paddy Field in Ehime, Japan .....	22
4.1.2.5. Available of Soil NH <sub>4</sub> <sup>+</sup> in Paddy Field in Ehime, Japan.....	23
4.1.2.6. Available of Soil NO <sub>3</sub> <sup>-</sup> in Paddy Field in Ehime, Japan.....	24
4.1.3. Growth and Plant Biomass of Paddy in The Study Site .....	26
4.1.3.1. Plant Growth of Paddy in The Study Site .....	26
4.1.3.2. Rice Yield in The Study Site .....	26
4.1.4. N <sub>2</sub> O Emission and Global Warming Potential .....	27
4.1.4.1. Seasonal Variation of N <sub>2</sub> O Flux .....	27
4.1.4.2. Daily N <sub>2</sub> O Flux and N <sub>2</sub> O Emission .....	29
4.1.4.3. Global Warming Potential of N <sub>2</sub> O.....	30
4.2. Discussion.....	30
4.2.1. Soil properties related to N <sub>2</sub> O flux and N <sub>2</sub> O emission .....	30
4.2.2. Effect of midseason drainage on N <sub>2</sub> O emission.....	32
4.2.3. Effect of green manure application on N <sub>2</sub> O emission.....	33
4.2.4. Effect of midseason drainage and green manure application on GWP and Rice Yield.....	33
<b>V. CONCLUSION AND RECOMMENDATION .....</b>	<b>35</b>
5.1. Conclusion .....	35
5.2. Recommendation .....	35
<b>REFERENCES .....</b>	<b>36</b>
<b>SUPPLEMENTARY .....</b>	<b>41</b>

## LIST OF TABLE

	Page
Table 3.1 Combination of the treatment on paddy field in Ehime, Japan .....	14
Tabel 4.1 Soil Properties in paddy field in Ehime, Japan on 2014-2015 .....	18
Tabel 4.2 Plant height, tiller number, and chlorophyll content (SPAD) (Mean $\pm$ SD) in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan.....	26
Tabel 4.3 Number of panicle, 1000-grain weight, and brown rice yield (Mean $\pm$ SD) in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan.....	26
Tabel 4.4 Daily nitrous oxide (N <sub>2</sub> O) flux and emission (Mean $\pm$ SD) in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan. Early and late mean irrigating the field before and after MD, respectively .....	29

## LIST OF FIGURE

	Page
Figure 2.1 A schematic diagram of N transformations in a submerged soil (Buresh et al., 2008).....	6
Figure 3.1 Outline of the chamber and base used in the closed chamber method during fallow season.....	15
Figure 4.1 Soil temperature during fallow season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	19
Figure 4.2 Soil temperature during rice growing season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	20
Figure 4.3 Soil Eh during growing season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan.....	21
Figure 4.4 Soil Eh during midseason drainage period in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	21
Figure 4.5 Soil pH in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	22
Figure 4.6 Soil water content during fallow season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	23
Figure 4.7 Soil water content during rice growing season season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan.....	23
Figure 4.8 Available soil NH <sub>4</sub> <sup>+</sup> during fallow season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	24
Figure 4.9 Available soil NH <sub>4</sub> <sup>+</sup> during rice growing season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	24

Figure 4.10	Available soil NO <sub>3</sub> <sup>-</sup> during fallow season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	25
Figure 4.11	Available soil NO <sub>3</sub> <sup>-</sup> during rice growing season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	25
Figure 4.12	Seasonal variation of N <sub>2</sub> O flux during fallow season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan.....	27
Figure 4.13	Seasonal variation of N <sub>2</sub> O flux during rice growing season in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan.....	28
Figure 4.14	Seasonal variation of N <sub>2</sub> O flux during midseason drainage period in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	28
Figure 4.15	Global warming potential of N <sub>2</sub> O emission in the treatments of conventional midseason drainage (MD) without (CM) or with green manure (GM) (CGM) and prolonged MD without (PM) or with GM (PGM) in paddy field in Ehime, Japan .....	30