UNIVERSITAS GADJAH MADA
EKSPRESI GEN KETAHANAN Xa7 PADA PADI HITAM (Oryza sativa L.) SETELAH INFEKSI Xanthomonas oryzae pv oryzae
PUTRI WIJAYANTI, Dr. Yekti Asih Purwestri,S.Si.,M.Si; Dr. Tri Joko, S.P.,M.Sc.
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

DAFTAR PUSTAKA

Balai Penelitian Tanaman Pangan.2011.*Varietas Padi Tahun 1978 – 2010*. Bogor; Pustaka

Bai, J., Choi, S.H., Ponciano, G., Leung, H. and Leach, J.E. (2000) Xanthomonas oryzae pv. oryzae avirulence genes contribute differently and specifically to pathogen aggressiveness. Mol. Plant–Microbe Interact.13, 1322–1329.

Brenner WG, Schmulling T (2012) Transcript profiling of cytokinin action in Arabidopsis roots and shoots discovers largely similar but also organ-specific responses. BMC Plant Biol 12: 112.

Budiman. 2012. *Model Pertumbuhan Padi Hitam Pada Sawah Organik Di Kecamatan Kepanjen Kabupaten Malang*. TESIS. Jurusan Biologi, Universitas Brawijaya: Malang

Burdman, S., Shen, Y., Lee, S.W., Xue, Q., Ronald, P. 2004. RaxH/RaxR: a two-component regulatory system in *Xanthomonas oryzae pv. oryzae* required for AvrXa21 activity. *Mol Plant Microbe Interact.* 17:602-612

Chen, X., and Ronald, P.C. 2011. Innate immunity in rice. *Trends Plant Sci*. 16, 451-459. EPPO. 2007. *Xanthomonas oryzae*. European and Mediterranean Plant-Protection Organization. *Bulletin OEPP/EPPO*. 37: 543-553.

Dossa GS, Torres R, Henry A,Oliva R,Maiss E, Cruz CV,Wydra K. 2016.Rice response to simultaneous bacterial blight and drought stress during compatible and incompatible interactions.Eur J Plant Pathol

EPPO. 2007. *Xanthomonas oryzae*. European and Mediterranean Plant-Protection Organization. *Bulletin OEPP/EPPO*. 37: 543-553.

Fraga D, Meulia T, Fenster S. 2008. Real-time PCR. In: Current protocols essential laboratory techniques. New York (US): John Wiley & Sons, Inc. p. 10.3.1-10.3.33.

Furutani, A., Tsuge, S., Oku, T., Tsuno, K., Inoue, Y., Ochiai, H., Kaku, H. and Kubo, Y. (2003) Hpa1 secretion via type III secretion system in Xanthomonas oryzae pv. oryzae. J. Gen. Plant Pathol. 69, 271–275.

Gibbs RA. 1990. DNA amplification by the polymerase chain reaction. Anal Chem. 62:1202-1214

Herlina, L. dan Tiur, S.S. 2011. Seleksi Lapang Ketahanan Beberapa Varietas Padi terhadap Infeksi Hawar Daun Bakteri Strain IV dan VIII. *Buletin Plasma Nutfah*. Vol. 17. No. 2.

Hewajuli DA, Dharmayanti NLPI. 2014.Perkembangan Teknologi *Reverse Transcriptase-Polymerase Chain Reaction* dalam Mengidentifikasi Genom *Avian Influenza* dan *Newcastle Diseases*.Balai Besar Penelitian Veteriner;Bogor

Hifni, H.R., S. Mihardja, E. Soetarwo, Yusida, and M.K. Kardin. 1996. Penyakit hawar daun bakteri pada padi sawah masalah dan pemecahannya. Bulletin AgroBio. 1(1):18-23

UNIVERSITAS GADJAH MADA
EKSPRESI GEN KETAHANAN Xa7 PADA PADI HITAM (Oryza sativa L.) SETELAH INFEKSI Xanthomonas oryzae pv oryzae
PUTRI WIJAYANTI, Dr. Yekti Asih Purwestri,S.Si.,M.Si; Dr. Tri Joko, S.P.,M.Sc.
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

IRRI. 2014. Bacterial blight. http://www. knowledgebank.irri.org/ ricebreedingcourse/ breeding_for_resistence_blight [Maret 2017]

Khaeruni,A., Taufik, M., Wijayanto, T., Johan, E.A., 2014.Perkembangan penyakit hawar daun bakteri pada tiga varietas padi sawah yang diinokulasi pada beberapa fase pertumbuhan. Fitopatologi 10(4); 119-125

Kim, SM, Jung-Pil S, Yang Q, Tae-Hwan N, Russel FR, Kshirod K.J. 2015. Identification and fine-mapping of a new resistance gene, *Xa40*, conferring resistance to bacterial blight races in rice (*Oryza sativa* L.). *Theor Appl Genet* 128:1933–1943.

Kristamtini. 2009. Keragaan Beras Hitam sebagai Sumberdaya Genetik Lokal. *Prosiding Risalah Aplikasi Paket Teknologi "Mendukung Hari Pangan Sedunia*". BPTP Yogyakarta.

Kristamtini, Taryono, Panjisakti B, dan Rudi H.M. 2014. Keragaman Genetik dan Korelasi Parameter Warna Beras dan Kandungan Antosianin Total Sebelas Kultivar Padi Beras Hitam Lokal. *Ilmu Pertanian* Vol. 17 No.1, 2014 : 57 – 70.

Liu, D.N., P.C. Ronald, and A.J. Bogdanove. 2006. *Xanthomonas oryzae* pathovars: Model pathogens of a model crop. Mol. Plant Pathol. 7:303-324.

Li, Z., Alma, S., Angeles, E., Sukhwider, S., Jessica, D., Ning, H., and Gurdev, S. K. 2001. Are the dominant and recessive plant disease resistance genes similar? : A case study of rice R genes and *Xanthomonas oryzae* pv. *oryzae* races. *Genetics* 159: 757-765.

Loan, L.C., V.T.T. Ngan, and P.V. Du. 2006. Preliminary evaluation on resistance genes against rice bacterial leaf blight in Can Tho province-Vietnam. Omonrice.14:44–47.

Mostafa, K.M, Q. hassan and E. H. Chowdhury.2015.Application of remore sensors in mapping rice area and forecasting its production: A review. *Depparttemnet of Geomatics Engineering, Schulich School of Engineering University of Calgary.* 15: 769-791

Mudingotto PJ, Tusiime G, Asea G, Rubaihayo PR, Gibson P, Tumutegyereize J, & Lamo J. 2010. Genetics of resistance to bacterial leaf blight in rice germplasm in Uganda. *Second RUFORUM Biennial Meeting 20 - 24 September 2010.* Kampala (UG): Makerere University

Nam, H., Chung, B. C., et al. 2009. Combining tissue transcriptomics and urine metabolomics for breast cancer biomarker identification. Bioinformatics, 25(23), 3151–3157.

Nino-Liu, D., Pamela, C. Ronald and Adam J. B. 2006. *Xanthomonas Oryzae* Pathovars: Model Pathogens Of A Model Crop. *Molecular Plant Pathology*. 7(5) : 303–324.

Ogawa T, Yamamoto T. 1987.Near-isogenic lines as internasional differentials for resistance to bacterial blight of rice. [terhubung berkala]
http://www.gramene.org/newsletters/rice_genetic (Maret 2017)

UNIVERSITAS GADJAH MADA
EKSPRESI GEN KETAHANAN Xa7 PADA PADI HITAM (Oryza sativa L.) SETELAH INFEKSI Xanthomonas oryzae pv oryzae
PUTRI WIJAYANTI, Dr. Yekti Asih Purwestri,S.Si.,M.Si; Dr. Tri Joko, S.P.,M.Sc.
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Onasaya A *et al*.2010. Development of combined molecular diagnostic and DNA fingerprinting technique for rice bacteria patogens in Africa.Biotechnology 9:89-105

Ou, S.H. 1985. Rice Disease 2nd. Commonwealth Mycological Institute, Kiew, Surrey, England. 380 p.

Ponciano, G., Yoshikawa, M., Lee, J.L., Ronad , P.C., & Whalen, M.C. (2006). Pathogenesis-related gene expression in rice is correlated with developmentally controlled *Xa21*-mediated resistance against *Xanthomonas oryzae* pv. *oryzae*. *Physiological and Molecular Plant Pathology*. *69*,131-139.

Qiagen. 2012. *RNeassy Mini Handbook 4th Edition*. Qiagen Sample and Assay Technologies. Pp. 1.

Retnoningrum, 2017. Analisis Transkriptomik Gen Ketahanan *Xa* Padi Hitam (*Oryza sativa* L) setelah Infeksi *Xanthomonas oryzae* pv *oryzae*;UGM.Yogyakarta

Sa'adah, IR, Supriyanta, dan Subejo. 2013. Keragaman Warna Gabah Dan Warna Beras Varietas Lokal Padi Beras Hitam (*Oryza sativa* L.) Yang Dibudidayakan Oleh Petani Kabupaten Sleman, Bantul, Dan Magelang. *Vegetalika*. Vol.2 No.3, 2013 : 13-20.

Salanoubat, M., S. Genin, F. Artiguenave, J. Gouzy, S. Mangenot, M. Arlat, A. Billault, P. Brottier, J.C. Camus, L. Cattolico, M. Chandler, N. Choisne, C. Claudel-Renard, S. Cunnac, N. Demange, P. Siguier, P. Thebault, M. Whalen, P. Wincker, M. Levy, J. Weissenbach, and C.A. Boucher. 2002. Genome sequence of the plant pathogen Ralstonia solanacearum. Nature 415:497-502

Schornack, S., Meyer, A., Romer, P., Jordan, T., Lahaye, T. 2006. Gene for gene mediated recognition of nuclear target AvrBs3-like bacterial efector proteins. *J Plant Physiol*. 163:256-272.

Suparyono, Sudir, and Suprihanto. 2004. Pathotype profile of *Xanthomonas oryzae* pv. *oryzae* Isolates from the rice ecosystem in Java. Indonesian J. Agric. Sci. 5:63-69.

Sudir dan Handoko. 2012. Komposisi dan penyebaran patotipe *Xanthmonas oryzae* pv. *oryzae*, penyebab penyakit hawar daun bakteri padi di beberapa daerah produksi padi di Jawa Timur. J. Pengkaj. Dan Pengem. Tekn. Pert. 15(1):25-39

Swing, J.M., Van den Mouten, L. Vantening, B. Hoste, M. Gillis, T.W. Mew, and K. Kerters. 1990. Reclassificasion of the causal agents of bacterial blight (*Xanthomonas Campestris* pv. *oryzae*) and *Buletin Plasma N* 60 *utfah Vol.19 No.2 Th.2013* bacterial leaf streack (*Xanthomonas Campestris* pv. *oryzicola*) of rice as pathovar of *Xanthomonas oryzae* sp. (ex Ishyama, 1992) Rev. inf. J. Syst Bacterial 40:309-311.

Tasliah. 2012. Gen Ketahanan Tanaman Padi Terhadap Bakteri Hawar Daun (*Xanthomonas oryzae* pv. *oryzae*). *J. Litbang Pert.* Vol. 31 No. 3 Hal : 103-112.

UNIVERSITAS
GADJAH MADA
EKSPRESI GEN KETAHANAN Xa7 PADA PADI HITAM (Oryza sativa L.) SETELAH INFEKSI Xanthomonas oryzae pv oryzae
PUTRI WIJAYANTI, Dr. Yekti Asih Purwestri,S.Si.,M.Si; Dr. Tri Joko, S.P.,M.Sc.
Universitas Gadjah Mada, 2017 | Diunduh dari http://etd.repository.ugm.ac.id/

Theopillus BDM, Rapley R. 2002.PCR Mutation Detection Protocols. New Jersey: Humana-Pr

Tjitrosoepomo, G. 1994. *Taksonomi Tumbuhan (Spermatophyta)*. Gadjah Mada University Press : Yogyakarta.

Triny, S. K., Suryadi, Y., Sudir, Machmud, M. 2011. Penyakit Bakteri Padi dan Cara Pengendaliannya. Balai Besar Penelitian Tanaman Padi, Bogor.

Utami DW, Sutoro, Nurul H, Andari R, Ida H. 2011a. Keragaman genetik 96 aksesi plasma nutfah padi berdasarkan 30 marka SSR terpaut gen pengatur waktu pembungaan (HD *Genes*). *J AgroBiogen.* 7(2):76-84.

Utami DW, Kadir TS, Yuriyah S. 2011b. Faktor virulensi *AvrBs3/PthA* pada Ras III, Ras IV, Ras VIII, dan IXO93-068 patogen hawar daun bakteri (*Xanthomonas oryzae* pv. *oryzae*). *J AgroBiogen*. 7(1):1-8.

Venkatachalam P, Thulaseedharan A, Raghothama K (2007) Identification of expression profiles of tapping panel dryness (TPD) associated genes from the latex of rubber tree (Hevea brasiliensis Muell. Arg.). Planta 226: 499-515.

Vera-Cruz, C.-M., Bai, J., Ona, I., Leung, H., Nelson, R.-J., Mew, T.-W. and Leach, J.-E. (2000) Predicting durability of a disease resistance gene based on an assessment of the fitness loss and epidemiological consequences of avirulence gene mutation. Proc. Natl Acad. Sci. USA,97, 13500–13505

Wahyudi, A.T., Siti M., dan Abdjad A.N. 2011. *Xanthomonas oryzae* pv. *oryzae* Bakteri Penyebab Hawar Daun pada Padi : Isolasi, Karakterisasi, Telaah Mutagenesis dengan Transposon. *Makara, Sains*. Vol. 15. No.1 : 89-96.

Wang, Y.S., Pi, L.Y., Chen, X., Chakrabarty, P.K., Jiang, J., Lopez-De Leon, A., Liu, G.Z., Li, L., Benny, U., Oard, J., Ronald, P.C., & Song, W.Y. (2006). Rice XA21 binding protein 3 is a ubiquitin ligase required for full *Xa21*-mediated disease resistance. *The Plant Cell. 18*,3635-3646.

Winandari, A. Tjahjoleksono, dan D.W. Utami. 2014. Identifikasi gen ketahanan hawar daun bakteri pada galur padi introduksi dan galur dihaploid. J. Hama dan Penyakit Tumbuhan Tropika 14(2):101−109.

Yang, B. and F.F. White. 2004. Diverse members of the *AvrBs3/PthA* family of type III effectors are major virulence determinants in Bacterial Blight Disease of rice. J. Molecular Plant Microbe Interactions 17(11):1192-1200.

Yusida, S. Miharja, H.R. Hifni, dan T. Soewito. 1994. Identifikasi gen ketahanan pada varieties padi IRBBN yang efektif terhadap strain *Xanthomonas oryzae* pv. *oryzae* Kelompok III dan IV. hlm. 169- 174. *Dalam* M. Machmud dan Y. Suryadi (*eds*.) Risalah Hasil Penelitian Tanaman Pangan No. 3, Balai Penelitian Tanaman Pangan, Bogor

Yuwono T. 2006. Teori dan aplikasi *polymerase chain reaction*. Yogyakarta (Indonesia): Penerbit Andi