

- Akash, M.S.H., Rehman, K., Chen, S., 2013. Role of inflammatory mechanisms in pathogenesis of type 2 diabetes mellitus. *J. Cell. Biochem.* 114, 525–531.
- Alberti, G., 2005. Introduction to the metabolic syndrome. *Eur. Heart J. Suppl.* 7, D3–D5.
- Aldhahi, W., Hamdy, O., 2003. Adipokines, inflammation, and the endothelium in diabetes. *Curr. Diab. Rep.* 3, 293–298.
- Anandh Babu, P., Sabitha, K., Shyamaladevi, C., 2006. Green Tea Extract Impedes Dyslipidemia and Development of Cardiac Dysfunction in Streptozocin-Diabetic Rats. *Clin. Exp. Pharmacol. Physiol.* 33, 1184–1189.
- Aoyagi, R., Funakoshi-Tago, M., Fujiwara, Y., Tamura, H., 2014. Coffee Inhibits Adipocyte Differentiation via Inactivation of PPAR γ . *Biol. Pharm. Bull.* 37, 1820–1825.
- Atanasovska, E., Tasic, V., Slaninka-Miceska, M., Alabakovska, S., Zafirov, D., Kostova, E., Pavlovska, K., Filipce, V., Labacevski, N., 2014. Six week follow-up of metabolic effects induced by a high-fat diet and streptozotocin in a rodent model of type 2 diabetes mellitus. *Contrib. Sec Med Sci* 35, 169–79.
- Bansal, S., Syan, N., Mathur, P., Choudhary, S., 2012. Pharmacological profile of green tea and its polyphenols: a review. *Med. Chem. Res.* 21, 3347–3360. doi:10.1007/s00044-011-9800-4
- Bantle, J.P., 2009. Dietary Fructose and Metabolic Syndrome and Diabetes. *J. Nutr.* 139, 1263S–1268S.
- Bernal-Mizrachi, C., Weng, S., Feng, C., Finck, B.N., Knutsen, R.H., Leone, T.C., Coleman, T., Mechem, R.P., Kelly, D.P., Semenkovich, C.F., 2003. Dexamethasone induction of hypertension and diabetes is PPAR- α dependent in LDL receptor-null mice. *Nat. Med.* 9, 1069–1075.
- Bose, M., Lambert, J.D., Ju, J., Reuhl, K.R., Shapses, S.A., Yang, C.S., 2008. The major green tea polyphenol, (-)-epigallocatechin-3-gallate, inhibits obesity, metabolic syndrome, and fatty liver disease in high-fat-fed mice. *J. Nutr.* 138, 1677–1683.
- Brown, A.L., Lane, J., Coverly, J., Stocks, J., Jackson, S., Stephen, A., Bluck, L., Coward, A., Hendrickx, H., 2009. Effects of dietary supplementation with the green tea polyphenol epigallocatechin-3-gallate on insulin resistance and associated metabolic risk factors: randomized controlled trial. *Br. J. Nutr.* 101, 886–894.
- Cameron, A.J., Shaw, J.E., Zimmet, P.Z., 2004. The metabolic syndrome: prevalence in worldwide populations. *Endocrinol. Metab. Clin. North Am.* 33, 351–375.
- Capeau, J., 2008. Insulin resistance and steatosis in humans. *Diabetes Metab.* 34, 649–657.
- Chakravarthy, M.V., Pan, Z., Zhu, Y., Tordjman, K., Schneider, J.G., Coleman, T., Turk, J., Semenkovich, C.F., 2005. “New” hepatic fat activates PPAR α to maintain glucose, lipid, and cholesterol homeostasis. *Cell Metab.* 1, 309–322.
- Chan, D.C., Barrett, P.H.R., Watts, G.F., 2004. Lipoprotein Kinetics in the Metabolic Syndrome: Pathophysiological and Therapeutic Lessons from Stable Isotope Studies. *Clin. Biochem. Rev.* 25, 31–48.
- Chen, Y.-K., Cheung, C., Reuhl, K.R., Liu, A.B., Lee, M.-J., Lu, Y.-P., Yang, C.S., 2011. Effects of Green Tea Polyphenol (-)-Epigallocatechin-3-gallate on Newly Developed High-Fat/Western-Style Diet-Induced Obesity and Metabolic Syndrome in Mice. *J. Agric. Food Chem.* 59, 11862–11871.
- Chinetti, G., Lestavel, S., Bocher, V., Remaley, A.T., Neve, B., Torra, I.P., Teissier, E., Minnich, A., Jaye, M., Duverger, N., Brewer, H.B., Fruchart, J.C., Clavey, V., Staels, B., 2001. PPAR- α and PPAR- γ activators induce cholesterol

- removal from human macrophage foam cells through stimulation of the ABCA1 pathway. *Nat. Med.* 7, 53–58.
- Chirumbolo, S., 2015. In vivo anti-diabetic potential of chlorogenic acid as a consequence of synergism with other phenolic compounds? *Br. J. Nutr.* 113, 546–547.
- Cho, A.-S., Jeon, S.-M., Kim, M.-J., Yeo, J., Seo, K.-I., Choi, M.-S., Lee, M.-K., 2010. Chlorogenic acid exhibits anti-obesity property and improves lipid metabolism in high-fat diet-induced-obese mice. *Food Chem. Toxicol.* 48, 937–943.
- Choi, B.-K., Park, S.-B., Lee, D.-R., Lee, H.J., Jin, Y.-Y., Yang, S.H., Suh, J.-W., 2016. Green coffee bean extract improves obesity by decreasing body fat in high-fat diet-induced obese mice. *Asian Pac. J. Trop. Med.* 9, 635–643.
- Chou, C.-W., Huang, W.-J., Tien, L.-T., Wang, S.-J., 2007. (-)-Epigallocatechin gallate, the most active polyphenolic catechin in green tea, presynaptically facilitates Ca²⁺-dependent glutamate release via activation of protein kinase C in rat cerebral cortex. *Synap. N. Y. N* 61, 889–902.
- Clifford, M.N., 2000. Chlorogenic acids and other cinnamates—nature, occurrence, dietary burden, absorption and metabolism. *J. Sci. Food Agric.* 80, 1033–1043.
- Crozier, T.W.M., Stalmach, A., Lean, M.E.J., Crozier, A., 2012. Espresso coffees, caffeine and chlorogenic acid intake: potential health implications. *Food Funct.* 3, 30–33.
- Dandona, P., Aljada, A., Bandyopadhyay, A., 2004. Inflammation: the link between insulin resistance, obesity and diabetes. *Trends Immunol.* 25, 4–7.
- de Sotillo, D.V.R., Hadley, M., 2002. Chlorogenic acid modifies plasma and liver concentrations of: cholesterol, triacylglycerol, and minerals in (fa/fa) Zucker rats. *J. Nutr. Biochem.* 13, 717–726.
- Elbedoui, J., Oak, M., Anglard, P., Schinikerth, V., 2005. Catechins prevent vascular smooth muscle cell invasion by inhibiting MT1-MMP activity and MMP-2 expression. *Cardiovasc. Res.* 67, 317–325.
- Festing, M.F.W., 2006. Design and Statistical Methods in Studies Using Animal Models of Development. *ILAR J.* 47, 5–14.
- Furuyashiki, T., Nagayasu, H., Aoki, Y., Bessho, H., Hashimoto, T., Kanazawa, K., Ashida, H., 2004. Tea Catechin Suppresses Adipocyte Differentiation Accompanied by Down-regulation of PPAR γ 2 and C/EBP α in 3T3-L1 Cells. *Biosci. Biotechnol. Biochem.* 68, 2353–2359.
- Grundy, S.M., 2004. Definition of Metabolic Syndrome: Report of the National Heart, Lung, and Blood Institute/American Heart Association Conference on Scientific Issues Related to Definition. *Circulation* 109, 433–438.
- Han, S.H., Quon, M.J., Koh, K.K., 2005. Beneficial Vascular and Metabolic Effects of Peroxisome Proliferator-Activated Receptor- Activators. *Hypertension* 46, 1086–1092.
- Hardie, D.G., Hawley, S.A., Scott, J.W., 2006. AMP-activated protein kinase - development of the energy sensor concept: AMPK - development of the energy sensor concept. *J. Physiol.* 574, 7–15.
- Ihm, S.-H., Jang, S.-W., Kim, O.-R., Chang, K., Oak, M.-H., Lee, J.-O., Lim, D.-Y., Kim, J.-H., 2012. Decaffeinated green tea extract improves hypertension and insulin resistance in a rat model of metabolic syndrome. *Atherosclerosis* 224, 377–383.
- Iwabu, M., Yamauchi, T., Okada-Iwabu, M., Sato, K., Nakagawa, T., Funata, M., Yamaguchi, M., Namiki, S., Nakayama, R., Tabata, M., Ogata, H., Kubota, N., Takamoto, I., Hayashi, Y.K., Yamauchi, N., Waki, H., Fukayama, M., Nishino, I., Tokuyama, K., Ueki, K., Oike, Y., Ishii, S., Hirose, K., Shimizu, T., Touhara, K., Kadowaki, T., 2010. Adiponectin and AdipoR1 regulate PGC-1 α and mitochondria by Ca(2+) and AMPK/SIRT1. *Nature* 464, 1313–1319.



- Jin, S., Chang, C., Zhang, L., Liu, Y., Huang, X., Chen, Z., 2015. Chlorogenic Acid Improves Late Diabetes through Adiponectin Receptor Signaling Pathways in db/db Mice. *PLOS ONE* 10, e0120842.
- Kadowaki, T., Yamauchi, T., Kubota, N., Hara, K., Ueki, K., Tobe, K., 2006. Adiponectin and adiponectin receptors in insulin resistance, diabetes, and the metabolic syndrome. *J. Clin. Invest.* 116, 1784–1792.
- Karnieli, E., Hissin, P.J., Simpson, I.A., Salans, L.B., Cushman, S.W., 1981. A possible mechanism of insulin resistance in the rat adipose cell in streptozotocin-induced diabetes mellitus. Depletion of intracellular glucose transport systems. *J. Clin. Invest.* 68, 811.
- Kersten, S., Seydoux, J., Peters, J.M., Gonzalez, F.J., Desvergne, B., Wahli, W., 1999. Peroxisome proliferator-activated receptor α mediates the adaptive response to fasting. *J. Clin. Invest.* 103, 1489–1498.
- Khitan, Z., Kim, D.H., 2013. Fructose: A Key Factor in the Development of Metabolic Syndrome and Hypertension. *J. Nutr. Metab.* 2013, 1–12.
- Klaus, S., Pültz, S., Thöne-Reineke, C., Wolfram, S., 2005. Epigallocatechin gallate attenuates diet-induced obesity in mice by decreasing energy absorption and increasing fat oxidation. *Int. J. Obes.* 2005 29, 615–623.
- Kolovou, G.D., Anagnostopoulou, K.K., Cokkinos, D.V., 2005. Pathophysiology of dyslipidaemia in the metabolic syndrome. *Postgrad. Med. J.* 81, 358–366. doi:10.1136/pgmj.2004.025601
- Koo, S.-H., Satoh, H., Herzig, S., Lee, C.-H., Hedrick, S., Kulkarni, R., Evans, R.M., Olefsky, J., Montminy, M., 2004. PGC-1 promotes insulin resistance in liver through PPAR- α -dependent induction of TRB-3. *Nat. Med.* 10, 530–534.
- Kuate, D., Kengne, A.P.N., Biapa, C.P.N., Azantsa, B.G.K., Wan Muda, W.A.M.B., 2015. Tetrapleura tetraptera spice attenuates high-carbohydrate, high-fat diet-induced obese and type 2 diabetic rats with metabolic syndrome features. *Lipids Health Dis.* 14.
- Lafontan, M., Viguerie, N., 2006. Role of adipokines in the control of energy metabolism: focus on adiponectin. *Curr. Opin. Pharmacol.* 6, 580–585.
- Levine, T.B., Levine, A.B., 2012. Metabolic syndrome and cardiovascular disease. John Wiley & Sons.
- Lin, W.-Y., Xaiver Pi-Sunyer, F., Chen, C.-C., Davidson, L.E., Liu, C.-S., Li, T.-C., Wu, M.-F., Li, C.-I., Chen, W., Lin, C.-C., 2011. Coffee consumption is inversely associated with type 2 diabetes in Chinese. *Eur. J. Clin. Invest.* 41, 659–666.
- Li, R.W., Douglas, T.D., Maiyoh, G.K., Adeli, K., Theriault, A.G., 2006. Green tea leaf extract improves lipid and glucose homeostasis in a fructose-fed insulin-resistant hamster model. *J. Ethnopharmacol.* 104, 24–31.
- Manach, C., Scalbert, A., Morand, C., Rémésy, C., Jiménez, L., 2004. Polyphenols: food sources and bioavailability. *Am. J. Clin. Nutr.* 79, 727–747.
- Matsuda, M., Shimomura, I., 2013. Increased oxidative stress in obesity: Implications for metabolic syndrome, diabetes, hypertension, dyslipidemia, atherosclerosis, and cancer. *Obes. Res. Clin. Pract.* 7, e330–e341.
- Matsuzawa, Y., 2004. Adiponectin and Metabolic Syndrome. *Arterioscler. Thromb. Vasc. Biol.* 24, 29–33.
- Ma, Y., Gao, M., Liu, D., 2015. Chlorogenic Acid Improves High Fat Diet-Induced Hepatic Steatosis and Insulin Resistance in Mice. *Pharm. Res.* 32, 1200–1209.
- Meng, S., Cao, J., Feng, Q., Peng, J., Hu, Y., 2013. Roles of Chlorogenic Acid on Regulating Glucose and Lipids Metabolism: A Review. *Evid. Based Complement. Alternat. Med.* 2013, 1–11.



- Montero, M., Farah, A., Perrone, D., Trugo, L.C., Donangelo, C., 2007. Chlorogenic acid compounds from coffee are differentially absorbed and metabolized in humans. *J. Nutr.* 137, 2196–2201.
- Novo, S., Peritore, A., Guarneri, F.P., Corrado, E., Macaione, F., Evola, S., Novo, G., 2012. Metabolic syndrome (MetS) predicts cardio and cerebrovascular events in a twenty years follow-up. A prospective study. *Atherosclerosis* 223, 468–472.
- Okamoto, Y., Kihara, S., Funahashi, T., Matsuzawa, Y., Libby, P., 2006. Adiponectin: a key adipocytokine in metabolic syndrome. *Clin. Sci.* 110, 267–278. doi:10.1042/CS20050182
- Ong, K.W., Hsu, A., Tan, B.K.H., 2012. Chlorogenic acid stimulates glucose transport in skeletal muscle via AMPK activation: a contributor to the beneficial effects of coffee on diabetes. *PloS One* 7, e32718.
- Orešič, M., Vidal-Puig, A. (Eds.), 2014. A Systems Biology Approach to Study Metabolic Syndrome. Springer International Publishing, Cham.
- Petersen, K.F., Dufour, S., Savage, D.B., Bilz, S., Solomon, G., Yonemitsu, S., Cline, G.W., Befroy, D., Zeman, L., Kahn, B.B., others, 2007. The role of skeletal muscle insulin resistance in the pathogenesis of the metabolic syndrome. *Proc. Natl. Acad. Sci.* 104, 12587–12594.
- Plant, S., Shand, B., Elder, P., Scott, R., 2008. Adiponectin attenuates endothelial dysfunction induced by oxidised low-density lipoproteins. *Diabetes Vasc. Dis. Res. Off. J. Int. Soc. Diabetes Vasc. Dis.* 5, 102.
- Potenza, M.A., Marasciulo, F.L., Tarquinio, M., Tiravanti, E., Colantuono, G., Federici, A., Kim, J.-A., Quon, M.J., Montagnani, M., 2007. EGCG, a green tea polyphenol, improves endothelial function and insulin sensitivity, reduces blood pressure, and protects against myocardial I/R injury in SHR. *Am. J. Physiol. Endocrinol. Metab.* 292, E1378–1387.
- Riegsecker, S., Wiczynski, D., Kaplan, M.J., Ahmed, S., 2013. Potential benefits of green tea polyphenol EGCG in the prevention and treatment of vascular inflammation in rheumatoid arthritis. *Life Sci.* 93, 307–312.
- Ringseis, R., Muschick, A., Eder, K., 2007. Dietary Oxidized Fat Prevents Ethanol-Induced Triacylglycerol Accumulation and Increases Expression of PPAR α Target Genes in Rat Liver. *J. Nutr.* 137, 77–83.
- Rohdiana, D., Shabri, S., 2016. Analysis of individual catechin from green tea extracted and fractionated by column chromatography. *J. Penelit. Teh Dan Kina* 15.
- Ryo, M., Nakamura, T., Kihara, S., Kumada, M., Shibazaki, S., Takahashi, M., Nagai, M., Matsuzawa, Y., Funahashi, T., 2004. Adiponectin as a biomarker of the metabolic syndrome. *Circ. J.* 68, 975–981.
- Santamarina, A.B., Oliveira, J.L., Silva, F.P., Carnier, J., Mennitti, L.V., Santana, A.A., Souza, G.H.I. de, Ribeiro, E.B., Nascimento, C.M.O. do, Lira, F.S., Oyama, L.M., 2015. Green Tea Extract Rich in Epigallocatechin-3-Gallate Prevents Fatty Liver by AMPK Activation via LKB1 in Mice Fed a High-Fat Diet. *PLOS ONE* 10, e0141227.
- Sargowo, D., Andarini, S., 2011. Pengaruh Komposisi Asupan Makanan Terhadap Komponen Sindrom Metabolik. *J Kardiologi Indones* 32, 14–23.
- Serisier, S., Leray, V., Poudroux, W., Magot, T., Ouguerram, K., Nguyen, P., 2008. Effects of green tea on insulin sensitivity, lipid profile and expression of PPAR α and PPAR γ and their target genes in obese dogs. *Br. J. Nutr.* 99.
- Shah, A., Rader, D.J., Millar, J.S., 2010. The effect of PPAR- α agonism on apolipoprotein metabolism in humans. *Atherosclerosis* 210, 35–40.



- Shimoda, H., Seki, E., Aitani, M., 2006. Inhibitory effect of green coffee bean extract on fat accumulation and body weight gain in mice. *BMC Complement. Altern. Med.* 6, 9.
- Smith, U., Yang, X., 2011. Adipocytokine Dysregulation, Obesity and the Metabolic Syndrome. *Metab. Syndr.* 2, 229–244.
- Song, S.J., Choi, S., Park, T., 2014. Decaffeinated Green Coffee Bean Extract Attenuates Diet-Induced Obesity and Insulin Resistance in Mice. *Evid. Based Complement. Alternat. Med.* 2014, 1–14.
- Sukohar, A., Wirakusumah, F.F., Sastramihardja, H.S., others, 2013. Isolation and Characterization Cytotoxic Compounds Caffeine and Chlorogenic Acid Seeds of Lampung Coffee Robusta. *J. Med. Planta* 1.
- Sundaram, R., Naresh, R., Shanthi, P., Sachdanandam, P., 2013. Modulatory effect of green tea extract on hepatic key enzymes of glucose metabolism in streptozotocin and high fat diet induced diabetic rats. *Phytomedicine* 20, 577–584.
- Tang, F.-Y., Nguyen, N., Meydani, M., 2003. Green tea catechins inhibit VEGF-induced angiogenesis in vitro through suppression of VE-cadherin phosphorylation and inactivation of Akt molecule. *Int. J. Cancer* 106, 871–878.
- Taylor, R., 2012. Insulin Resistance and Type 2 Diabetes. *Diabetes* 61, 778–779.
- Tontonoz, P., Mangelsdorf, D.J., 2003. Liver X Receptor Signaling Pathways in Cardiovascular Disease. *Mol. Endocrinol.* 17, 985–993.
- Tsubakio-Yamamoto, K., Matsuura, F., Koseki, M., Oku, H., Sandoval, J.C., Inagaki, M., Nakatani, K., Nakaoka, H., Kawase, R., Yuasa-Kawase, M., Masuda, D., Ohama, T., Maeda, N., Nakagawa-Toyama, Y., Ishigami, M., Nishida, M., Kihara, S., Shimomura, I., Yamashita, S., 2008. Adiponectin prevents atherosclerosis by increasing cholesterol efflux from macrophages. *Biochem. Biophys. Res. Commun.* 375, 390–394.
- Van Dam, R.M., 2008. Coffee consumption and risk of type 2 diabetes, cardiovascular diseases, and cancer. *Appl. Physiol. Nutr. Metab. Physiol. Appliquée Nutr. Métabolisme* 33, 1269–1283.
- Volek, J.S., Feinman, R.D., 2005. Carbohydrate restriction improves the features of Metabolic Syndrome. Metabolic Syndrome may be defined by the response to carbohydrate restriction. *Nutr. Metab.* 2, 1.
- Wan, C.-W., Wong, C.N.-Y., Pin, W.-K., Wong, M.H.-Y., Kwok, C.-Y., Chan, R.Y.-K., Yu, P.H.-F., Chan, S.-W., 2013. Chlorogenic acid exhibits cholesterol lowering and fatty liver attenuating properties by up-regulating the gene expression of PPAR- α in hypercholesterolemic rats induced with a high-cholesterol diet. *Phytother. Res.* PTR 27, 545–551.
- Wang, Y., Ho, C.-T., 2009. Polyphenolic Chemistry of Tea and Coffee: A Century of Progress. *J. Agric. Food Chem.* 57, 8109–8114.
- Wild, S.H., Byrne, C.D., 2011. The Epidemiology of the Metabolic Syndrome and its Association with Diabetes, Cardiovascular Disease and Other Conditions. *Metab. Syndr.* 1.
- Wolfram, S., Raederstorff, D., Wang, Y., Teixeira, S.R., Elste, V., Weber, P., 2005. TEAVIGO (epigal vlocatechin gallate) supplementation prevents obesity in rodents by reducing adipose tissue mass. *Ann. Nutr. Metab.* 49, 54–63.
- Wu, L.-Y., Juan, C.-C., Hwang, L.S., Hsu, Y.-P., Ho, P.-H., Ho, L.-T., 2004. Green tea supplementation ameliorates insulin resistance and increases glucose transporter IV content in a fructose-fed rat model. *Eur. J. Nutr.* 43, 116–124.



UNIVERSITAS
GADJAH MADA

**EFEK KOMBINASI EKSTRAK KOPI HIJAU DAN EKSTRAK TEH HIJAU TERHADAP PROFIL
METABOLIK, ADIPONEKTIN, DAN**

EKSPRESI GEN PPAR-alfa PADA TIKUS MODEL SINDROMA METABOLIK

DWI ADI NUGROHO, dr. Arta Farmawati, PhD.; Dr. Pramudji Hastuti., M.S., Apt

Universitas Gadjah Mada, 2017 | Diunduh dari <http://etd.repository.ugm.ac.id/>

Yang, M.-H., Wang, C.-H., Chen, H.-L., 2001. Green, oolong and black tea extracts modulate lipid metabolism in hyperlipidemia rats fed high-sucrose diet. *J. Nutr. Biochem.* 12, 14–20.

Yan, J., Zhao, Y., Zhao, B., 2013. Green tea catechins prevent obesity through modulation of peroxisome proliferator-activated receptors. *Sci. China Life Sci.* 56, 804–810.

Zaveri, N.T., 2006. Green tea and its polyphenolic catechins: Medicinal uses in cancer and noncancer applications. *Life Sci.* 78, 2073–2080.