

DAFTAR PUSTAKA

- Abad CL, Kumar A, Safdar N. Antimicrobial therapy of sepsis and septic shock-when are two drugs better than one?. *Crit Care Clin* 2011;27:e1-27
- Akech SO, Karisa J, Nakamya P, Boga M, Maitland K. Phase II trial of isotonic fluid resuscitation in Kenyan children with severe malnutrition and hypovolaemia. *BMC Pediatrics* 2010;10:1-10
- Annane D. Corticosteroids for severe sepsis: an evidence based guide for physicians. *Annals of Intensive Care* 2011;1:7
- Annane D, Bellissant E, Bollaert PE, Briegel J, Keh D, Kupfer Y. Corticosteroids for severe sepsis and septic shock: a systematic review and meta-analysis. *BMJ* 2004;8:1-10
- Arabi YM., Dara SI, Memish Z, Al Abdulkareem A, Tamim HM, Kumar A, *et al.* Antimicrobial therapeutic determinants of outcomes from septic shock among patients with cirrhosis. *Hepatology* 2012;56:2305-15
- Ariano RE.2008.Acute respiratory distress syndrome: Pharmacotherapy Self Assessment. 6th edition. London: Blackwell Publishing
- Arifin N, Mangunatmadja I, Pudjiadi A, Pardede SO. Correlation between brain injury biomarkers and Glasgow coma scale in pediatric sepsis. *Paediatr Indones* 2012;52:111-7
- Assenzio B, Martin-Conte EL.2012.Cellular mechanisms of MOF during severe sepsis and septic shock-Understanding a serious killer. London: InTech Publishing
- Banks JG, Foulis AK, Ledingham I, Macsween RNM. Liver function in septic shock. *J Clin Pathol* 2002;35:1249-52
- Ben-Ari J, Samra Z, Nahum E, Levy I, Ashkenazi S, Schonfeld TM. Oral amphotericin B for the prevention of *Candida* bloodstream infection in critically ill children. *Pediatr Crit Care Med* 2006;7:115-8
- Bennett TD, Hayward KN, Farris RW, Ringold S, Wallace CA, Brogan TV. Very high serum ferritin levels are associated with increased mortality and critical care in pediatric patients. *Pediatr Crit Care* 2011;12:e233-6

- Biban P, Gaffuri M, Spaggiari S, Zaglia F, Serra A, Santuz P. Early recognition and management of septic shock in children. *Pediatric Reports* 2012;4:48-51
- Bigoniya P, Singh CS, Shukla A. A comprehensive review of different liver toxicants used in experimental pharmacology. *IJPSDR* 2009;1:124-35
- Branco RG, Garcia PCR, Piva JP, Casartelli CH, Seibel V, Tasker RC. Glucose level and risk of mortality in pediatric septic shock. *Ped Crit Care Med* 2005;6:470-2
- Branco RG, Tasker RC, Garcia PCR., Piva JP, Xavier LD. Glycemic control and insulin therapy in sepsis and critical illness. *Jornal de Pediatria* 2007;83:S128-36
- Brierley J, Carcillo JA, Choong K, Cornell T, De Caen A, Deymann A. Clinical practice parameters for hemodynamic support of pediatric and neonatal septic shock: 2007 update from the American College of Critical Care Medicine. *Crit Care Med* 2007;37:666-88
- Carcillo JA, Fields AI. Clinical practice parameters for hemodynamic support of pediatric and neonatal patients in septic shock. *Crit Care Med* 2002;30:1365-78
- [Carcillo JA](#), [Davis AL](#), [Zaritsky A](#). Role of early fluid resuscitation in pediatric septic shock. *JAMA* 1999;266:1242-5
- Cardoso LTQ, Grion CMC, Matsuo T, Anami EHT, Kauss IAM, Bonametti AM, *et al*. Impact of delayed admission to intensive care units on mortality of critically ill patients: a cohort study. *Crit Care* 2011;15:1-8
- Casartelli CH, Garcia PCR, Piva JP, Branco RG. Adrenal insufficiency in children with septic shock. *J Pediatr* 2003;79:S169-S76
- Ceneviva G, Paschall JA, Maffei F, Carcillo JA. Hemodynamic support in fluid refractory pediatric septic shock. *Pediatrics* 1998;102:1-6
- Chesnut RM, Ghajar J, Maas AIR, Marion DW, Servadei F, Zitney GA. Early indicators of prognosis in severe traumatic brain injury. *J Trauma* 1993;34:216-22
- Clec'h C, Ferriere F, Karoubi P, Fosse JP, Cupa M, Cohen Y, *et al*. Diagnostic and prognostic value of procalcitonin in patients with septic shock. *Crit Care Med* 2004;32:1166-9
- Deitch EA, Livingston DH, Lavery RF, Monaghan SF, Bongu A, Machiedo GW. Hormonally active women tolerate shock-trauma better than do men: a prospective study of over 4000 trauma patients. *Ann Surg* 2007;246:447-53

- Dellinger RP, Levy MM, Carlet JM, Bion J, Parker MM, Jaeschke R, *et al.* Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock. *Intensive Care Med* 2008;34:17-60
- Dellinger RP, Levy, MM, Rhodes A, Annane D, Gerlach H, Moreno R, *et al.* Surviving sepsis campaign: International guidelines for management of severe sepsis and septic shock. *Critical Care Medicine* 2012;41:580-637
- Dewi R. Sepsis pada anak: Pola kuman dan uji kepekaan. *Maj Kedokt Indon* 2011;61:101-6
- Dowd MD. Management of pediatric septic shock in the emergency department. *Pem-Database.Org* 2003;1;1-12
- Edwards J D, Houtrow AJ, Vasilevskis EE, Rehm RS, Markovitz BP, Dudley RA. Chronic conditions among children admitted to U.S. PICUs: their prevalence and impact on risk for mortality and prolonged length of stay. *Crit Care Med* 2012;40:2196–203
- Eidelman LA, Putterman D, Putterman C, Sprung CL. The spectrum of septic encephalopathy. Definitions, etiologies, and mortalities. *JAMA* 1996;275:470-3
- Enrionne MA, Powell KR. 2007. *Nelson Textbook of Pediatrics*. Edisi ke-18. Philadelphia: Saunders Elsevier
- Feudtner C, Christakis DA, Connell FA. Pediatric Deaths Attributable to Complex Chronic Conditions: A Population-Based Study of Washington State, 1980-1997. *Pediatrics* 2000; 106:205-9
- Finfer S, Chittock DR, Su SYS, Blair D, Foster D, Ronco JJ. Intensive versus conventional glucose control in critical ill patients. *N Engl J Med* 2009;360:1283-97
- Fioretto JR, Borin FC, Bonatto RC, Ricchetti SM, Kurokawa CS, De Moraes MA, *et al.* Procalcitonin in children with sepsis and septic shock. *J Pediatr Rio J* 2007;83:323-8
- Gaieski DF, Pines JM, Band RA, Mikkelsen ME, Massone R, Goyal M, *et al.* Impact of time to antibiotics on survival in patients with severe sepsis or septic shock in whom early goal-directed therapy was initiated in the emergency department. *Crit Care Med* 2010;38:1-9
- Garcia PCR, Longhi F, Branco RG, Piva JP, Lacks D, Tasker RC. Ferritin levels in children with severe sepsis and septic shock. *Acta Paediatrica* 2007;96:1829-31

- Ghuman AK, Newth CJL, Khemani RG. Impact of gender on sepsis mortality and severity of illness for prepubertal and postpubertal children. *J Pediatr* 2013;163:835-40
- Goldstein B, Giroir B, Randolph A. International pediatric sepsis consensus conference: Definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 2005.;6:1-8
- Guidet B, Martinet O, Boulain T, Philippart F, Poussel JP, Aken HV, *et al.* Assessment of hemodynamic efficacy and safety of 6% hydroxyethylstarch 130/0.4 vs. 0.9% NaCl fluid replacement in patients with severe sepsis: The CRYSTMAS study. *Crit Care* 2012;16:R94
- Han YY, Carcillo JA, Dragotta MA, Bills DM, Watson RS, Orr RA, *et al.* Early reversal of pediatric-neonatal septic shock by community physicians is associated with improved outcome. *Pediatrics* 2003;112:793-9
- Hazelzet JA, Risseuw-Appelt IM, Kornelisse RF, Hop WCJ, Dekker I, Hack CE, *et al.* Age-related differences in outcome and severity of DIC in children with septic shock and purpura. *Thrombosis and hemostasis* 1996;76:932-8
- He SJ, Jin YM, Huang AR, Wang CX, Zhou AH, Shan XO. Clinical analysis of community-acquired *Pseudomonas aeruginosa* septic shock. *Zhonghua Er Ke Za Zhi.* 2008;46:333-9
- Hollenberg SM. Inotrope and vasopressor therapy of septic shock. *Crit Care Med* 2009;25:781-802
- Honna L, Triratna S, Triwani, Theodorus. Use of pediatric logistic organ dysfunction (PELOD) in determining prognostic among pediatric intensive care unit patients. *Pediatr Indones* 2010;50:347-50
- [Inwald DP](#), [Tasker RC](#), [Peters MJ](#), [Nadel S](#). Emergency management of children with severe sepsis in the United Kingdom: the results of the Paediatric Intensive Care Society sepsis audit. *Arch Dis Child.*2009;94:348-53.
- Jaramillo-Bustamante JC, Marín-Agudelo A, Fernández-Laverde M., Bareño-Silva J. Epidemiology of sepsis in pediatric intensive care units: First Colombian multicenter study. *Pediatr Crit Care Med* 2012;13:501-8
- Jat KR, Jhamb U, Gupta VK. Serum lactate levels as the predictor of outcome in pediatric septic shock. *Indian Journal of Critical Care Medicine* 2011;15:102-7

- Keh D, Sprung CL. Use of corticosteroid therapy in patients with sepsis and septic shock: An evidence-based review. *Crit Care Med* 2004;32:S527-33
- Kim YA, Ha EJ, Jhang WK, Park SJ. Early blood lactate area as a prognostic marker in pediatric septic shock. *Intensive Care Med* 2013;39:1818-23
- Kissoon N. Sepsis and septic shock: A global perspective and initiative. *Saudi Med J* 2008;29:1383-7
- Kissoon N, Carcillo JA, Espinosa V, Argent A, Devictor D, Latour J, *et al.* World Federation of Pediatric Intensive Care and Critical Care Societies: Global sepsis initiative. *Pediatr Crit Care Med* 2011;12:494-503
- Kumar A, Roberts D, Wood KE, Light B, Parrillo JE, Cheang M, *et al.* Duration of hypotension before initiation of effective antimicrobial therapy is the critical determinant of survival in human septic shock. *Crit Care Med* 2006;34:1589-96
- Kumar A, Zarychanski R, Light B, Parillo JE, Maki D, Doucette S, *et al.* Early combination antibiotic therapy yields improved survival compared to monotherapy in septic shock: A propensity-matched analysis. *Crit Care Med* 2010;38:1-12
- Kutko MC, Calarco MP, Flaherty MB, Helmrich RF, Ushay HM, Pon S, Greenwald BM. Mortality rates in pediatric septic shock with and without multiple organ system failure. *Pediatr Crit Care Med* 2003;4:333-7
- Latief A, Pudjiadi AH, Somasetia DH, Alwy EH, Mulyo GD, Triratna S, *et al.* 2010. Diagnosis dan tatalaksana sepsis pada anak. Jakarta: Unit Kerja Koordinasi Pediatri Gawat Darurat Ikatan Dokter Anak Indonesia.
- Landry DW, Oliver JA. Mechanisms of disease. *New Engl J Med* 2001;345:588-95
- Larsen GY, Mechem N, Greenberg R. An emergency department septic shock protocol and care guideline for children initiated at triage. *Pediatrics* 2011;127:e1585-92
- Leteurtre S, Leclerc F, Duhamel A, Grandbastien B, Proulx F, Lacroix J, *et al.* Cumulative influence of organ dysfunctions and septic state on mortality of critically ill. *Am J Respir Critical Care Med* 2005;171:348-53
- Levi M, Toh CH, Tachil J, Watson HG. Guidelines for the diagnosis and management of disseminated intravascular coagulation. *Br J Haematol* 2009;145:24-33

- Lundberg JS, Perl TM, Wiblin T, Costigan MD, Dawson J, Wenzel RP, *et al.* Septic shock: An analysis of outcomes for patients with onset on hospital wards versus intensive care units. *Crit Care Med* 1998;26:1020-4
- Maat M, Buysse CMP, Emonts M, Spanjaard L, Joosten KFM, Hazelzet JA, *et al.* Improved survival of children with sepsis and purpura: effects of age, gender, and era. *Critical Care* 2007;11:R112
- Madiyono B, Moeslichan S, Sastroasmoro S, Budiman I, Purwanto HS. 2011. Dalam Sastroasmoro, S., dan Ismail S. Dasar-dasar Metodologi Penelitian Klinis. Edisi Keempat. Jakarta: CV Sagung Seto
- Melendez E, Bachur R. Advances in the emergency management of pediatric sepsis: *Curr Opin Pediatr* 2006;18:245-53
- Merenstein GB, Adams K, Weisman LE. 2002. Handbook of neonatal intensive care. Edisi ke-5. Philadelphia: Mosby
- Metta, D., Soebardja, D., Hudaya, D.S. 2006. The use of pediatric logistic organ dysfunction (PELOD) scoring system to determine the prognosis of patients in pediatric intensive care units: *Pediatrica Indonesiana* (46):1-6
- Minneci PC, Deans KJ, Eichacker PQ, Natanson C. The effects of steroids during sepsis depend on dose and severity of illness: an updated meta-analysis. *Clin Microbiol Infect* 2009;15: 308–18
- Muller L, Jaber S, Molinari N, Favier L, Larche J, Motte G *et al.* Fluid management and risk factors for renal dysfunction in patients with severe sepsis and/or septic shock. *Crit Care* 2012;16:R34
- Myburgh JA, Finfer S, Bellomo R, Billot L, Cass A, Webb SAR. Hydroxyethyl starch or saline for fluid resuscitation in intensive care. *N Engl J Med* 2012;367:1901–11
- Nadel S, Britto J, Booy R., Maconochie I, Habibi P, Levin M. Avoidable deficiencies in the delivery of health care to children with meningococcal disease. *J Accid Emerg Med* 1998;15:298-303
- Nduka OO. The pathophysiology of septic shock. *Crit Care Clin* 2009;25:677-702
- Oliveira CF, Nogueira de Sá FR, Oliveira DSF, Gottschald AFC, Moura JDG, Carcillo JA, *et al.* Time- and fluid-sensitive resuscitation for hemodynamic support of children in septic shock: Barriers to the implementation of the American College of Critical Care Medicine/Pediatric Advanced Life Support

- Guidelines in a Pediatric Intensive Care Unit in a developing world. *Pediatr Emerg Care* 2008;24:810-5
- Odetola F, Gebremariam A, Freed GL. Patient and hospital correlates of clinical outcomes and resource utilization in severe pediatric sepsis. *Pediatrics* 2007;119:487
- Perner A, Haase N, Guttormsen AB, Tenhunen J, Klemnzson G, Wetterslev J, *et al.* Hydroxyethyl starch 130/0.42 versus Ringer's acetate in severe sepsis. *N Engl J Med* 2012;367:124–34
- Pizarro CF, Troster EJ, Damiani D, Carcillo JA. Absolute and relative adrenal insufficiency in children with septic shock *Crit Care Med* 2005;33:885-9
- Po'voa PR, Carneiro AH, Ribeiro OS, Pereira AC. Influence of vasopressor agent in septic shock mortality. *Crit Care Med* 2009;37:410-6
- Rai R, Singh DK. Intravenous adrenaline for shock in neonates. *Indian Pediatrics* 2010;1:1-2.
- Ranjit S, Kissoon N, Jayakumar I. Aggressive management of dengue shock syndrome may decrease mortality rate. *Pediatr Crit Care Med* 2005;6:412-9
- Rosário C, Zandman-Goddard G, Meyron-Holtz EG, D'Cruz DP, Shoenfeld Y. The hyperferritinemic syndrome: macrophage activation syndrome, Still's disease, septic shock and catastrophic antiphospholipid syndrome. *BMC Medicine* 2013;11:185
- Rosenberg DI, Moss MM. 2004. Section on Critical Care and Committee on Hospital Care. Guidelines and Levels of Care for Pediatric Intensive Care Units: *Pediatrics* (114):1114-25
- Sakr Y, Reinhart K, Vincent JL. Does dopamine administration in shock influence outcome? Results of the Sepsis Occurrence in Acutely Ill Patients (SOAP) study. *Crit Care Med* 2006;34:589 –97
- Sarathi M, Lodha R, Vivekanandhan S, Arora NK. Adrenal status in children with septic shock using low-dose stimulation test. *Pediatr Crit Care Med* 2007;8:1-6
- Schlichting, D., Collam, J.S. 2007. Recognizing and managing severe sepsis: A common and deadly threat. *Southern Medical Journal* 2007;100:594-600
- Setiadi TS. Penatalaksanaan syok septik pada anak. Banjarmasin: Bagian Ilmu Kesehatan Anak RSUD Ulin.2005

- Suradi R, Siahaan CM, Boedjang RF, Sudiyanto, Setyaningsih I, Soedibjo S. 2002. Dalam Sastroasmoro, S., dan Ismail S. Dasar-dasar Metodologi Penelitian Klinis. Edisi Kedua. CV Sagung Seto, Jakarta.
- Shime N, Kawasaki T, Saito O, Akamine Y, Toda Y, Nakagawa S, *et al.* Incidence and risk factors for mortality in paediatric severe sepsis: results from the national paediatric intensive care registry in Japan: *Intensive Care Med* 2012;28:1191-7
- Smith WS, Sumnicht GE, Sharpe RW, Samuelson D, Millard FE. Granulocyte colony-stimulating factor versus placebo in addition to penicillin G in a randomized blinded study of gram-negative pneumonia sepsis: Analysis of survival and multisystem organ failure: *Blood* 1995;86:1301-9
- Sperry JL, Nathens AB, Frankel HL, Vanek SL, Moore EE, Maier RV, *et al.* Characterization of the gender dimorphism after injury and hemorrhagic shock: are hormonal differences responsible?. *Crit Care Med* 2008;36:1838-45
- Stoll BJ, Hansen N, Fanaroff AA, Wright LL. Changes in pathogens causing early-onset sepsis in very low birth weight infants. *New Engl J Med* 2002;347:240-7
- Tanurahardja AG, Pudjiadi A, Kaban R, Alatas H, Gayatri P, Pulungan A. Profil hormon tiroid pada anak dengan sepsis dan hubungan dengan skor PELOD sebagai parameter penentu prognosis pada sepsis. *International Journal of Pediatric Endocrinology* 2010;11:1-100
- Umegaki T, Ikai H, Imanaka Y. The impact of acute organ dysfunction on patients' mortality with severe sepsis. *J Anesthesiol Clin Pharmacol* 2011;27:180-4
- Unal D, Orhan O, Eroglu C, Kaplan B. Prealbumin is a more sensitive marker than albumin to assess the nutritional status in patients undergoing radiotherapy for head and neck cancer. *Wspolczesna Onkol* 2013; 17 (3): 276–280
- Wheeler DS, Basu RK. Pediatric shock: An overview. *The Open Pediatric Medicine Journal* 2013;7:2-9
- Winiarti E, Kosim MS, Supriatna M. Predicting PICU patient survival prognosis: Pediatric Logistic Organ Dysfunction vs Pediatric Index of Mortality scores. *Paediatr Indones* 2012;52:165-9
- Yokoyama Y, Schwacha MG, Samy TS, Bland KI, Chaudry IH. Gender dimorphism in immune responses following trauma and hemorrhage. *Immunol Res* 2003;26:63–76