

DAFTAR PUSTAKA

- ADA. (2015). *Standards of Medical Care in Diabetes 2015* (Vol. 38). Alexandria: American Diabetes Association.
- Aditama, W. (2011). Hubungan self-care, self-efficacy dan social support terhadap pengendalian kadar gula darah (HbA1c) penderita diabetes mellitus tipe 2 di puskesmas banyudono 1 dan ngeplak kabupaten boyolali provinsi jawa tengah. *Tesis*. Yogyakarta: Universitas Gadjah Mada.
- ADS. (2009). Australian Diabetes Society Position Statement : Individualization of HbA1c Targets for Adults with Diabetes Mellitus. (N. W. Cheung, J. J. Conn, M. C. D'Emden, J. E. Gunton, A. J. Jenkins, G. P. Ross, ... S. M. Twigg, Eds.). The Australian Diabetes Society.
- Alonso-morán, E., Satyrganova, A., Orueta, J. F., & Nuño-solinis, R. (2014). Prevalence of depression in adults with type 2 diabetes in the Basque Country : relationship with glycaemic control and health care costs. *BMC Public Health*, *14*(769), 1–8.
- Amir, N. (2012). Luaran (Outcome) Terapi pada Gangguan Depresi Mayor. *Cermin Dunia Kedokteran*, *39*(2), 92–96.
- APA. (1994). *Diagnostic and Statistical manual of Mental Disorder IV-Text Revision* (IV). Washington DC: American Psychiatric Association. <http://doi.org/10.1073/pnas.0703993104>
- Arisman. (2013). *Buku Ajar Ilmu Gizi (Obesitas, Diabetes Mellitus & Dislipidemia)*. Jakarta: EGC.
- Beck, A. T., Steer, R. A., & Brown, G. K. (n.d.). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Brown, A. F., Ettner, S. L., Piette, J., Weinberger, M., Gregg, E., Shapiro, F., ... Arbor, A. (2004). Socioeconomic Position and Health among Persons with Diabetes Mellitus : A Conceptual Framework and Review of the Literature, *26*, 63–77. <http://doi.org/10.1093/epirev/mxh002>
- Chacko, M., & Begum, K. (2016). Correlates of Glycemic Control - A Cross Sectional Study among Diabetic Population, (6), 1458–1465.
- Clare-Salzler, M. J., Crawford, J. M., & Kumar, V. (2007). Pankreas. In V. Kumar, R. S. Cotran, & S. L. Robbins (Eds.), *Buku Ajar Patologi* (7th ed., pp. 711–734). Jakarta: EGC.
- Dahlan, M. S. (2010). *Mendiagnosis dan Menatalaksana 13 Penyakit Statistik : Disertai Aplikasi Program Stata*. Jakarta: Sagung Set.
- de Groot, M., Anderson, R., Freedland, K. E., Clouse, R. E., & Lustman, P. J. (2001). Association of depression and diabetes complications: a meta-analysis. *Psychosomatic Medicine*, *63*(4), 619–630.
- Depkes. (2007). *Pharmaceutical Care Untuk Penderita Gangguan Depresif*.

Jakarta: Departemen Kesehatan Republik Indonesia.

- Diabetes UK. (2010). *Diabetes in the UK 2010: Key statistics on diabetes. Diabetes* (Vol. 692). <http://doi.org/http://www.diabetes.org.uk/Professionals/Publications-reports-and-resources/Reports-statistics-and-case-studies/Reports/Diabetes-prevalence-2010/>
- Dinkes Kabupaten Bantul. (2012). *Profil Kesehatan Kabupaten Bantul Tahun 2012*. Bantul: Dinas Kesehatan Kabupaten Bantul.
- Dinkes Kabupaten Bantul. (2013). *Profil Kesehatan Kabupaten Bantul Tahun 2013*. Bantul: Dinas Kesehatan Kabupaten Bantul.
- Dinkes Kabupaten Bantul. (2014). *Profil Kesehatan Kabupaten Bantul Tahun 2014*. Bantul: Dinas Kesehatan Kabupaten Bantul.
- FBOP. (2014). *Management of Major Depressive Disorder*. Federal Bureau of Prisons.
- Foster, D. W. (2014). Diabetes Mellitus. In K. J. Isselbacher, E. Braunwald, Jean D Wilson, J. B. Martin, A. S. Fauci, & D. L. Kasper (Eds.), *Harrison Prinsip-Prinsip Ilmu Penyakit Dalam* (13th ed., pp. 2196–2217). Jakarta: EGC.
- Ghazali, M. V., Sastromiharjo, S., Soedjarwo, S. R., Soelaryo, T., & Pramulyo, H. S. (2010). Studi cross-sectional. In S. Sastroasmoro & S. Ismael (Eds.), *Dasar-Dasar Metodologi Penelitian Klinis* (3rd ed., pp. 112–126). Jakarta: Sagung Seto.
- Goto, M., Goto, A., Ikeda, N., Noda, H., & Shibuya, K. (2015). Factors Associated with Untreated Diabetes: Analysis of Data from 20 , 496 Participants in the Japanese National Health and Nutrition Survey, 748, 1–8. <http://doi.org/10.1371/journal.pone.0118749>
- Hoggatt, K. J. (2002). *Biostatistical Methods in Epidemiology*. New York: John Wiley & Sons Inc.
- Huang, N., Duggan, K., & Harman, J. (2008). Lifestyle management of hypertension. *Australian Prescriber*, 31(6), 150–153.
- IDF. (2012). *Global Guideline for Type 2 Diabetes*. Brussels: International Diabetes Federation.
- Ilyas, E. I. (2013). Manfaat Latihan Jasmani Bagi Penyandang Diabetes. In S. Soegondo, P. Soewondo, & I. Subekti (Eds.), *Penatalaksanaan Diabetes Melitus Terpadu* (2nd ed., pp. 289–301). Jakarta: FKUI.
- Jiwo, T. (2012). *Depresi: Panduan bagi pasien, keluarga dan teman dekat* (1st ed.). Purworejo: Pusat Pemulihan dan Pelatihan Bagi Penderita Gangguan Jiwa.
- Juarez, D. T., Sentell, T., Tokumar, S., Goo, R., Davis, J. W., & Mau, M. M. (2012). Factors associated with poor glycemic control or wide glycemic variability among diabetes patients in Hawaii, 2006–2009. *Preventing*

Chronic Disease, 9, 120065. <http://doi.org/10.5888/pcd9.120065>

- Kabo, P. (2010). *Bagaimana menggunakan obat-obat kardiovaskular secara rasional*. Jakarta: FKUI.
- Kaplan, H. I., Sadock, B. J., & Grebb, J. A. (2010). *Sinopsis Psikiatri*. Ciputat: Binarupa Aksara.
- Katz, M. H. (2006). *Multivariable Analysis: A Practical Guide for Clinicians Second Edition* (2nd ed.). Cambridge: Cambridge University Press. <http://doi.org/10.1017/CBO9781107415324.004>
- Kemendes. (2013). *Riset Kesehatan Dasar Tahun 2013*. Jakarta: Badan Pengembangan dan Penelitian Kesehatan Kementerian Kesehatan RI.
- KEMENKES RI. (2013). *Riset Kesehatan Dasar*. Jakarta: Kementerian Kesehatan Republik Indonesia.
- Khattab, M., Khader, Y. S., Al-Khawaldeh, A., & Ajlouni, K. (2010). Factors associated with poor glycemic control among patients with type 2 diabetes. *Journal of Diabetes and Its Complications*, 24(2), 84–9. <http://doi.org/10.1016/j.jdiacomp.2008.12.008>
- Lee, H., Ahn, S., & Kim, Y. (2009). Self-care, Self-efficacy, and Glycemic Control of Koreans With Diabetes Mellitus. *Asian Nursing Research*, 3(3), 139–146. [http://doi.org/10.1016/S1976-1317\(09\)60025-6](http://doi.org/10.1016/S1976-1317(09)60025-6)
- Lemeshow, S., Jr, D. W. H., Klar, J., & Lwanga, S. K. (1990). Sample Size for Case-Control Studies. In S. Lemeshow, D. W. H. Jr, J. Klar, & S. K. Lwanga (Eds.), *Adequacy of Sample Size in Health Studies* (pp. 16–20). Chichester: John Wiley & Sons.
- Lind, M., Odén, A., Fahlén, M., & Eliasson, B. (2008). A systematic review of HbA1c variables used in the study of diabetic complications. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, 2(4), 282–293. <http://doi.org/10.1016/j.dsx.2008.04.006>
- Lunghi, C., Moisan, J., & Gre, J. (2016). Incidence of Depression and Associated Factors in Patients With Type 2 Diabetes in Quebec , Canada, 95(21), 1–10. <http://doi.org/10.1097/MD.0000000000003514>
- Mackay, J., & Mensah, G. A. (2004). *The Atlas of Heart Disease and Stroke*. Geneva: WHO & CDC.
- Mihardja, L. (2009). Faktor yang Berhubungan dengan Pengendalian Gula Darah pada Penderita Diabetes Mellitus di Perkotaan Indonesia. *Majalah Kedokteran Indonesia*, 59(September), 418–24.
- Misra, R., & Lager, J. (2009). Ethnic and gender differences in psychosocial factors, glycemic control, and quality of life among adult type 2 diabetic patients. *Journal of Diabetes and Its Complications*, 23(1), 54–64. <http://doi.org/10.1016/j.jdiacomp.2007.11.003>
- Murti, B. (2003). *Prinsip dan Metode Riset* (2nd ed.). Yogyakarta: Gadjah Mada University Press.

- NIMH. (2009). *Depression and diabetes. Psychiatry* (Vol. 8). National Institute of Mental Health. <http://doi.org/10.1016/j.mppsy.2009.03.009>
- Nyamu, P. N., Otieno, C. F., Amayo, E. O., & Mcligeyo, S. O. (2003). Risk factors and prevalence of diabetic foot ulcers at Kenyatta National Hospital, Nairobi. *East African Medical Journal*, 80(1). <http://doi.org/10.4314/eamj.v80i1.8664>
- Nyunt, S. W., Howteerakul, N., Suwannapong, N., & Rajatanun, T. (2010). Self-efficacy, self-care behaviors and glycemic control among type-2 diabetes patients attending two private clinics in Yangon, Myanmar. *Southeast Asian Journal of Tropical Medicine and Public Health*, 41(4), 943–951.
- NZGG. (2003). *Guideline Management of Type 2. Review Literature And Arts Of The Americas*. Wellington: New Zealand Guidelines Group.
- O'Brien, R. M. (2007). A caution regarding rules of thumb for variance inflation factors. *Quality and Quantity*, 41(5), 673–690. <http://doi.org/10.1007/s11135-006-9018-6>
- Ozcelik, F., Yiginer, O., Arslan, E., Serdar, M. A., Uz, O., Kardesoglu, E., & Kurt, I. (2010). Association between glycemic control and the level of knowledge and disease awareness in type 2 diabetic patients. *Polskie Archiwum Medycyny Wewnetrznej*, 120(10), 399–406.
- Papelbaum, M., Moreira, R. O., Coutinho, W., Kupfer, R., Zagury, L., Freitas, S., & Appolinário, J. C. (2011). Depression, glycemic control and type 2 diabetes. *Diabetology & Metabolic Syndrome*, 3(1), 26. <http://doi.org/10.1186/1758-5996-3-26>
- PERKENI. (2011). *Konsensus Pengendalian dan Pencegahan Diabetes Mellitus Tipe 2 di Indonesia 2011*. Perkumpulan Endokrinologi Indonesia.
- Phaswana-Mafuya, N., Peltzer, K., Chirinda, W., Musekiwa, A., & Kose, Z. (2013). Sociodemographic predictors of multiple non-communicable disease risk factors among older adults in South Africa. *Global Health Action*, 6(1), 1–8. <http://doi.org/10.3402/gha.v6i0.20680>
- Powers, A. C. (2012). Diabetes Mellitus. In D. L. Longo, D. L. Kasper, J. L. Jameson, A. S. Fauci, S. L. Hauser, & J. Loscalzo (Eds.), *Harrison's Principles of Internal Medicine* (18th ed., pp. 2968– 3002). New York: McGraw-Hill Medical.
- Purwanti, O. S. (2013). Analisis Faktor - Faktor Risiko Terjadi Ulkus Kaki pada Pasien Diabetes Mellitus di RSUD Dr. Moewardi. *Tesis*. Depok: Universitas Indonesia.
- Roy, T., & Lloyd, C. E. (2012). Epidemiology of depression and diabetes : A systematic review. *Journal of Affective Disorders*, 142, S8–S21. [http://doi.org/10.1016/S0165-0327\(12\)70004-6](http://doi.org/10.1016/S0165-0327(12)70004-6)
- Schteingrat, D. E. (2006). Pankreas : Metabolisme Glukosa dan Diabetes Mellitus. In S. A. Price & L. M. Wilson (Eds.), *Patofisiologi Konsep Klinis Proses-Proses Penyakit* (6th ed., pp. 1259–1274). Jakarta: EGC.

- Segler, C. (2009). Diabetes, Smoking, and Foot Problems: Is It All Related? Retrieved January 1, 2015, from <http://www.articlesbase.com/health-articles/diabetes-smoking-and-foot-problems-is-it-all-related-976255.html>
- Semiardji, G. (2013). Stres Emosional pada Penyandang Diabetes. In S. Soegondo, P. Soewondo, & I. Subekti (Eds.), *Penatalaksanaan Diabetes Melitus Terpadu* (2nd ed., pp. 337–347). Jakarta: FKUI.
- Siddiqui, F. J., Avan, B. I., Mahmud, S., Nanan, D. J., Jabbar, A., & Assam, P. N. (2015). Uncontrolled diabetes mellitus: Prevalence and risk factors among people with type 2 diabetes mellitus in an Urban District of Karachi, Pakistan. *Diabetes Research and Clinical Practice*, *107*(1), 148–156. <http://doi.org/10.1016/j.diabres.2014.09.025>
- Soegondo, S. (2013). Diagnosis dan Klasifikasi Diabetes Meliitus Terkini. In S. Soegondo, P. Soewondo, & I. Subekti (Eds.), *Penatalaksanaan Diabetes Melitus Terpadu* (2nd ed., pp. 19–30). Jakarta: FKUI.
- Swedish National Institute of Public Health. (2010). *Physical Activity in the Prevention and Treatment of Disease. Medicine & Science in Sports & Exercise*. Östersund: Professional Associations for Physical Activity. Retrieved from http://journals.lww.com/acsm-msse/Abstract/1999/11001/Physical_activity_in_the_prevention_and_treatment.22.aspx
- Tridjaja, B. (2009). *Konsensus Nasional Pengelolaan Diabetes Melitus Tipe 1* (2nd ed.). Jakarta: Ikatan Dokter Anak Indonesia.
- Trief, P. M., Teresi, J. a., Eimicke, J. P., Shea, S., & Weinstock, R. S. (2008). Improvement in diabetes self-efficacy and glycaemic control using telemedicine in a sample of older, ethnically diverse individuals who have diabetes: the IDEATel project. *Age and Ageing*, *38*(2), 219–225. <http://doi.org/10.1093/ageing/afn299>
- Tumbelaka, A. R., Riono, P., Sastroasmoro, S., Wirjodiarjo, M., Pudjiastuti, P., & Firman, K. (2010). Pemilihan uji hipotesis. In S. Sastroasmoro & S. Ismael (Eds.), *Dasar-dasaar Metodologi Penelitian Klinis* (3rd ed., pp. 279–300). Jakarta: Sagung Seto.
- Vogelzangs, N., Kritchevsky, S. B., Beekman, A. T. F., Newman, A. B., Satterfield, S., Simonsick, E. M., ... Penninx, B. W. J. H. (2008). Depressive symptoms and change in abdominal obesity in older persons. *Archives of General Psychiatry*, *65*(12), 1386–93. <http://doi.org/10.1001/archpsyc.65.12.1386>
- Walker, R. J., Gebregziabher, M., Martin-harris, B., & Egede, L. E. (2014). Relationship between social determinants of health and processes and outcomes in adults with type 2 diabetes : validation of a conceptual framework, *14*(1), 1–10. <http://doi.org/10.1186/1472-6823-14-82>
- WHO. (1992). *The ICD-10 Classification of Mental and Behavioural Disorders. International Classification* (Vol. 10). Geneva: World Health Organization. [http://doi.org/10.1002/1520-6505\(2000\)9:5<201::AID-EVAN2>3.3.CO;2-P](http://doi.org/10.1002/1520-6505(2000)9:5<201::AID-EVAN2>3.3.CO;2-P)

- WHO. (1994). *Prevention of Diabetes Mellitus: Report of a WHO Study Group. World Health Organization technical report series* (Vol. 844). Geneva: World Health Organization. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7941615>
- WHO. (2003). Depression. In *Disease-Specific Reviews : Section III* (pp. 87–92). Geneva: World Health Organization. Retrieved from http://www.who.int/chp/knowledge/publications/adherence_section3.pdf
- WHO. (2010). *Global recommendations on physical activity for health*. Geneva: World Health Organization. <http://doi.org/10.1080/11026480410034349>
- WHO, S. (2011). *Noncommunicable Diseases in the South-East Asia Region*. New Delhi: World Health Organization, Regional Office for South-East Asia. <http://doi.org/978-92-9022-413-6>
- Wu, W. (2010). Confounding , interaction , and mediation in multivariable / multivariate regression modeling logo. *Cancer Biostatistics*.
- Yosep, H. I., & Sutini, T. (2014). *Buku Ajar Keperawatan Jiwa* (6th ed.). Bandung: Refika Aditama.